

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
5 August 2004 (05.08.2004)

PCT

(10) International Publication Number  
**WO 2004/065561 A2**

- (51) International Patent Classification<sup>7</sup>: **C12N**
- (21) International Application Number:  
PCT/US2004/001643
- (22) International Filing Date: 21 January 2004 (21.01.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/441,637 21 January 2003 (21.01.2003) US
- (71) Applicant (for all designated States except US): **PTC THERAPEUTICS, INC.** [US/US]; 100 Corporate Court, South Plainfield, NJ 07080 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **CAO, Liangxian** [CN/US]; 6 Mioduski Court, Parlin, NJ 08859 (US). **TRIFILLIS, Panayiota** [US/US]; 309 Rivercrest Drive, Piscataway, NJ 08854 (US).
- (74) Agents: **CORUZZI, Laura, A.** et al.; Jones Day, 222 East 41st Street, New York, NY 10017-6702 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

[Continued on next page]

(54) Title: METHODS FOR IDENTIFYING COMPOUNDS THAT MODULATE UNTRANSLATED REGION-DEPENDENT GENE EXPRESSION AND METHODS OF USING SAME



(57) Abstract: The present invention relates to methods for identifying compounds that modulate untranslated region-dependent expression of a target gene. The invention particularly relates to using untranslated regions of a target gene or fragments thereof linked to a reporter gene to identify compounds that modulate untranslated region-dependent expression of a target gene. The methods of the present invention provide a simple, sensitive assay for high-throughput screening of libraries of compounds to identify pharmaceutical leads.



---

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**METHODS FOR IDENTIFYING COMPOUNDS  
THAT MODULATE UNTRANSLATED REGION-DEPENDENT GENE  
EXPRESSION AND METHODS OF USING SAME**

---

[0001] This application is entitled to and claims priority benefit to U.S. provisional application Serial No. 60/441,637, filed January 21, 2003, which is incorporated herein by reference in its entirety.

**1. INTRODUCTION**

[0002] The present invention relates to a method for screening and identifying compounds that modulate untranslated region-dependent expression of any gene. In particular, the invention provides reporter gene-based assays for the identification of compounds that modulate untranslated region-dependent expression of a gene. The methods of the present invention provide a simple, sensitive assay for high-throughput screening of libraries of compounds to identify pharmaceutical leads.

**2. BACKGROUND OF THE INVENTION**

**2.1. Gene Expression**

[0003] Every living organism is a product of expression of its genes in response to a developmental program (encoded in the genome itself) and environmental factors. Gene expression can be defined as the conversion of the nucleotide sequence of a gene into the amino acid sequence of a protein or into the nucleotide sequence of a stable RNA.

[0004] In eukaryotes, gene expression begins in the nucleus with the transcription of a gene into a premessenger-RNA, also referred to as a primary transcript. While still in the nucleus, the pre-mRNA is extensively modified. Each primary transcript is capped at the 5' end, associates with hnRNP proteins to form messenger RNA-protein particles ("mRNPs"), acquires a polyadenylic acid tail at the 3' end, and undergoes splicing to remove introns. In addition, the nucleotide sequence of certain pre-mRNAs can be altered post-transcriptionally in a process known as RNA editing. Thus processed, the mature mRNA is exported to the cytoplasm. Upon export, mRNA dissociates from hnRNP proteins and binds a set of cytosol-specific mRNA-binding proteins. Once in the cytoplasm, the mRNA either immediately associates with ribosomes and templates for protein synthesis or is localized to discrete cellular foci to direct compartment-specific protein synthesis. Degradation of mRNA and protein, which occurs both in the nucleus and the cytoplasm, concludes the list of processes that comprise gene expression.

## **2.2. Post-transcriptional Gene Expression Regulation**

[0005] Gene expression is very tightly regulated. To produce the desired phenotype, each gene must be expressed at a defined time and at a defined rate and amount. Extensive experimental evidence indicates that post-transcriptional processes such as mRNA decay, translation, and mRNA localization constitute major control points in gene expression.

[0006] An aberration in the expression of one or more genes can be the cause or a downstream effect of a disease or other abnormality. Understanding gene expression regulation mechanisms in the normal/healthy/wild-type cell/body and during pathology will permit rational therapeutic intervention.

[0007] Regulation of gene expression both at the mRNA stability and translation levels is important in cellular responses to development or environmental stimuli such as nutrient levels, cytokines, hormones, and temperature shifts, as well as environmental stresses like hypoxia, hypocalcemia, viral infection, and tissue injury (reviewed in Guhaniyogi & Brewer, 2001, *Gene* 265(1-2):11-23). Furthermore, alterations in mRNA stability have been causally connected to specific disorders, such as neoplasia, thalassemia, and Alzheimer's disease, (reviewed in Guhaniyogi & Brewer, 2001, *Gene* 265(1-2):11-23 and *Translational Control of Gene Expression*, Sonenberg, Hershey, and Mathews, eds., 2000, CSHL Press). In contrast, regulation of gene expression at the mRNA localization level is primarily used by the cell to create and maintain polarity (internal gradients of protein concentration) (reviewed in *Translational Control of Gene Expression*, Sonenberg, Hershey, and Mathews, eds., 2000, CSHL Press).

## **2.3. mRNA Untranslated Regions in Gene Expression Regulation**

[0008] A typical mRNA contains a 5' cap, a 5' untranslated region ("5' UTR") upstream of a start codon, an open reading frame, also referred to as coding sequence, that encodes a stable RNA or a functional protein, a 3' untranslated region ("3' UTR") downstream of the termination codon, and a poly(A) tail. Most studied *cis*-dependent RNA-based gene expression regulation elements map to the 5' or 3' UTRs.

[0009] Examples of 5' UTR regulatory elements include the iron response element ("IRE"), internal ribosome entry site ("IRES"), upstream open reading frame ("uORF"), male specific lethal element ("MSL-2"), G-quartet element, and 5'-terminal oligopyrimidine tract ("TOP") (reviewed in Keene & Tenenbaum, 2002, *Mol Cell* 9:1161 and *Translational Control of Gene Expression*, Sonenberg, Hershey, and Mathews, eds., 2000, CSHL Press).

[0010] Examples of 3' UTR regulatory elements include AU-rich elements ("AREs"), Selenocysteine insertion sequence ("SECIS"), histone stem loop, cytoplasmic



polyadenylation elements ("CPEs"), nanos translational control element, amyloid precursor protein element ("APP"), translational regulation element ("TGE")/direct repeat element ("DRE"), bruno element ("BRE"), 15-lipoxygenase differentiation control element (15-LOX-DICE), and G-quartet element (reviewed in Keene & Tenenbaum, 2002, Mol Cell 9:1161).

**[0011]** The internal ribosome entry site ("IRES") is one of the 5' UTR-based *cis*-acting elements of post-transcriptional gene expression control. IRESes facilitate cap-independent translation initiation by recruiting ribosomes directly to the mRNA start codon. IRESes are commonly located in the 3' region of a 5' UTR and are, as recent work has established, frequently composed of several discrete sequences. IRESes do not share significant primary structure homology, but do form distinct RNA secondary and tertiary structures. Some IRESes contain sequences complementary to 18S RNA and therefore may form stable complexes with the 40S ribosomal subunit and initiate assembly of translationally competent complexes. A classic example of an "RNA-only" IRES is the internal ribosome entry site from Hepatitis C virus. However, most known IRESes require protein co-factors for activity. More than 10 IRES *trans*-acting factors ("ITAFs") have been identified so far. In addition, all canonical translation initiation factors, with the sole exception of 5' end cap-binding eIF4E, have been shown to participate in IRES-mediated translation initiation (reviewed in Vagner et al., 2001, EMBO reports 2:893 and Translational Control of Gene Expression, Sonenberg, Hershey, and Mathews, eds., 2000, CSHL Press).

**[0012]** AU-rich elements ("AREs") are 3' UTR-based regulatory signals. AREs are the primary determinant of mRNA stability and one of the key determinants of mRNA translation initiation efficiency. A typical ARE is 50 to 150 nucleotides long and contains 3 to 6 copies of AU<sub>3</sub>A pentamers embedded in a generally A/U-enriched RNA region. The AU<sub>3</sub>A pentamers can be scattered within the region or can stagger or even overlap (see, *e.g.*, Chen et al., 1995, Trends Biol Sci 20:465). One or several AU<sub>3</sub>A pentamers can be replaced by expanded versions such as AU<sub>4</sub>A or AU<sub>5</sub>A heptamers (see, *e.g.*, Wilkund et al., 2002, J Biol Chem 277:40462 and Tholanikunnel and Malborn, 1997, J Biol Chem 272:11471). Single copies of the AU<sub>n</sub>A (where n = 3, 4, or 5) elements placed in a random sequence context are inactive. The minimal active ARE has the sequence U<sub>2</sub>AU<sub>n</sub>A(U/A)(U/A) (where n = 3, 4, or 5) (see, *e.g.* Worthington et al., 2002, J Biol Chem, 277:48558-64). The activity of certain AU-rich elements in promoting mRNA degradation is enhanced in the presence of distal uridine-rich sequences. These U-rich elements do not affect mRNA stability when present alone and thus have been termed "ARE enhancers" (see, *e.g.*, Chen et al., 1994, Mol. Cell. Biol. 14:416).

**[0013]** Most AREs function in mRNA decay regulation and translation initiation regulation by interacting with specific ARE-binding proteins (“AUBPs”). There are at least 14 known cellular proteins that bind to AU-rich elements. AUBP functional properties determine ARE involvement in one or both pathways. For example, ELAV/HuR binding to *c-fos* ARE inhibits *c-fos* mRNA decay (see, *e.g.*, Brennan & Steitz, 2001, Cell Mol Life Sci. 58:266), association of tristetraprolin with TNF $\alpha$  ARE dramatically enhances TNF $\alpha$  mRNA hydrolysis (see, *e.g.*, Carballo et al., 1998, Science 281:1001), whereas interaction of TIA-1 with the TNF $\alpha$  ARE does not alter the TNF $\alpha$  mRNA stability but inhibits TNF $\alpha$  translation (see, *e.g.*, Piecyk et al., 2000, EMBO J. 19:4154). Given its size, it is very likely that one copy of a typical ARE is capable of interacting with several AUBPs molecules. Therefore, it is contended that in the cell the competition of multiple AUBPs for the limited set of AUBP-binding sites in an ARE and the resulting “ARE proteome” determines the ARE regulatory output.

**[0014]** The mechanism of ARE-mediated mRNA decay is poorly understood. It has been established that mammalian mRNA degradation proceeds in 3' to 5' direction and that the first step is deadenylation by poly(A)-specific ribonuclease (“PARN”). Recent work indicates that following deadenylation a stable multi-ribonuclease complex, termed exosome, degrades the body of the message. Exosome alone is capable of initiating and accomplishing mRNA decay. However, the presence of certain AREs upregulates degradation efficiency. Available evidence suggests that AREs alone or bound by AUBPs help recruit exosome to the RNA (see, *e.g.*, Chen et al., 2001, Cell 107:451 and Mukherjee et al., 2002, EMBO J. 21:165).

**[0015]** It has been reported that degradation of some mRNAs depends on ongoing translation. Thus, the translation machinery can also serve as a ribonuclease-recruiting or stabilizing AUBPs-removing entity. Supporting evidence indicates that this mechanism may operate only on a subset of mRNAs under special cell growth conditions (see, *e.g.*, Curatola et al., 1995, Mol. Cell. Biol. 15:6331; Chen et al., 1995, Mol. Cell. Biol. 15:5777; Koeller et al., 1991, Proc. Natl. Acad. Sci. 88:7778; Savant-Bhonsale et al., 1992, Genes Dev. 6:1927; and Aharon & Schneider, 1993, Mol. Cell. Biol. 13:1971).

**[0016]** The mechanism of ARE-dependent translation regulation is understood even less well than that of ARE-mediated mRNA decay. It is not clear how a 3' UTR-localized element can affect translation initiation, a process that takes place in the 5' UTR. One plausible explanation comes from recent work showing that most or all cytoplasmic mRNPs are circularized via eIF4F - poly(A)-binding protein (“PABP”) interaction. This interaction

can bring AREs in the 3' UTR into close proximity to the translation initiation site (see, e.g., Wells et al., 1998, Mol. Cell. 2:135).

[0017] Citation or identification of any reference in Section 2 of this application is not an admission that such reference is available as prior art to the present invention.

### 3. SUMMARY OF THE INVENTION

[0018] The present invention provides methods for identifying a compound that modulates untranslated region-dependent expression of a target gene. In particular, the invention provides methods for identifying compounds that down-regulate the translation or the stability of an mRNA of a target gene that is associated with or has been linked to the onset, development, progression or severity of a particular disease or disorder, said compounds functioning, at least in part, by targeting one or more aspects of untranslated region-dependent expression of the target gene. The invention also provides methods for identifying compounds that upregulate the translation or the stability of an mRNA of a target gene whose expression is beneficial to a subject with a particular disease or disorder, said compounds functioning, at least, in part, by targeting one or more aspects of untranslated region-dependent expression of the target gene. The invention encompasses the use of the compounds identified utilizing the methods of the invention for modulating the expression of a target gene *in vitro* and *in vivo*. In particular, the invention encompasses the use of the compounds identified utilizing the methods of the invention for the prevention, treatment or amelioration of a disease or disorder or a symptom thereof.

[0019] The invention provides reporter gene-based assays for the identification of a compound that modulates untranslated region-dependent expression of a target gene. The reporter gene-based assays may be conducted by contacting a compound with a cell genetically engineered to express a nucleic acid comprising a reporter gene operably linked to one or more untranslated regions of a target gene, and measuring the expression of said reporter gene. Alternatively, the reporter gene-based assays may be conducted by contacting a compound with a cell-free translation mixture and a nucleic acid comprising a reporter gene operably linked to one or more untranslated regions of a target gene, and measuring the expression of said reporter gene. The alteration in reporter gene expression relative to a previously determined reference range or a control in such reporter-gene based assays indicates that a particular compound modulates untranslated region-dependent expression of a target gene. In a specific embodiment, a compound identified utilizing a reporter gene-based assay described herein alters the expression of the reporter gene by at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least

90%, at least 95%, or at least 99%, or at least 1.5 fold, at least 2 fold, at least 2.5 fold, at least 5 fold, at least 7.5 fold or at least 10 fold relative to a control (*e.g.*, PBS), the absence of a control or a previously determined reference range in an assay described herein or well-known in the art. In order to exclude the possibility that a particular compound is functioning solely by modulating the expression of a target gene in an untranslated region-independent manner, one or more mutations (*i.e.*, deletions, insertions, or nucleotide substitutions) may be introduced into the untranslated regions operably linked to a reporter gene and the effect on the expression of the reporter gene in a reporter gene-based assay described herein can be determined.

**[0020]** In one embodiment, the invention provides a method for identifying a compound that modulates untranslated region-dependent expression of a target gene, said method comprising: (a) expressing a nucleic acid comprising a reporter gene operably linked to two, three or more untranslated regions of said target gene in a cell; (b) contacting said cell with a member of a library of compounds; and (c) detecting the expression of said reporter gene, wherein a compound that modulates untranslated region-dependent regulation of expression is identified if the expression of said reporter gene in the presence of a compound is altered as compared to the expression of said reporter gene in the absence of said compound or the presence of a control (*e.g.*, phosphate buffered saline ("PBS")). In an alternative embodiment, the invention provides a method for identifying a compound that modulates untranslated region-dependent expression of a target gene, said method comprising: (a) expressing a nucleic acid comprising a reporter gene operably linked two, three or more untranslated regions of said target gene in a cell; (b) contacting said cell with a member of a library of compounds; and (c) detecting the expression of said reporter gene, wherein a compound that modulates untranslated region-dependent expression is identified if the expression of said reporter gene is altered in the presence of the compound relative to a previously determined reference range.

**[0021]** In another embodiment, the invention provides a method for identifying a compound that modulates untranslated region-dependent expression of a target gene, said method comprising: (a) expressing a nucleic acid consisting of a reporter gene operably linked to one, two, three or more untranslated regions of the target gene in a cell; (b) contacting said cell with a member of a library of compounds; and (c) detecting the expression of said reporter gene, wherein a compound that modulates untranslated region-dependent regulation of expression is identified if the expression of said reporter gene in the presence of a compound is altered as compared to the expression of said reporter gene in the absence of said compound or the presence of a control (*e.g.*, PBS). In an alternative

embodiment, the invention provides a method for identifying a compound that modulates untranslated region-dependent expression of a target gene, said method comprising: (a) expressing a nucleic acid consisting of a reporter gene operably linked to one, two, three or more untranslated regions of the target gene in a cell; (b) contacting said cell with a member of a library of compounds; and (c) detecting the expression of said reporter gene, wherein a compound that modulates untranslated region-dependent expression is identified if the expression of said reporter gene in the presence of the compound is altered relative to a previously determined reference range.

**[0022]** In another embodiment, the invention provides a method for identifying a compound that modulates untranslated region-dependent expression of a target gene, said method comprising: (a) contacting a member of a library of compounds with a cell-free translation mixture and a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of said target gene; and (b) detecting the expression of said reporter gene, wherein a compound that modulates untranslated region-dependent expression is identified if the expression of said reporter gene in the presence of a compound is altered as compared to the expression of said reporter gene in the absence of said compound or the presence of a control (*e.g.*, PBS). In an alternative embodiment, the invention provides a method for identifying a compound that modulates untranslated region-dependent expression of a target gene, said method comprising: (a) contacting a member of a library of compounds with a cell-free translation mixture and a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of said target gene; and (b) detecting the expression of said reporter gene, wherein a compound that modulates untranslated region-dependent expression of a target gene is identified if the expression of said reporter gene in the presence of a compound is altered relative to a previously determined reference range.

**[0023]** In another embodiment, the invention provides a method for identifying a compound that modulates untranslated region-dependent expression of a target gene, said method comprising: (a) contacting a member of a library of compounds with a cell containing a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of said target gene; and (b) detecting a reporter protein translated from said reporter gene, wherein a compound that modulates untranslated region-dependent expression is identified if the expression of said reporter gene in the presence of a compound is altered as compared to the expression of said reporter gene in the absence of said compound or the presence of a control (*e.g.*, PBS). In an alternative embodiment, the invention provides a method for identifying a compound that modulates untranslated

region-dependent expression of a target gene, said method comprising: (a) contacting a member of a library of compounds with a cell containing a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of said target gene; and (b) detecting expression of said reporter gene, wherein a compound that modulates untranslated region-dependent expression is identified if the expression of said reporter gene in the presence of a compound is altered relative to a previously determined reference range.

**[0024]** The invention also provides methods of identifying compounds that upregulate untranslated region-dependent expression of a target gene utilizing the reporter gene-based assays described herein. In a specific embodiment, the invention provides a method of upregulating untranslated region-dependent expression of a target gene, said method comprising (a) contacting a compound with a cell containing a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of said target gene; and (b) detecting a reporter gene protein translated from said reporter gene, wherein a compound that upregulates untranslated region dependent expression is identified if the expression of said reporter gene in the presence of a compound is increased relative to the absence of the compound or a previously determined reference range. In another embodiment, the invention provides a method of upregulating untranslated region-dependent expression of a target gene, said method comprising (a) contacting a compound with a cell-free translation mixture and a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of said target gene; and (b) detecting a reporter gene protein translated from said reporter gene, wherein a compound that upregulates untranslated region dependent expression is identified if the expression of said reporter gene in the presence of a compound is increased relative to the absence of the compound or a previously determined reference range.

**[0025]** The invention also provides methods of identifying compounds that down-regulate untranslated region-dependent expression of a target gene utilizing the reporter gene-based assays described herein. In a specific embodiment, the invention provides a method of down-regulating untranslated region-dependent expression of a target gene, said method comprising (a) contacting a compound with a cell containing a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of said target gene; and (b) detecting a reporter gene protein translated from said reporter gene, wherein a compound that down-regulates untranslated region dependent expression is identified if the expression of said reporter gene in the presence of a compound is decreased relative to the absence of the compound or a previously determined reference range. In

another embodiment, the invention provides a method of down-regulating untranslated region-dependent expression of a target gene, said method comprising (a) contacting a compound with a cell-free translation mixture and a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of said target gene; and (b) detecting a reporter gene protein translated from said reporter gene, wherein a compound that down-regulates untranslated region dependent expression is identified if the expression of said reporter gene in the presence of a compound is decreased relative to the absence of the compound or a previously determined reference range.

[0026] In accordance with the invention, the step of contacting a compound with a cell, or cell-free translation mixture and a nucleic acid in the reporter gene-based assays described herein is preferably conducted in an aqueous solution comprising a buffer and a combination of salts. In a specific embodiment, the aqueous solution approximates or mimics physiologic conditions. In another specific embodiment, the aqueous solution further comprises a detergent or a surfactant.

[0027] The present invention provides methods of identifying environmental stimuli (e.g., exposure to different concentrations of CO<sub>2</sub> and/or O<sub>2</sub>, stress, temperature shifts, and different pHs) that modulate untranslated region-dependent expression of a target gene utilizing the reporter gene-based assays described herein. In particular, the invention provides a method of identifying an environmental stimulus, said method comprising (a) contacting a cell containing a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of said target gene with an environmental stimulus; and (b) detecting a reporter gene protein translated from said reporter gene, wherein a compound that modulates untranslated region dependent expression is identified if the expression of said reporter gene in the presence of an environmental stimuli is altered relative to the absence of the compound or a previously determined reference range. In a specific embodiment, the environmental stimuli is not hypoxia. In another embodiment, the environmental stimuli does not include a compound.

[0028] The reporter gene constructs utilized in the reporter gene-based assays described herein may comprise a 5' untranslated region ("UTR") of a target gene, a 3' UTR of a target gene, or a 5' UTR and a 3' UTR of a target gene operably linked to a reporter gene. In a specific embodiment, a reporter gene construct utilized in the reporter gene-based assays described herein comprises a 5' UTR of a target gene with a stable hairpin secondary structure operably linked to a reporter gene. In a preferred embodiment, a reporter gene construct utilized in the reporter gene-based assays described herein comprises a 5' UTR and a 3' UTR of a target gene. The untranslated regions of a target gene utilized to

construct a reporter gene construct may comprise one or more of the following elements: an iron response element ("IRE"), Internal ribosome entry site ("IRES"), upstream open reading frame ("uORF"), male specific lethal element ("MSL-2"), G quartet element, 5'-terminal oligopyrimidine tract ("TOP"), AU-rich element ("ARE"), selenocysteine insertion sequence ("SECIS"), histone stem loop, cytoplasmic polyadenylation element ("CPE"), nanos translational control element, amyloid precursor protein element ("APP"), translational regulation element ("TGE")/direct repeat element ("DRE"), Bruno element ("BRE"), and a 15-lipoxygenase differentiation control element ("15-LOX-DICE").

[0029] In addition to untranslated regions, the reporter gene constructs utilized in the reporter gene-based assays described herein may comprise one, two, three or more introns within the open reading frame ("ORF") of the reporter gene. Further, the 3' end of a reporter gene may be polyadenylated and/or the 5' end may be capped. In a specific embodiment, the 5' end of the reporter gene is not capped.

[0030] The reporter gene constructs utilized in the reporter gene-based assays described herein may comprise an untranslated region of a gene whose expression is associated with or has been linked to the onset, development, progression or severity of a particular disease or disorder. Alternatively, the reporter gene constructs utilized in the reporter gene-based assays described herein may comprise an untranslated region of a gene whose expression is beneficial to a subject with a particular disease or disorder. Examples of genes from which the untranslated regions may be obtained include, but are not limited, the gene encoding tumor necrosis factor alpha ("TNF-a"), the gene encoding granulocyte-macrophage colony stimulating factor ("GM-CSF"), the gene encoding granulocyte colony stimulating factor ("G-CSF"), the gene encoding interleukin 2 ("IL-2"), the gene encoding interleukin 6 ("IL-6"), the gene encoding vascular endothelial growth factor ("VEGF"), the genome encoding hepatitis C virus ("HCV"), the gene encoding survivin, or the gene encoding Her-2. In a specific embodiment, an untranslated region is obtained or derived from Her-2 and/or VEGF. In another embodiment, an untranslated region is not obtained or derived from the gene encoding Her-2. In another embodiment, an untranslated region is not obtained or derived from the gene encoding VEGF. In another embodiment, an untranslated region is not obtained or derived from the genes encoding VEGF and Her-2.

[0031] Any reporter gene well-known to one of skill in the art may be utilized in the reporter gene constructs described herein. Examples of reporter genes include, but are not limited to, the gene encoding firefly luciferase, the gene coding renilla luciferase, the genes encoding click beetle luciferase, the gene encoding green fluorescent protein, the gene encoding yellow fluorescent protein, the gene encoding red fluorescent protein, the gene



encoding cyan fluorescent protein, the gene encoding blue fluorescent protein, the gene encoding beta-galactosidase, the gene encoding beta-glucoronidase, the gene encoding beta-lactamase, the gene encoding chloramphenicol acetyltransferase, and the gene encoding alkaline phosphatase.

[0032] The reporter gene-based assays described herein may be conducted in a cell genetically engineered to express a reporter gene or *in vitro* utilizing a cell-free translation mixture. Any cell or cell line of any species well-known to one of skill in the art may be utilized in accordance with the methods of the invention. Further, a cell-free translation mixture may be derived from any cell or cell line of any species well-known to one of skill in the art. Examples of cells and cell types include, but are not limited to, human cells (*e.g.*, HeLa cells and 293 cells), yeast, mouse cells (*e.g.*, cultured mouse cells), rat cells (*e.g.*, cultured rat cells), Chinese hamster ovary ("CHO") cells, *Xenopus* oocytes, cancer cells (*e.g.*, undifferentiated cancer cells), primary cells, reticulocytes, wheat germ, rye embryo, or bacterial cells.

[0033] The compounds utilized in the reporter gene-based assays described herein may be members of a library of compounds. In specific embodiment, the compound is selected from a combinatorial library of compounds comprising peptoids; random biooligomers; diversomers such as hydantoins, benzodiazepines and dipeptides; vinylogous polypeptides; nonpeptidal peptidomimetics; oligocarbamates; peptidyl phosphonates; peptide nucleic acid libraries; antibody libraries; carbohydrate libraries; and small organic molecule libraries. In a preferred embodiment, the small organic molecule libraries are libraries of benzodiazepines, isoprenoids, thiazolidinones, metathiazanones, pyrrolidines, morpholino compounds, or diazepindiones.

[0034] Once a compound that modulates untranslated region-dependent expression of a target gene is identified, the structure of the compound may be determined utilizing well-known techniques or by referring to a predetermined code. For example, the structure of the compound may be determined by mass spectroscopy, NMR, vibrational spectroscopy, or X-ray crystallography.

[0035] A compound identified in accordance with the methods of the invention may directly bind to an RNA transcribed from a target gene. Alternatively, a compound identified in accordance with the methods of invention may bind to one or more *trans*-acting factors (such as, but not limited to, proteins) that modulate untranslated region-dependent expression of a target gene. Further, a compound identified in accordance with the methods of invention may disrupt an interaction between the 5' UTR and the 3' UTR.

**[0036]** In a specific embodiment, a compound identified in accordance with the methods of the invention reduces the translation efficiency and/or stability of an mRNA transcribed from a target gene by at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99%, or at least 1.5 fold, at least 2 fold, at least 2.5 fold, at least 5 fold, at least 7.5 fold or at least 10 fold relative to a control (*e.g.*, PBS), the absence of a control or a previously determined reference range in an assay described herein or well-known in the art. In another embodiment, a compound identified in accordance with the methods of the invention reduces the translation efficiency and/or stability of an mRNA transcribed from a target gene by at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99%, or at least 1.5 fold, at least 2 fold, at least 2.5 fold, at least 5 fold, at least 7.5 fold or at least 10 fold relative to a control (*e.g.*, PBS), the absence of a control or a previously determined reference range in an assay described herein or well-known in the art.

**[0037]** A compound that modulates untranslated region-dependent expression in a reporter gene-based assay described herein may be subsequently tested in *in vitro* assays (*e.g.*, cell-free assays) or *in vivo* assays (*e.g.*, cell-based assays) well-known to one of skill in the art or described herein for the effect of said compound on the expression of the target gene from which the untranslated regions of the reporter gene construct were derived. Further, to assess the specificity of a particular compound's effect on untranslated region-dependent expression of a target gene, the effect of said compound on the expression of one or more genes (preferably, a plurality of genes) can be determined utilizing assays well-known to one of skill in the art or described herein. In a preferred embodiment, a compound identified utilizing the reporter gene-based assays described herein has a specific effect on the expression of only one gene or a group of genes within the same signaling pathway.

**[0038]** In a specific embodiment, the specificity of a particular compound for an untranslated region of a target gene is determined by (a) contacting the compound of interest with a cell containing a nucleic acid comprising a reporter gene operably linked to an UTR of a different gene; and (b) detecting a reporter gene protein translated from the reporter gene, wherein the compound is specific for the untranslated region of the target gene if the expression of said reporter gene in the presence of the compound is not altered or is not substantially altered relative to a previously determined reference range, or the expression in the absence of the compound or the presence of a control (*e.g.*, PBS). In

another embodiment, the specificity of a particular compound for an untranslated region of a target gene is determined by (a) contacting the compound of interest with a panel of cells, each cell in a different well of a container (*e.g.*, a 48 or 96 well microtiter plate) and each cell containing a nucleic acid comprising a reporter gene operably linked to an UTR of a different gene; and (b) detecting a reporter gene protein translated from the reporter gene, wherein the compound is specific for the untranslated region of the target gene if the expression of said reporter gene in the presence of the compound is not altered or is not substantially altered relative to a previously determined reference range, or the expression in the absence of the compound or the presence of a control (*e.g.*, PBS). In accordance with this embodiment, the panel may comprise 5, 7, 10, 15, 20, 25, 50, 75, 100 or more cells. In another embodiment, the specificity of a particular compound for an untranslated region of a target gene is determined by (a) contacting the compound of interest with a cell-free translation mixture and a nucleic acid comprising a reporter gene operably linked to an UTR of a different gene; and (b) detecting a reporter gene protein translated from the reporter gene, wherein the compound is specific for the untranslated region of the target gene if the expression of said reporter gene in the presence of the compound is not altered or is not substantially altered relative to a previously determined reference range, or the expression in the absence of the compound or the presence of a control (*e.g.*, PBS). As used herein, the term “not substantially altered” means that the compound alters the expression of the reporter gene or target gene by less than 20%, less than 15%, less than 10%, less than 5%, or less than 2% relative to a negative control such as PBS.

[0039] The invention provides for methods for treating, preventing or ameliorating one or more symptoms of a disease or disorder associated with the aberrant expression of a target gene, said method comprising administering to a subject in need thereof a therapeutically or prophylactically effective amount of a compound, or a pharmaceutically acceptable salt thereof, identified according to the methods described herein. In one embodiment, the target gene is aberrantly overexpressed. In another embodiment, the target gene is expressed at an aberrantly low level. In particular, the invention provides for a method of treating or preventing a disease or disorder or ameliorating a symptom thereof, said method comprising administering to a subject in need thereof an effective amount of a compound, or a pharmaceutically acceptable salt thereof, identified according to the methods described herein, wherein said effective amount increases the expression of a target gene beneficial in the treatment or prevention of said disease or disorder. The invention also provides for a method of treating or preventing a disease or disorder or ameliorating a symptom thereof, said method comprising administering to a subject in need

thereof an effective amount of a compound, or a pharmaceutically acceptable salt thereof, identified according to the methods described herein, wherein said effective amount decreases the expression of a target gene whose expression is associated with or has been linked to the onset, development, progression or severity of said disease or disorder. In a specific embodiment, the disease or disorder is a proliferative disorder, an inflammatory disorder, an infectious disease, a genetic disorder, an autoimmune disorder, a cardiovascular disease, or a central nervous system disorder. In an embodiment wherein the disease or disorder is an infectious disease, the infectious disease can be caused by a fungal infection, a bacterial infection, a viral infection, or an infection caused by another type of pathogen.

**[0040]** The invention provides a method for identifying a compound that inhibits or reduces angiogenesis, said method comprising: (a) contacting a member of a library of compounds with a cell containing a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of a target gene; and (b) detecting the expression of said reporter gene, wherein if a compound that reduces the expression of said reporter gene relative to a previously determined reference range, or to the expression of said reporter gene in the absence of said compound or in the presence of a control (*e.g.*, PBS) is detected in (b), then (c) contacting the compound with a tumor cell and detecting the proliferation of said tumor cell, so that if the compound reduces or inhibits the proliferation of the tumor cell, the compound is identified as a compound that inhibits or reduces angiogenesis. The invention provides a method for identifying a compound that inhibits or reduces angiogenesis, said method comprising: (a) contacting a cell-free translation mixture with a member of a library of compounds and a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of a target gene; and (b) detecting the expression of said reporter gene, wherein if a compound that reduces the expression of said reporter gene relative to a previously determined reference range, or to the expression of said reporter gene in the absence of said compound or in the presence of a control (*e.g.*, PBS) is detected in (b), then (c) contacting the compound with a tumor cell and detecting the proliferation of said tumor cell, so that if the compound reduces or inhibits the proliferation of the tumor cell, the compound is identified as a compound that inhibits or reduces angiogenesis. In a specific embodiment, the compound is further tested in an animal model for angiogenesis by, *e.g.*, administering said compound to said animal model and verifying that angiogenesis is inhibited by said compound in said animal model. In a preferred embodiment, the target gene is VEGF. In another embodiment, the compound identified in accordance with the methods of the invention inhibits or reduces angiogenesis by at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least

50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or at least 99%, or at least 1.5 fold, at least 2 fold, at least 2.5 fold, at least 5 fold, at least 7.5 fold, or at least 10 fold relative to a control (*e.g.*, PBS) in an assay described herein or well-known in the art.

**[0041]** The invention provides for a method for identifying a therapeutic agent for the treatment or prevention of cancer, or amelioration of a symptom thereof, said method comprising: (a) contacting a member of a library of compounds with a cell containing a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of a target gene; and (b) detecting the expression of said reporter gene, wherein if a compound that reduces the expression of said reporter gene relative to a previously determined reference range, or the expression of said reporter gene in the absence of said compound or the presence of a control (*e.g.*, PBS) is detected in (b), then (c) contacting the compound with a cancer cell and detecting the proliferation of said cancer cell, so that if the compound reduces or inhibits the proliferation of the cancer cell, the compound is identified as a therapeutic agent for the treatment or prevention of cancer, or amelioration of a symptom thereof. The invention also provides for a method for identifying a therapeutic agent for the treatment or prevention of cancer, or amelioration of a symptom thereof, said method comprising: (a) contacting a cell-free translation mixture with a member of a library of compounds and a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of a target gene; and (b) detecting the expression of said reporter gene, wherein if a compound that reduces the expression of said reporter gene relative to a previously determined reference range, or the expression of said reporter gene in the absence of said compound or the presence of a control (*e.g.*, PBS) is detected in (b), then (c) contacting the compound with a cancer cell and detecting the proliferation of said cancer cell, so that if the compound reduces or inhibits the proliferation of the cancer cell, the compound is identified as a therapeutic agent for the treatment or prevention of cancer, or amelioration of a symptom thereof. In a specific embodiment, the compound is further tested in an animal model for cancer by, *e.g.*, administering said compound to said animal model and verifying that the compound is effective in reducing the proliferation or spread of cancer cells in said animal model. In a preferred embodiment, the target gene is survivin.

**[0042]** In a specific embodiment, the invention provides for a method of identifying a therapeutic agent for the treatment or prevention of breast cancer, or amelioration of a symptom thereof, said method comprising: (a) contacting a member of a library of compounds with a cell containing a nucleic acid comprising a reporter gene operably linked

to one, two, three or more untranslated regions of a target gene; and (b) detecting the expression of said reporter gene, wherein if a compound that reduces the expression of said reporter gene relative a previously determined reference range, or the expression of said reporter gene in the absence of said compound or the presence of a control is detected in (b), then (c) contacting the compound with a breast cancer cell and detecting the proliferation of said breast cancer cell, so that if the compound reduces or inhibits the proliferation of the breast cancer cell, the compound is identified as a therapeutic agent for the treatment or prevention of breast cancer, or amelioration of a symptom thereof. In another embodiment, the invention provides for a method of identifying a therapeutic agent for the treatment or prevention of breast cancer, or amelioration of a symptom thereof, said method comprising: (a) contacting a cell-free translation mixture with a member of a library of compounds and a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of a target gene; and (b) detecting the expression of said reporter gene, wherein if a compound that reduces the expression of said reporter gene relative to a previously determined reference range, or the expression of said reporter gene in the absence of said compound or the presence of a control is detected in (b), then (c) contacting the compound with a breast cancer cell and detecting the proliferation of said breast cancer cell, so that if the compound reduces or inhibits the proliferation of the breast cancer cell, the compound is identified as a therapeutic agent for the treatment or prevention of breast cancer, or amelioration of a symptom thereof. In accordance with these embodiments, the compound may be further tested in an animal model for breast cancer by, *e.g.*, administering said compound to said animal model and verifying that the compound is effective in reducing the proliferation or spread of breast cancer cells in said animal model. In a preferred embodiment, the target gene is Her-2.

[0043] The invention also provides methods for upregulating or downregulating the expression of a target gene utilizing a compound identified in accordance with the methods described herein. The upregulation or downregulation of a target gene is particularly useful *in vitro* when attempting to produce a protein encoded by said target gene for use as a therapeutic or prophylactic agent, or in experiments conducted to, *e.g.*, identify the function or efficacy of said protein. In particular, the invention provides a method of modulating the expression of a target gene, said method comprising contacting a cell with an effective amount of a compound or pharmaceutically acceptable derivative thereof, identified according to the methods described herein. In one embodiment, the cell is a eucaryotic cell. In another embodiment, the cell is a procaryotic cell.

[0044] The invention further provides methods for verifying or confirming the ability of a compound to modulate untranslated region-dependent expression of a target gene. The ability of a compound to modulate untranslated region-dependent expression of a target gene can be verified or confirmed utilizing any of the assays described herein to identify such a compound. In a first embodiment, the invention provides a method for verifying the ability of a compound to modulate untranslated region-dependent expression of a target gene, said method comprising: (a) expressing a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of said target gene in a cell; (b) contacting said cell with a compound; and (c) detecting the expression of said reporter gene, wherein a compound that modulates untranslated region-dependent expression is verified if the expression of said reporter gene in the presence of a compound is altered as compared to a previously determined reference range or the expression of said reporter gene in the absence of said compound or the presence of a control.

[0045] In a second embodiment, the invention provides a method for verifying the ability of a compound to modulate untranslated region-dependent expression of a target gene, said method comprising: (a) contacting a compound with a cell-free translation mixture and a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of said target gene; and (b) detecting the expression of said reporter gene, wherein a compound that modulates untranslated region-dependent expression is verified if the expression of said reporter gene in the presence of a compound is altered as compared to a previously determined reference range or the expression of said reporter gene in the absence of said compound or the presence of a control.

[0046] In a third embodiment, the invention provides a method for verifying the ability of a compound to modulate untranslated region-dependent expression of a target gene, said method comprising: (a) contacting a compound with a cell containing a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of said target gene; and (b) detecting the expression of said reporter gene, wherein a compound that modulates untranslated region-dependent expression is verified if the expression of said reporter gene in the presence of a compound is altered as compared to a previously determined reference range or the expression of said reporter gene in the absence of said compound or the presence of a control.

### 3.1. Terminology

[0047] As used herein, the term "5' cap" refers to a methylated guanine cap, *e.g.*, a 7 methylguanosine (5'-5') RNA triphosphate, that is added to the 5' end of a pre-mRNA.

[0048] As used herein, the term “ARE” refers to an adenylate uridylate rich element in the the 3' UTR of a mRNA.

[0049] As used herein, the term “compound” refers to any agent or complex that is being tested for its ability to modulate untranslated region-dependent expression of a target gene, or any agent or complex identified by the methods described herein. Examples of compounds include, but are not limited to, proteins, polypeptides, peptides, peptide analogs (including peptides comprising non-naturally occurring amino acids, *e.g.*, D-amino acids, phosphorous analogs of amino acids, such as  $\alpha$ -amino phosphoric acids and  $\alpha$ -amino phosphoric acids, or amino acids having non-peptide linkages), nucleic acids, nucleic acid analogs such as phosphorothioates and PNAs, hormones, antigens, antibodies, lipids, fatty acids, synthetic or naturally occurring drugs, opiates, dopamine, serotonin, catecholamines, thrombin, acetylcholine, prostaglandins, organic molecules, pheromones, adenosine, sucrose, glucose, lactose and galactose.

[0050] As used herein, the term “CUG repeat” refers to a repeat of a cytosine-uracil-guanine triplet in the 3' UTR of a mRNA.

[0051] As used herein, the term “cytosine rich element” refers to cytosine-rich stability determinant sequences in the 3' UTR of a mRNA.

[0052] As used herein, the terms “disorder” and “disease” refer to a condition in a subject.

[0053] As used herein, the term “effective amount” refers to the amount of a compound which is sufficient to reduce or ameliorate the severity, duration and/or a disease or disorder or a symptom thereof, prevent the advancement of a disease or disorder, cause regression of a disease or disorder, prevent the recurrence, development, or onset of one or more symptoms associated with a disease or disorder, or enhance or improve the prophylactic or therapeutic effect(s) of another therapy (*e.g.*, prophylactic or therapeutic agent).

[0054] As used herein, the term “fragment” refers to a nucleotide sequence comprising an nucleic acid sequence of at least 5 contiguous nucleic acid residues, at least 10 contiguous nucleic acid residues, at least 15 contiguous nucleic acid residues, at least 20 contiguous nucleic acid residues, at least 25 contiguous nucleic acid residues, at least 40 contiguous nucleic acid residues, at least 50 contiguous nucleic acid residues, at least 60 contiguous nucleic acid residues, at least 70 contiguous nucleic acid residues, at least contiguous 80 nucleic acid residues, at least contiguous 90 nucleic acid residues, at least contiguous 100 nucleic acid residues, at least contiguous 125 nucleic acid residues, at least 150 contiguous nucleic acid residues, at least contiguous 175 nucleic acid residues, at least contiguous 200 nucleic acid residues, or at least contiguous 250 nucleic acid residues of the



nucleotide sequence of untranslated region of a target gene. In a specific embodiment, a fragment of a untranslated region of a target gene retains at least one element of the untranslated region (*e.g.*, an IRES).

[0055] As used herein, the term “target RNA” refers to an RNA of interest, *i.e.*, the RNA transcribed from a target gene or a gene of interest. In a preferred embodiment, the target RNA contains one or more untranslated regions, and more preferably, contains at least one element of the untranslated region (*e.g.*, an IRES).

[0056] As used herein, the term “host cell” includes a particular subject cell transformed or transfected with a nucleic acid molecule and the progeny or potential progeny of such a cell. Progeny of such a cell may not be identical to the parent cell transfected with the nucleic acid molecule due to mutations or environmental influences that may occur in succeeding generations or integration of the nucleic acid molecule into the host cell genome.

[0057] As used herein, the term “in combination” refers to the use of more than one therapies (*e.g.*, prophylactic and/or therapeutic agents). The use of the term “in combination” does not restrict the order in which therapies (*e.g.*, prophylactic and/or therapeutic agents) are administered to a subject with a particular disease or disorder. A first therapy (*e.g.*, a prophylactic or therapeutic agent such as, *e.g.*, a compound identified in accordance with the methods of the invention) can be administered prior to (*e.g.*, 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), concomitantly with, or subsequent to (*e.g.*, 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of a second therapy (*e.g.*, a prophylactic or therapeutic agent such as, *e.g.*, a chemotherapeutic agent, an anti-inflammatory agent or a TNF- $\alpha$  antagonist) to a subject with a particular disease or disorder.

[0058] As used herein, the term “IRE” refers to an iron response element in the 5' UTR or 3' UTR of a mRNA.

[0059] As used herein, the term “IRES” refers to an internal ribosome entry site in the 5' UTR of a mRNA.

[0060] As used herein, the term “library” refers to a plurality of compounds. A library can be a combinatorial library, *e.g.*, a collection of compounds synthesized using combinatorial chemistry techniques, or a collection of unique chemicals of low molecular weight (less than 1000 daltons) that each occupy a unique three-dimensional space.

[0061] As used herein, the term “ORF” refers to the open reading frame of a mRNA, *i.e.*, the region of the mRNA that is translated into protein.

[0062] As used herein, the terms “non-responsive” and “refractory” describe patients treated with a currently available therapy (*e.g.*, a prophylactic or therapeutic agent) for a disease or disorder, which is not clinically adequate to relieve one or more symptoms associated with such disease or disorder. Typically, such patients suffer from severe, persistently active disease and require additional therapy to ameliorate the symptoms associated with their disease or disorder.

[0063] As used herein, the phrase “pharmaceutically acceptable salt(s),” includes, but is not limited to, salts of acidic or basic groups that may be present in compounds identified using the methods of the present invention. Compounds that are basic in nature are capable of forming a wide variety of salts with various inorganic and organic acids. The acids that can be used to prepare pharmaceutically acceptable acid addition salts of such basic compounds are those that form non-toxic acid addition salts, *i.e.*, salts containing pharmacologically acceptable anions, including but not limited to sulfuric, citric, maleic, acetic, oxalic, hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, isonicotinate, acetate, lactate, salicylate, citrate, acid citrate, tartrate, oleate, tannate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate, gluconate, glucuronate, saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate and pamoate (*i.e.*, 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)) salts. Compounds that include an amino moiety may form pharmaceutically acceptable salts with various amino acids, in addition to the acids mentioned above. Compounds that are acidic in nature are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include alkali metal or alkaline earth metal salts and, particularly, calcium, magnesium, sodium lithium, zinc, potassium, and iron salts.

[0064] As used herein, the term “poly(A) tail” refers to a polyadenylic acid tail that is added to the 3' end of a pre-mRNA.

[0065] As used herein, the terms “prophylactic agent” and “prophylactic agents” refer to any agent(s) which can be used in the prevention of a particular disease or disorder. In certain embodiments, the term “prophylactic agent” refers to a compound identified in the screening assays described herein. In certain other embodiments, the term “prophylactic agent” does not refer a compound identified in the screening assays described herein.

[0066] As used herein, the phrase “prophylactically effective amount” refers to the amount of a therapy (*e.g.*, a prophylactic agent) which is sufficient to result in the

prevention of the development, recurrence or onset of a disease or disorder or one or more symptoms associated thereof.

[0067] As used herein, the terms “prevent”, “preventing” and “prevention” refer to the prevention of the development, recurrence or onset of a disease or disorder or one or more symptoms thereof resulting from the administration of one or more compounds identified in accordance the methods of the invention or the administration of a combination of such a compound and a known therapy for a particular disease or disorder.

[0068] As used herein, the term “previously determined reference range” refers to a reference range for the expression and/or the activity of a reporter gene or a target gene by a particular cell or in a particular cell-free translation mixture. Each laboratory will establish its own reference range for each particular assay, each cell type and each cell-free translation mixture. In a preferred embodiment, at least one positive control and at least one negative control are included in each batch of compounds analyzed.

[0069] As used herein, the term “small molecules” and analogous terms include, but are not limited to, peptides, peptidomimetics, amino acids, amino acid analogs, polynucleotides, polynucleotide analogs, nucleotides, nucleotide analogs, organic or inorganic compounds (*i.e.*, including heteroorganic and organometallic compounds) having a molecular weight less than about 10,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 5,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 1,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 500 grams per mole, organic or inorganic compounds having a molecular weight less than about 100 grams per mole, and salts, esters, and other pharmaceutically acceptable forms of such compounds. Salts, esters, and other pharmaceutically acceptable forms of such compounds are also encompassed.

[0070] As used herein, the terms “subject” and “patient” are used interchangeably herein. The terms “subject” and “subjects” refer to an animal, preferably a mammal including a non-primate (*e.g.*, a cow, pig, horse, cat, dog, rat, and mouse) and a primate (*e.g.*, a monkey such as a cynomolgous monkey and a human), and more preferably a human. In one embodiment, the subject is refractory or non-responsive to current therapies for a disease or disorder (*e.g.*, viral infections, fungal infections, bacterial infections, proliferative diseases or inflammatory diseases). In another embodiment, the subject is a farm animal (*e.g.*, a horse, a cow, a pig, *etc.*) or a household pet (*e.g.*, a dog or a cat). In a preferred embodiment, the subject is a human.

[0071] As used herein, the term “synergistic” refers to a combination of a compound identified using one of the methods described herein, and another therapy (preferably, a therapy which has been or is currently being used to prevent or treat a particular disease or disorder) which is more effective than the additive effects of the therapies. A synergistic effect of a combination of therapies (*e.g.*, prophylactic or therapeutic agents) permits the use of lower dosages of one or more of the therapies and/or less frequent administration of said therapies to a subject with a particular disease or disorder. The ability to utilize lower dosages of a therapy (*e.g.*, prophylactic or therapeutic agent) and/or to administer said therapy less frequently reduces the toxicity associated with the administration of said therapy to a subject without reducing the efficacy of said therapies in the prevention or treatment of a particular disease or disorder. In addition, a synergistic effect can result in improved efficacy of therapies in the prevention or treatment of a particular disease or disorder. Finally, a synergistic effect of a combination of therapies (*e.g.*, prophylactic or therapeutic agents) may avoid or reduce adverse or unwanted side effects associated with the use of either therapy alone.

[0072] As used herein, the term “target gene” refers to a gene or nucleotide sequence encoding a protein or polypeptide of interest. In a preferred embodiment, the gene or nucleotide sequence comprises an untranslated region.

[0073] As used herein, a “target nucleic acid” refers to RNA, DNA, or a chemically modified variant thereof. In a preferred embodiment, the target nucleic acid is RNA. In a preferred embodiment, the target nucleic acid refers to the untranslated region of an mRNA, such as, but not limited to, a 5' UTR and a 3' UTR. In another embodiment, the target nucleic acid refers to an open reading frame of an mRNA. A target nucleic acid also refers to tertiary structures of the nucleic acids, such as, but not limited to loops, bulges, pseudoknots, guanosine quartets and turns. A target nucleic acid also refers to RNA elements such as, but not limited to, the HIV TAR element, internal ribosome entry site, instability elements, and adenylate uridylate-rich elements, which are described in Section 5.1. Non-limiting examples of target nucleic acids are presented in Section 5.1 and Section 6.

[0074] As used herein, the terms “therapeutic agent” and “therapeutic agents” refer to any agent(s) which can be used in the prevention, treatment, management or amelioration of one or more symptoms of a particular disease or disorder. In certain embodiments, the term “therapeutic agent” refers to a compound identified in the screening assays described herein. In other embodiments, the term “therapeutic agent” does not refer to a compound identified in the screening assays described herein.

[0075] As used herein, the term “therapeutically effective amount” refers to that amount of a therapy (*e.g.*, a therapeutic agent) sufficient to reduce the severity of a disease or disorder, reduce the duration of a disease or disorder, ameliorate one or more symptoms of a disease or disorder, prevent advancement of a disease or disorder, cause regression of the disease or disorder, or to enhance or improve the therapeutic effect(s) of another therapeutic agent. In a specific embodiment, with respect to the treatment of cancer, a therapeutically effective amount refers to the amount of a therapy (*e.g.*, a therapeutic agent) that inhibits or reduces the proliferation of cancerous cells, inhibits or reduces the spread of tumor cells (metastasis), inhibits or reduces the onset, development or progression of one or more symptoms associated with cancer, or reduces the size of a tumor. Preferably, with respect to the treatment of cancer, a therapeutically effective amount of a therapy (*e.g.*, a therapeutic agent) reduces the proliferation of cancerous cells or the size of a tumor by at least 5%, preferably at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% relative to a control (*e.g.*, PBS) in an assay described herein or well-known in the art.

[0076] In another embodiment, with respect to the treatment of a viral infection, a therapeutically effective amount refers to the amount of a therapy (*e.g.*, a therapeutic agent) sufficient to reduce or inhibit the replication of a virus, inhibit or reduce the spread of the virus to other tissues or subjects, or ameliorate one or more symptoms associated with the viral infection. Preferably, with respect to a viral infection, a therapeutically effective amount of a therapy (*e.g.*, a therapeutic agent) reduces the replication or spread of a virus by at least 5%, preferably at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% relative to a control (*e.g.*, PBS) in an assay described herein or well-known in the art.

[0077] In another embodiment, with respect to the treatment of a fungal infection, a therapeutically effective amount refers to the amount of a therapy (*e.g.*, a therapeutic agent) sufficient to inhibit or reduce the replication of the fungus, inhibit or reduce the replication or spread of the fungus to other tissues or subjects, or ameliorate one or more symptoms associated with the fungal infection. Preferably, with respect to a fungal infection, a therapeutically effective amount of a therapy (*e.g.*, a therapeutic agent) reduces the spread of a fungus by at least 5%, preferably at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least

60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% relative to a control (*e.g.*, PBS) in an assay described herein or well-known in the art.

[0078] In another embodiment, with respect to the treatment of a bacterial infection, a therapeutically effective amount refers to the amount of a therapy (*e.g.*, a therapeutic agent) sufficient to inhibit or reduce the replication of the bacteria, to inhibit or reduce the replication or spread of the bacteria to other tissues or subjects, or ameliorate one or more symptoms associated with the bacterial infection. Preferably, with respect to a bacterial infection, a therapeutically effective amount of a therapy (*e.g.*, a therapeutic agent) reduces the spread of a bacteria by at least 5%, preferably at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% relative to a control (*e.g.*, PBS) in an assay described herein or well-known in the art.

[0079] In another embodiment, with respect to the treatment of an inflammatory disorder, a therapeutically effective amount refers to the amount of a therapy (*e.g.*, a therapeutic agent) that reduces the inflammation of a joint, organ or tissue. Preferably, with respect to an inflammatory disorder, a therapeutically effective amount of a therapy (*e.g.*, a therapeutic agent) reduces the inflammation of a joint, organ or tissue by at least 5%, preferably at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% relative to a control (*e.g.*, PBS) in an assay described herein or well-known in the art.

[0080] As used herein, the terms “therapies” and “therapy” can refer to any protocol(s), method(s), and/or agent(s) that can be used in the prevention, treatment, management, or amelioration of a disease or disorder or one or more symptoms thereof.

[0081] As used herein, the terms “treat”, “treatment” and “treating” refer to the reduction or amelioration of the progression, severity and/or duration of a disease or disorder or one or more symptoms thereof resulting from the administration of one or more compounds identified in accordance the methods of the invention, or the administration of a combination of therapies (*e.g.*, a compound identified in accordance with the methods of the invention and another therapeutic agent). In certain embodiments, such terms refer to the inhibition or reduction in the proliferation of cancerous cells, the inhibition or reduction the spread of tumor cells (metastasis), the inhibition or reduction in the onset, development or progression of one or more symptoms associated with cancer, or the reduction in the size of

a tumor. In other embodiments, such terms refer to the reduction or inhibition of the replication of a virus, the inhibition or reduction in the spread of a virus to other tissues or subjects, or the amelioration of one or more symptoms associated with a viral infection. In other embodiments, such terms refer to the reduction or inhibition of the replication of a fungus, the reduction or inhibition in the spread of a fungus to other tissues or subjects, or the amelioration of one or more symptoms associated with a fungal infection. In other embodiments, such terms refer to the inhibition or reduction of the replication of a bacteria, the inhibition or reduction in the spread of a bacteria to other tissues or subjects, or the amelioration of one or more symptoms associated with a bacterial infection. In other embodiments, such terms refer to a reduction in the swelling of one or more joints, organs or tissues, or a reduction in the pain associated with an inflammatory disorder.

[0082] As used herein, the term “UTR” refers to the untranslated region of a mRNA, *i.e.*, the region of the mRNA that is not translated into protein. In a preferred embodiment, the UTR is a 5' UTR, *i.e.*, upstream of the coding region, or a 3' UTR, *i.e.*, downstream of the coding region. In another embodiment, the term UTR corresponds to a reading frame of the mRNA that is not translated. In another embodiment, a UTR contains a fragment of an untranslated region of a mRNA. In a preferred embodiment, a UTR contains one or more regulatory elements that modulate untranslated region-dependent regulation of gene expression.

[0083] As used herein, the term “uORF” refers to an upstream open reading frame that is in the 5' UTR of the main open reading frame, *i.e.*, that encodes a functional protein, of a mRNA.

[0084] As used herein, the term “untranslated region-dependent expression” or “UTR-dependent expression” refers to the regulation of gene expression through untranslated regions at the level of mRNA expression, *i.e.*, after transcription of the gene has begun until the protein or the RNA product(s) encoded by the gene has degraded. In a preferred embodiment, the term “untranslated region-dependent expression” or “UTR-dependent expression” refers to the regulation of mRNA stability or translation. In a more preferred embodiment, the term “untranslated region-dependent expression” refers to the regulation of gene expression through regulatory elements present in an untranslated region(s).

#### 4. DESCRIPTION OF DRAWINGS

[0085] FIGS. 1A-1B: Schematic representation of the VEGF 5'- and 3'-UTRs generated by PCR. A. VEGF 5'UTR was amplified from human genomic DNA by two separate PCR reactions. 5'UTR1, from position 337 to the 3' end plus first 45 nucleotides of VEGF open reading frame, was generated using primers 3 and 4. 5'UTR2, covered from

position 1 to 498, was generated with primers 1 and 2. In the overlap region of 5'UTR1 and 5'UTR2, the unique enzyme site BamH I was used to assemble the full length 5'UTR in the subsequent cloning. B. The full length VEGF 3'UTR was directly amplified from genomic DNA using primers 5 and 6. The two enzyme sites close to 5' end and 3' end of 3'UTR (Bgl II and EcoR I) were used for subsequent cloning.

**[0086]** FIGS. 2A-2C: Identification of VEGF IRES domain in the VEGF mRNA 5'UTR. A. Dual luciferase vector used for mapping IRES function (Grentzmann et al., 1998, RNA 4:479-486). B. Schematic representation of the dicistronic plasmids used for transfection experiments. P2luc/veg5utr1 is the dicistronic plasmid containing the VEGF 5'UTR1, in which nucleotides 337 to 1083 of the VEGF cDNA were fused to the firefly luciferase coding sequence; P2luc/veg5utr-fl was generated by subcloning VEGF 5'UTR2 into the plasmid p2luc/veg5utr1 between Sal I and BamH I; plasmid p2luc/veg5'utr-delta51-476 is derived from p2luc/veg5'utr-fl by removing the Nhe I fragment (nt 51 to 746); plasmid p2luc/veg5utr-delta476-1038 was derived from p2luc/veg5utr-fl by removing the sequence from BamH I site to the 3'end of 5'UTR; plasmid p2luc/veg5utr-delta1-476 was derived from p2luc/veg5utr-fl by removing the sequence from BamH I to the 5'end of 5'UTR. P2luc-e used as negative control in this study. C. The constructs depicted in panel A were transfected into 293T cells in the triplicate format and expression was analyzed by monitoring luciferase activity.

**[0087]** FIGS. 3A-3B: Generation of stable cell lines for cell based high throughput screening ("HTS"). A. Schematic representation of the monocistronic plasmid used in this study for generation of stable cell lines. B. Screening of stable cell lines. The plasmid depicted in panel A was transfected into 293T cells. 48 hours later, the transfected cells were seeded in 96 well plates at 100-500 cells per well and 200 mg/ml hygromycin was added for selection. The culture media plus hygromycin was changed every 3 to 4 days. After 2 weeks of selection, cells were screened under a microscope and single colony wells were expanded for further luciferase assays. The chart in panel shows the luciferase activities for 19 stable clones.

**[0088]** FIG. 4: Side by side comparison of luciferase activities for three stable clones (B9, D3 and H6). For each cell line,  $5 \times 10^5$  cells per well were seeded in 24 well plate. 48 hours later, cells were lysed and assayed for luciferase activities. The luciferase activities were normalized against protein concentration.

**[0089]** FIG. 5: Sustained high expression of luciferase by B9 cells. B9 cells were continuously cultured in vitro for more than 3 months. At the time points indicated,



Luciferase activity was tested with Promega's Bright Glow substrate and normalized against the protein concentration.

[0090] FIGS. 6A-6B: Reporter gene integration in B9 cells. The integration levels of the reporter gene were determined using semi-quantitative PCR. Series diluted plasmid plus 5' + 3' vegf-UTR were included as positive control to make sure the reaction for sample (genomic DNA from B9 cells) was in the linear range, *i.e.*, not saturated. Panel A shows the PCR results for sample and positive control. The PCR band intensity for each reaction is at the bottom of the picture. Panel B shows the PCR standard curve, plotted with PCR band intensity against the amount of positive control plasmid loaded for PCR.

[0091] FIGS. 7A-7B: The 5' UTR of survivin can function as an internal ribosome entry site (IRES). A. Firefly luciferase assays on 293T cells transiently transfected with the survivin expression vectors in the absence of a stem-loop secondary structure. "5' + 3' UTR" represents the survivin expression vector containing the firefly luciferase reporter gene surrounded by both the 5' and 3' untranslated regions of survivin. "5' UTR" represents the survivin expression vector containing the firefly luciferase reporter gene preceded only by the 5' UTR of survivin. "3' UTR" represents the survivin expression vector containing the firefly luciferase reporter gene followed only by the 3' UTR of survivin. "no UTR" represents the survivin expression vector containing the firefly luciferase reporter gene lacking any surrounding untranslated regions of survivin. The survivin expression vectors were transiently transfected into 293T cells in duplicate (represented by the two bars for each construct in the graph) and firefly luciferase activity (measured in quadruplicate) was normalized to total protein concentration in each of the cell lysates. B. As in FIG. 7A, except that the survivin expression vectors containing the stem-loop secondary structure to separate cap-dependent from cap-independent translation were used.

[0092] FIGS. 8A-8C: Expression Vectors. A. Schematic representation of pCMR1, a high-level stable and transient mammalian expression vector designed to randomly integrate into the genome. B. Schematic representation of pMCP1, a high level stable and transient mammalian expression vector designed to site-specifically integrate into the genome of cells genetically engineered to contain the FRT site-specific recombination site via the Flp recombinase (see, *e.g.*, Craig, 1988, Ann. Rev. Genet. 22: 77-105; and Sauer, 1994, Curr. Opin. Biotechnol. 5: 521-527). C. Schematic representation of pCMR2, an episomal mammalian expression vector.

## 5. DETAILED DESCRIPTION OF THE INVENTION

[0093] The present invention provides methods for identifying compounds that modulate the untranslated region-dependent expression of any target gene. In particular, the invention provides simple, rapid and sensitive methods for identifying compounds that modulate untranslated region-dependent expression of a target gene utilizing reporter gene-based constructs comprising one or more mRNA untranslated regions ("UTRs") of the target gene. The reporter gene-based assays described herein can be utilized in a high throughput format to screen libraries of compounds to identify those compounds that modulate untranslated region-dependent expression of a target gene.

[0094] The reporter gene-based assays of the invention reduce the bias introduced by competitive binding assays which require the identification of use of a host cell factor (presumably essential for modulating RNA function) as a binding partner for the target RNA. The reporter gene-based assays of the invention are designed to detect any compound that modulates untranslated region-dependent expression of a target gene under physiologic conditions.

[0095] The reporter gene-based assays may be conducted by contacting a compound with a cell genetically engineered to express a nucleic acid comprising a reporter gene operably linked to one or more untranslated regions (preferably, the 5' and/or 3' UTRs) of a target gene, and measuring the expression of said reporter gene. Alternatively, the reporter gene-based assays may be conducted by contacting a compound with a cell-free translation mixture and a nucleic acid comprising a reporter gene operably linked to one or more untranslated regions of a target gene, and measuring the expression of said reporter gene. The alteration in reporter gene expression relative to a previously determined reference range or a control in such reporter-gene based assays indicates that a particular compound modulates untranslated region-dependent expression of a target gene. In order to exclude the possibility that a particular compound is functioning solely by modulating the expression of a target gene in an untranslated region-independent manner, one or more mutations (*i.e.*, deletions, insertions, or nucleotide substitutions) may be introduced into the untranslated regions operably linked to a reporter gene and the effect on the expression of the reporter gene in a reporter gene-based assay described herein can be determined.

[0096] The compounds identified in the reporter gene-based assays described herein that modulate untranslated region-dependent expression may be tested in *in vitro* assays (*e.g.*, cell-free assays) or *in vivo* assays (*e.g.*, cell-based assays) well-known to one of skill in the art or described herein for the effect of said compounds on the expression of the target gene from which the untranslated regions of the reporter gene construct were derived. Further, the specificity of a particular compound's effect on untranslated region-dependent

expression of one or more other genes (preferably, a plurality of genes) can be determined utilizing assays well-known to one of skill in the art or described herein. In a preferred embodiment, a compound identified utilizing the reporter gene-based assays described herein has a specific effect on the expression of only one gene or a group of genes within the same signaling pathway.

[0097] The structure of the compounds identified in the reporter gene-based assays described herein that modulate untranslated region-dependent expression can be determined utilizing assays well-known to one of skill in the art or described herein. The methods used will depend, in part, on the nature of the library screened. For example, assays or microarrays of compounds, each having an address or identifier, may be deconvoluted, *e.g.*, by cross-referencing the positive sample to an original compound list that was applied to the individual test assays. Alternatively, the structure of the compounds identified herein may be determined using mass spectrometry, nuclear magnetic resonance (“NMR”), X ray crystallography, or vibrational spectroscopy.

[0098] The invention encompasses the use of the compounds identified in accordance with the methods described herein for the modulation (*i.e.*, upregulation or downregulation) of the expression of a target gene. The upregulation or downregulation of a target gene is particularly useful *in vitro* when attempting to produce a protein encoded by said target gene for use as a therapeutic or prophylactic agent, or in experiments conducted to, *e.g.*, identify the function or efficacy of said protein. The invention also encompasses the use of the compounds identified in accordance with the methods described herein for the prevention, treatment or amelioration of a disease or disorder or a symptom thereof. Examples of diseases and disorders which may be prevented, treated or ameliorated utilizing a compound identified in accordance with the invention include, but are not limited to, proliferative disorders, disorders associated with aberrant angiogenesis, inflammatory disorders, infectious diseases, genetic disorders, autoimmune disorders, cardiovascular diseases, and central nervous system disorders. In an embodiment wherein the disease or disorder is an infectious disease, the infectious disease can be caused by a fungal infection, a bacterial infection, a viral infection, or an infection caused by another type of pathogen.

### 5.1. Untranslated Regions

[0099] Any untranslated region may be utilized in the reporter gene constructs described herein. An untranslated gene region(s) may be obtained or derived from a gene from any species, including, but not limited to, plants (*e.g.*, soybean, canola, cotton, wheat, corn, rice, potato, and tomato plants), viruses, bacteria, fungus and animals (including, but not limited to, mammals (primates and non-primates), farm animals (*e.g.*, horses, pigs,

cows, donkeys, etc.), pets (*e.g.*, guinea pigs, cats, and dogs), and humans). Untranslated regions may be obtained and the nucleotide sequence of the untranslated regions determined by any method well-known to one of skill in the art. The nucleotide sequence of an untranslated region for a target gene can be obtained, *e.g.*, from the literature or a database such as GenBank. Alternatively, the nucleotide sequence of the untranslated regions of a target gene may be generated from nucleic acid from a suitable source. If a clone containing a nucleic acid of an untranslated region of a target gene is not available, but the sequence of the untranslated region is known, a nucleic acid of the untranslated region may be chemically synthesized or obtained from a suitable source (*e.g.*, a cDNA library) by PCR amplification. Once the nucleotide sequence of an untranslated region is determined, the nucleotide sequence of the untranslated region may be manipulated using methods well-known in the art for the manipulation of nucleotide sequences, *e.g.*, recombinant DNA techniques, site directed mutagenesis, PCR, *etc.* (see, for example, the techniques described in Sambrook *et al.*, 1990, *Molecular Cloning, A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY and Ausubel *et al.*, eds., 1998, *Current Protocols in Molecular Biology*, John Wiley & Sons, NY, which are both incorporated by reference herein in their entireties), to generate an untranslated region having a different nucleotide sequence.

**[00100]** In one embodiment, an untranslated gene region(s) is obtained or derived from a gene whose expression is associated with or has been linked to the onset, development, progression or severity of a particular disease or disorder. In another embodiment, an untranslated gene region(s) is obtained or derived from a gene whose expression is beneficial to a subject with a particular disease or disorder. Examples of genes from which the untranslated regions may be obtained or derived from include, but are not limited to, cytokines, cytokine receptors, T cell receptors, B cell receptors, co-stimulatory molecules, clotting cascade factors, cyclins, cyclin inhibitors, oncogenes, growth factors, growth factor receptors, tumor suppressors, apoptosis inhibitor proteins, cell adhesion molecules, hormones, GTP-binding proteins, glycoproteins, ion channel receptors, calcium channel pumps, steroid receptors, opioid receptors, sodium channel pumps, heat shock proteins, MHC proteins, and tumor-associated antigens ("TAAs").

**[00101]** Specific examples of genes from which the untranslated regions may be obtained or derived from include, but are not limited, to the gene encoding *abl*, the gene encoding acetyl CoA carboxylase beta ("ACC2"; see, *e.g.*, OMIM accession number 601557, locus link accession number 32), the gene encoding acetylcholinesterase ("ACHE"; see, *e.g.*, OMIM accession number 100740, locus link accession number 43, GenBank

accession number NM 0006 65), the gene encoding actin, alpha cardiac ("ACTC"; see, *e.g.*, OMIM accession number 102540, locus link accession number 70), the gene encoding acyl-CoA dehydrogenase ("ACADVL"; see, *e.g.*, OMIM accession number 201475, locus link accession number 37), the gene encoding adiponectin ("ACRP30"; see, *e.g.*, OMIM accession number 605441, locus link accession number 9370, GenBank accession number NM 0047 97), the gene encoding ADP-ribosylation factor-4 ("ARF4"; see, *e.g.*, OMIM accession number 601177, locus link accession number 378, GenBank accession number NM 0017 ev 25), the gene encoding alpha-glucosidase, the gene encoding Alzheimer's disease amyloid A4 ("APP" or "A4" or "CVAP" or "AD1"; see, *e.g.*, OMIM accession number 104760, locus link accession number 351), the gene encoding angiogenin ("ANG" or "RNASE5"; see, *e.g.*, OMIM accession number 105850, locus link accession number 283, GenBank accession number NM 0011 45), the gene encoding angiopoietin1 ("ANG1"; see, *e.g.*, OMIM accession number 601667, locus link accession number 284), the gene encoding angiopoietin2 ("ANG2"; see, *e.g.*, OMIM accession number 601922, locus link accession number 285), the gene encoding angiostatin, the gene encoding angiotensin 1-converting enzyme ("DCP1"; see, *e.g.*, OMIM accession number 106180, locus link accession number 1636), the gene encoding antigen CD82 ("KAI1"; see, *e.g.*, OMIM accession number 600623, locus link accession number 3732, GenBank accession number NM 0022 31), the gene encoding APC, the gene encoding atrial natriuretic factor, the gene encoding bactericidal/permeability-increasing protein ("BPI"; see, *e.g.*, OMIM accession number 109195, locus link accession number 671, GenBank accession number NM 0017 ev 25), the gene encoding bcl-2, the gene encoding beta-catenin ("CTNNB1"; see, *e.g.*, OMIM accession number 116806, locus link accession number 1499), the gene encoding beta-site APP-cleaving enzyme 2 ("BASE2"; see, *e.g.*, OMIM accession number 605668, locus link accession number 25825, GenBank accession number NM 1389 92), the gene encoding bile salt export pump ("ABCB1 1"; see, *e.g.*, OMIM accession number 603201, locus link accession number 8647), the gene encoding BMP, the gene encoding BDNF, the gene encoding bombesin receptor, the gene encoding brca1, the gene encoding brca2, the gene encoding C1q complement receptor (see, *e.g.*, OMIM accession number 120577, locus link accession number 22918), the gene encoding c-fms, the gene encoding c-myc, the gene encoding calcitonin, the gene encoding calcium-binding protein in macrophages ("MRP14"; see, *e.g.*, OMIM accession number 123886, locus link accession number 6280, GenBank accession number NM 0029 ev 65), the gene encoding calsenilin ("DREAM/CSEN" or "CREAM" or "KCh IP3"; see, *e.g.*, OMIM accession number 604662, locus link accession number 30818, GenBank accession number NM 0134), the gene encoding carnitine o-

palmitoyltransferase ("CPT2"; see, *e.g.*, OMIM accession number 600650, locus link accession number 1376), the gene encoding catechol-o-methyltransferase ("COMT"; see, *e.g.*, OMIM accession number 116790, locus link accession number 1312, GenBank accession number NM 000754, NM 007310), the gene encoding cathepsin K, the gene encoding CD40 ligand ("TNFSF5"; see, *e.g.*, OMIM accession number 300386, locus link accession number 959), the gene encoding cdk4 inhibitor, the gene encoding chemokine (C-C) receptor ("IL13R"; see, *e.g.*, OMIM accession number 601268, locus link accession number 1232), the gene encoding chemokine (C-X3-C) receptor 1 ("CX3CR1"; see, *e.g.*, OMIM accession number 601470, locus link accession number 1524), the gene encoding CLCA homolog ("hCLCA2"; see, *e.g.*, OMIM accession number 604003, locus link accession number 9635, GenBank accession number NM 0065 36), the gene encoding complement decay-accelerating factor ("DAF/CD55"; see, *e.g.*, OMIM accession number 125240, locus link accession number 1604), the gene encoding connective tissue growth factor ("CTGF"; see, *e.g.*, OMIM accession number 121009, locus link accession number 1490), the gene encoding corticotrophin releasing factor, the gene encoding CTLA4, the gene encoding cyclin D1, the gene encoding cyclin E, the gene encoding cyclin T1 (see, *e.g.*, OMIM accession number 602506, locus link accession number 904, GenBank accession number NM 0012 40), the gene encoding cyclin-dependent kinase inhibitor 1A ("p21" or "WAF1" or "CDKN1A" or "Cip1"; see, *e.g.*, OMIM accession number 116899, locus link accession number 1026, GenBank accession number NM 0784 67), the gene encoding cyclin-dependent kinase inhibitor 2A ("CDKN2A"; see, *e.g.*, OMIM accession number 600160, locus link accession number 1029), the gene encoding cystic fibrosis transmembrane conductance regulator ("CFTR"), the gene encoding cytochrome P-450, the gene encoding D-1 dopamine receptor ("DRD1"; see, *e.g.*, OMIM accession number 126449, locus link accession number 1812, GenBank accession number NM 00794, X589987), the gene encoding D-amino-acid oxidase ("DAO"; see, *e.g.*, OMIM accession number 124050, locus link accession number 1610, GenBank accession number NM 0019 17), the gene encoding damage specific DNA binding protein ("DDB1"; see, *e.g.*, OMIM accession number 600045, locus link accession number 1642), the gene encoding DCC, the gene encoding desmoglein 1 ("DSG1"; see, *e.g.*, OMIM accession number 125670, locus link accession number 1828), the gene encoding a dihydrofolate reductase ("DHFR"; see, *e.g.*, OMIM accession number 126060, locus link accession number 1719, GenBank accession number NM 0007 91), the gene encoding a disintegrin and metallo proteinase domain 33 ("ADAM 33"; see, *e.g.*, OMIM accession number 607114, locus link accession number 80332), the gene encoding DNA methyltransferase ("DNMT3b"; see, *e.g.*, OMIM

accession number 602900, locus link accession number 1789), the gene encoding DPP-IV, the gene encoding drebrin-1 dendritic spine protein ("DBN1"; see, *e.g.*, OMIM accession number 126660, locus link accession number 1627, GenBank accession number NM 004395, NM 080881), the gene encoding E-cadherin, the gene encoding effector cell protease receptor ("EPR1"; see, *e.g.*, OMIM accession number 603411, locus link accession number 8475), the gene encoding EGF, the gene encoding EGFR (see, *e.g.*, OMIM accession number 131550, locus link accession number 1956), the gene encoding an EGFR subunit, the gene encoding EIF4BP (see, *e.g.*, OMIM accession number 602223, locus link accession number 1978, GenBank accession number NM 0040 95), the gene encoding EMMPRIN (see, *e.g.*, OMIM accession number 109480, locus link accession number 682, GenBank accession number NM 0017 28), the gene encoding emotakin ATP-binding cassette, sub-family a, member 1 ("ABCA1"; see, *e.g.*, OMIM accession number 600046, locus link accession number 19), the gene encoding endostatin, the gene encoding eotaxin ("CCL11"; see, *e.g.*, OMIM accession number 601156, locus link accession number 6356, GenBank accession number NM 0029 86), the gene encoding erythropoietin ("EPO"; see, *e.g.*, OMIM accession number 133170, locus link accession number 2056, GenBank accession number NM 0007 99), the gene encoding estrogen receptor, the gene encoding factor IX, the gene encoding factor VIII, the gene encoding farnesyl transferase, the gene encoding FGF, the gene encoding FGF1 (see, *e.g.*, OMIM accession number 131220, locus link accession number 2246, GenBank accession number ), the gene encoding FGF2 (see, *e.g.*, OMIM accession number 134920, locus link accession number 2247, GenBank accession number NM 0020 06), the gene encoding FGFR, the gene encoding fibrillin ("FBN1"; see, *e.g.*, OMIM accession number 134797, locus link accession number 2200), the gene encoding FMS-related tyrosine kinase 1 ("FLT1"; see, *e.g.*, OMIM accession number 165070, locus link accession number 2321, GenBank accession number NM 0020 ev 19), the gene encoding forkhead box C2 ("FOXC2"; see, *e.g.*, OMIM accession number 602402, locus link accession number 2303, GenBank accession number NM 0052 51), the gene encoding fos (see, *e.g.*, OMIM accession number 164810, locus link accession number 2353, GenBank accession number NM 0052 52), the gene encoding G-CSF, the gene encoding G-CSF 3 ("CSF3"; see, *e.g.*, OMIM accession number 138970, locus link accession number 1440), the gene encoding a GABA receptor, the gene encoding galanin ("GAL"; see, *e.g.*, OMIM accession number 137035, locus link accession number 2586), the gene encoding gastric inhibitory polypeptide ("GIP"; see, *e.g.*, OMIM accession number 137240, locus link accession number 2695), the gene encoding GDNF, the gene encoding GGF, the gene encoding GGRP, the gene encoding ghrelin ("GHRL"; see, *e.g.*, OMIM

accession number 605353, locus link accession number 51738), the gene encoding gip, the gene encoding glucagon, the gene encoding glucagon receptor ("GCGR"; see, *e.g.*, OMIM accession number 138033, locus link accession number 2642), the gene encoding glucagon-like peptide-1 ("GLP1"; see, *e.g.*, OMIM accession number 138030, locus link accession number 2641), the gene encoding glucokinase ("GCK"; see, *e.g.*, OMIM accession number 138079, locus link accession number 2645, GenBank accession number NM 0001 62), the gene encoding glutamic acid decarboxylase 2 (see, *e.g.*, OMIM accession number 138275), the gene encoding glutamic acid decarboxylase 3 (see, *e.g.*, OMIM accession number 138276), the gene encoding glutamic acid decarboxylase, brain, membrane form (see, *e.g.*, OMIM accession number 138277), the gene encoding glycogen synthase kinase-3A ("GSK-3A"; see, *e.g.*, OMIM accession number 606784, locus link accession number 2931), the gene encoding glycogen synthase kinase-3B ("GSK-3B"; see, *e.g.*, OMIM accession number 605004, locus link accession number 2932), the gene encoding GM-CSF (see, *e.g.*, OMIM accession number 138960, locus link accession number 1437), the gene encoding gonadotropin, the gene encoding gonadotropin releasing hormone, the gene encoding GRO2 oncogene or macrophage inflammatory protein-2-alpha precursor ("CXCL2"; see, *e.g.*, OMIM accession number 139110, locus link accession number 2920), the gene encoding growth hormone releasing factor, the gene encoding growth hormone, the gene encoding gsp, the gene encoding H-ras, the gene encoding heat shock protein ("HSP")-70, the gene encoding heparanase ("HPA"; see, *e.g.*, OMIM accession number 604724, locus link accession number 10855), the gene encoding hepatitis A virus cellular receptor ("HAVCR"; see, *e.g.*, OMIM accession number 606518, locus link accession number 26762), the gene encoding hepatitis B virus X interacting protein ("HBXIP"), the gene encoding hepsin ("HPN"; see, *e.g.*, OMIM accession number 142440, locus link accession number 3249, GenBank accession number NM 0021 51), the gene encoding Her-2 ("ERBB2"; see, *e.g.*, OMIM accession number 164870, locus link accession number 2064), the gene encoding HGF, the gene encoding histone acetyltransferase ("HAT1"; see, *e.g.*, OMIM accession number 603053, locus link accession number 8520, GenBank accession number NM 0036 42), the gene encoding histone deacetylase 1 ("HDAC1"; see, *e.g.*, OMIM accession number 601241, locus link accession number 3065), the gene encoding histone deacetylase 3 ("HDAC3"; see, *e.g.*, OMIM accession number 605166, locus link accession number 8841, GenBank accession number NM 0038 ev 83), the gene encoding HIV Tat Specific Factor 1 ("HTATSF1"; see, *e.g.*, OMIM accession number 300346, locus link accession number 27336), the gene encoding HMG CoA synthetase, the gene encoding HSP-90, the gene encoding huntingtin ("HD"; see, *e.g.*, OMIM accession number 143100,



Tocus link accession number 3064, GenBank accession number NM 0021 11), the gene encoding Hu antigen R ("HUR"; see, *e.g.*, OMIM accession number 603466, locus link accession number 1994, GenBank accession number NM 0014 19), the gene encoding 3-hydroxy-3-methylglutaryl-CoA reductase ("HMGCR"; see, *e.g.*, OMIM accession number 142910, locus link accession number 3156), the gene encoding hypoxia-inducible factor 1 ("HIF-1A"; see, *e.g.*, OMIM accession number 603348, locus link accession number 3091), the gene encoding hypoxia-inducible factor 1-alpha inhibitor ("HIF1AN"; see, *e.g.*, OMIM accession number 606615, locus link accession number 55662), the gene encoding iduronate 2-sulfatase ("IDS"; see, *e.g.*, OMIM accession number 309900, locus link accession number 3423), the gene encoding IGF-1 (see, *e.g.*, OMIM accession number 147440, locus link accession number 3486), the gene encoding IGF-1R (see, *e.g.*, OMIM accession number 147370, locus link accession number 3480, GenBank accession number NM 0008 ev 75), the gene encoding IGF-2, the gene encoding IGF binding protein-2 ("IGFBP2"; see, *e.g.*, OMIM accession number 146731, locus link accession number 3485), the gene encoding IκB kinase ("IKBKB"; see, *e.g.*, OMIM accession number 603258, locus link accession number 3551), the gene encoding inositol polyphosphate phosphatase-like 1 ("SHIP-2"; see, *e.g.*, OMIM accession number 600829, locus link accession number 3636, GenBank accession number NM 0015 67), the gene encoding insulin, the gene encoding interferon inducible protein ("CXCL10 (IP10)"; see, *e.g.*, OMIM accession number 147310, locus link accession number 3627, GenBank accession number NM 0015 65), the gene encoding interferon ("IFN")-α, the gene encoding interferon-α 1/13 precursor, the gene encoding interferon-α 5 precursor ("IFNA5"; see, *e.g.*, OMIM accession number 147565, locus link accession number 3442), the gene encoding interferon-α-16 precursor ("IFNA16"; see, *e.g.*, OMIM accession number 147580, locus link accession number 3449), the gene encoding IFN-β, the gene encoding IFN-β 1 ("IFNB1"; see, *e.g.*, OMIM accession number 147640, locus link accession number 3456), the gene encoding IFN-γ (see, *e.g.*, OMIM accession number 147440, locus link accession number 3479), the gene encoding insulin receptor ("INSR"; see, *e.g.*, OMIM accession number 147670, locus link accession number 3643, GenBank accession number NM 0002 08), the gene encoding interleukin-1 b ("IL1B"; see, *e.g.*, OMIM accession number 147720, locus link accession number 3553), the gene encoding interleukin-2 ("IL-2"; see, *e.g.*, OMIM accession number 147680, locus link accession number 3558), the gene encoding interleukin-3 ("IL-3"), the gene encoding interleukin-4 ("IL-4"; see, *e.g.*, OMIM accession number 147780, locus link accession number 3565, GenBank accession number NM 0005 89), the gene encoding interleukin-4 receptor ("IL4R"; see, *e.g.*, OMIM accession number 147781, locus link accession number

3566, GenBank accession number NM 0004 18), the gene encoding interleukin-5 ("IL-5"), the gene encoding interleukin-6 ("IL-6"; see, *e.g.*, OMIM accession number 147620, locus link accession number 3569), the gene encoding interleukin-7 ("IL-7"), the gene encoding interleukin-8 ("IL-8"; see, *e.g.*, OMIM accession number 146930, locus link accession number 3576), the gene encoding interleukin-9 ("IL-9"; see, *e.g.*, OMIM accession number 146931, locus link accession number 3578), the gene encoding interleukin-10 ("IL-10"; see, *e.g.*, OMIM accession number 124092, locus link accession number 3586, GenBank accession number NM 0005 72), the gene encoding interleukin-12 ("IL-12"), the gene encoding interleukin-12 beta chain precursor ("IL12B"; see, *e.g.*, OMIM accession number 161561, locus link accession number 3593), the gene encoding interleukin-13 ("IL-13"; see, *e.g.*, OMIM accession number 147683, locus link accession number 3596, GenBank accession number NM 0021 88), the gene encoding interleukin-15 ("IL-15"), the gene encoding interleukin-17F ("ML1"; see, *e.g.*, OMIM accession number 606496, locus link accession number 11274), the gene encoding interleukin-18 ("IL-18"; see, *e.g.*, OMIM accession number 600953, locus link accession number 3606), the gene encoding INI1/hSNF5 (see, *e.g.*, OMIM accession number 601607, locus link accession number 6598), the gene encoding jun, the gene encoding kallikrein 6 ("KLK6"; see, *e.g.*, OMIM accession number 602652, locus link accession number 5653, GenBank accession number NM 0027 74), the gene encoding KGF, the gene encoding ki-ras, the gene encoding kit ligand, stem cell factor ("KITLG (SCF)"; see, *e.g.*, OMIM accession number 184745, locus link accession number 4254, GenBank accession number NM 0008 99), the gene encoding klotho ("KL"; see, *e.g.*, OMIM accession number 604824, locus link accession number 9365, GenBank accession number NM 0047 95), the gene encoding L-myc, the gene encoding large tumor suppressor ("LATS1"; see, *e.g.*, OMIM accession number 603473, locus link accession number 9113, GenBank accession number NM 0046 ev 90), the gene encoding LDL receptor ("LDLR"; see, *e.g.*, OMIM accession number 606945, locus link accession number 3949, GenBank accession number NM 0005 27), the gene encoding leptin ("LEP"; see, *e.g.*, OMIM accession number 164160, locus link accession number 3952, GenBank accession number NM 0002 30), the gene encoding leptin receptor ("LEPR"; see, *e.g.*, OMIM accession number 601007, locus link accession number 3953), the gene encoding leucine amino peptidase-3 ("LAP3"; see, *e.g.*, OMIM accession number 606832, locus link accession number 51056), the gene encoding leukemia inhibitory factor ("LIF"; see, *e.g.*, OMIM accession number 159540, locus link accession number 3976), the gene encoding leukemia inhibitory factor receptor ("LIFR"; see, *e.g.*, OMIM accession number 151443, locus link accession number 3977), the gene encoding linker for activation

of T cells ("LAT"; see, *e.g.*, OMIM accession number 602354, locus link accession number 27040), the gene encoding livin (see, *e.g.*, OMIM accession number 605737, locus link accession number 79444, GenBank accession number NM 1393 ev 17), the gene encoding luteinizing hormone, the gene encoding luteinizing hormone releasing hormone, the gene encoding macrophage migration inhibitory factor ("MIF"; see, *e.g.*, OMIM accession number 153620, locus link accession number 4282, GenBank accession number NM 0024 15), the gene encoding major histocompatibility complex class I chain-related gene A ("MICA"; see, *e.g.*, OMIM accession number 600169, locus link accession number 4276, GenBank accession number NM 0002 47), the gene encoding major histocompatibility complex class I chain-related gene B ("MICB"; see, *e.g.*, OMIM accession number 602436, locus link accession number 4277, GenBank accession number NM 0059 31), the gene encoding matrix metalloproteinase 9 ("MMP9"; see, *e.g.*, OMIM accession number 120361, locus link accession number 4318), the gene encoding matrix metalloproteinase 12 ("MMP12"; see, *e.g.*, OMIM accession number 601046, locus link accession number 4321), the gene encoding max interacting protein 1 ("MXI1"; see, *e.g.*, OMIM accession number 600020, locus link accession number 4601), the gene encoding MCC, the gene encoding MDM2 (see, *e.g.*, OMIM accession number 164785, locus link accession number 4193, GenBank accession number NM 0023 92), the gene encoding METH-1, the gene encoding METH-2, the gene encoding methyl-CpG-binding endonuclease ("MBD4"; see, *e.g.*, OMIM accession number 603574, locus link accession number 8930, GenBank accession number NM 0039 ev 25), the gene encoding monoamine oxidase-A ("MAOA"; see, *e.g.*, OMIM accession number 309850, locus link accession number 4128, GenBank accession number NM 0002 ev 40), the gene encoding monoamine oxidase-B ("MAOB"; see, *e.g.*, OMIM accession number 309860, locus link accession number 4129), the gene encoding monocyte chemotactic protein 1 ("MCP1"; see, *e.g.*, OMIM accession number 158105, locus link accession number 6347), the gene encoding mos, the gene encoding MTS1, the gene encoding myc, the gene encoding myotrophin, the gene encoding N-acetyltransferase, the gene encoding N-cadherin, the gene encoding N-methyl D-aspartate ("NMDA") receptor, the gene encoding NAD(P)-dependent steroid dehydrogenase ("NSDHL"; see, *e.g.*, OMIM accession number 300275, locus link accession number 50814), the gene encoding natural resistance-associated macrophage protein ("NRAMP"; see, *e.g.*, OMIM accession number 600266, locus link accession number 6556), the gene encoding neural cell adhesion molecule 1 ("NCAM1"; see, *e.g.*, OMIM accession number 116930, locus link accession number 4684), the gene encoding neuron growth associated protein 43 ("GAP-43"; see, *e.g.*, OMIM accession number 162060, locus link accession number 2596),

the gene encoding NF1, the gene encoding NF2, the gene encoding NGF, the gene encoding a NGFR subunit, the gene encoding nm23, the gene encoding nuclear factor of kappa light polypeptide gene enhancer in B-cells 1 ("NFKB1"; see, *e.g.*, OMIM accession number 164011, locus link accession number 4790), the gene encoding OSM, the gene encoding osteopontin ("OPN"; see, *e.g.*, OMIM accession number 166490, locus link accession number 6696), the gene encoding P-glycoprotein-1 ("PGY1"; see, *e.g.*, OMIM accession number 171050, locus link accession number 5243, GenBank accession number NM 0009 27), the gene encoding p38 MAP kinase ("p38" or "MAPK14"; see, *e.g.*, OMIM accession number 600289, locus link accession number 1432), the gene encoding p53, the gene encoding p300/CBP associated factor("PCAF"; see, *e.g.*, OMIM accession number 602303, locus link accession number 8850), the gene encoding parathyroid hormone, the gene encoding PDGF, the gene encoding PDGF, beta chain ("PDGF2"; see, *e.g.*, OMIM accession number 190040, locus link accession number 5155), the gene encoding a PDGFR subunit, the gene encoding peroxin-1 ("PEX1"; see, *e.g.*, OMIM accession number 602136, locus link accession number 5189), the gene encoding peroxisome assembly factor-2 ("PEX6"; see, *e.g.*, OMIM accession number 601498, locus link accession number 5190), the gene encoding peroxisome proliferator-activated receptor-gamma ("PPARg"; see, *e.g.*, OMIM accession number 601487, locus link accession number 5468), the gene encoding phenylalanine hydroxylase, the gene encoding phosphodiesterase, the gene encoding human phosphotyrosyl-protein phosphatase ("PTP-1B"; see, *e.g.*, OMIM accession number 176885, locus link accession number 5770, GenBank accession number NM 0028 27), the gene encoding placental growth factor ("PGF"; see, *e.g.*, OMIM accession number 601121, locus link accession number 5228, GenBank accession number NM 0026 ev 32), the gene encoding plasminogen activator inhibitor protein ("PAI1"; see, *e.g.*, OMIM accession number 173360, locus link accession number 5054), the gene encoding pleiotrophin ("PTN"; see, *e.g.*, OMIM accession number 162095, locus link accession number 5764), the gene encoding poly(rC) binding protein 2 ("PCBP2"; see, *e.g.*, OMIM accession number 601210, locus link accession number 5094), the gene encoding progranulin ("PCDGF" or "GRN"; see, *e.g.*, OMIM accession number 138945, locus link accession number 2896), the gene encoding prolactin ("PRL"; see, *e.g.*, OMIM accession number 176760, locus link accession number 5617, GenBank accession number NM 0009 48), the gene encoding proliferating cell nuclear antigen ("PCNA"; see, *e.g.*, OMIM accession number 176740, locus link accession number 5111), the gene encoding protein kinase B/Akt ("AKT1"; see, *e.g.*, OMIM accession number 164730, locus link accession number 207), the gene encoding protein kinase C, gamma ("PKCg"; see, *e.g.*, OMIM accession number 176980,

locus link accession number 5582), the gene encoding protein-tyrosine phosphatase, 4A, 3 ("PTP4A3"; see, *e.g.*, OMIM accession number 606449, locus link accession number 11156, GenBank accession number NM 0326 11), the gene encoding psoriasin ("PSOR1"; see, *e.g.*, OMIM accession number 600353, locus link accession number 6278, GenBank accession number NM 0029 63), the gene encoding ras, the gene encoding resistin ("Fizz3"; see, *e.g.*, OMIM accession number 605565, locus link accession number 56729, GenBank accession number NM 0204 15), the gene encoding retinoblastoma ("Rb"; see, *e.g.*, OMIM accession number 180200, locus link accession number 5925, GenBank accession number NM 0003 21), the gene encoding retinoblastoma 1 ("Rb1"), the gene encoding retinoblastoma-binding protein 1-like 1 ("RBBP1L1"; see, *e.g.*, locus link accession number 51742), the gene encoding 5-a reductase, the gene encoding ribonuclease/angiogenin inhibitor ("RNH"; see, *e.g.*, OMIM accession number 173320, locus link accession number 6050), the gene encoding S100 calcium-binding protein A8 ("MRP8"; see, *e.g.*, OMIM accession number 123885, locus link accession number 6279, GenBank accession number NM 0029 ev 64), the gene encoding signal transducer and activator of transcription 6 ("STAT6"; see, *e.g.*, OMIM accession number 601512, locus link accession number 6778), the gene encoding soluble-type polypeptide FZD4S ("FZD4S"; see, *e.g.*, OMIM accession number 604579, locus link accession number 8322), the gene encoding somatotrophin or somatotropin, the gene encoding src (see, *e.g.*, OMIM accession number 190090, locus link accession number 6714, GenBank accession number NM 0054 ev 17), the gene encoding survivin, the gene encoding T-cell lymphoma invasion and metastasis 1 ("TIAM1"; see, *e.g.*, OMIM accession number 600687, locus link accession number 7074), the gene encoding TEK tyrosine kinase ("TIE2"; see, *e.g.*, OMIM accession number 600221, locus link accession number 7010), the gene encoding telomerase, the gene encoding TGF- $\beta$ , the gene encoding TGF- $\beta$ 1 (see, *e.g.*, OMIM accession number 190180, locus link accession number 7040), the gene encoding thrombomodulin ("THBD" or "THRM"; see, *e.g.*, OMIM accession number 188040, locus link accession number 7056), the gene encoding thrombopoietin ("THPO" or "TPO"; see, *e.g.*, OMIM accession number 600044, locus link accession number 7066), the gene encoding human trisosephosphate isomerase ("TPI1"; see, *e.g.*, OMIM accession number 109450, locus link accession number 7167), the gene encoding thyroid hormone, the gene encoding thyroid stimulating hormone, the gene encoding tissue factor, the gene encoding tissue inhibitor of metalloprotease 1 ("TIMP1"; see, *e.g.*, OMIM accession number 305370, locus link accession number 7076), the gene encoding tissue inhibitor of metalloprotease 2 ("TIMP2"; see, *e.g.*, OMIM accession number 188825, locus link accession number 7077, GenBank accession number NM 0032

55), the gene encoding tissue inhibitor of metalloprotease 4 (“TIMP4”; see, *e.g.*, OMIM accession number 601915, locus link accession number 7079, GenBank accession number NM 0032 56), the gene encoding TNF- $\alpha$  (see, *e.g.*, OMIM accession number 191160, locus link accession number 7124), the gene encoding troponin T (“TnT”), the gene encoding uncoupling protein 2 (“UCP2”; see, *e.g.*, OMIM accession number 601693, locus link accession number 7351, GenBank accession number NM 0033 55), the gene encoding urokinase plasminogen activator (“uPA”; see, *e.g.*, OMIM accession number 191840, locus link accession number 5328), the gene encoding utrophin (“UTRN”; see, *e.g.*, OMIM accession number 128240, locus link accession number 7402), the gene encoding v-myc myelocytomatosis viral oncogene homolog (“c-MYC”; see, *e.g.*, OMIM accession number 190080, locus link accession number 4609), the gene encoding vanilloid receptor subunit 1 (“VR1”; see, *e.g.*, OMIM accession number 602076, locus link accession number 7442, GenBank accession number NM 0187 ev 27, NM 08 0704, NM 0807 05, NM 0807 06), the gene encoding vascular endothelial growth factor (“VEGF”), the gene encoding virion infectivity factor (“VIF”), and the gene encoding VLA-4.

**[00102]** In a specific embodiment, an untranslated region is obtained or derived from the gene encoding Her-2. In another embodiment, an untranslated region is not obtained or derived from the gene encoding Her-2.

**[00103]** In one embodiment, an untranslated region is obtained or derived from the gene encoding VEGF. In another embodiment, an untranslated region is not obtained or derived from the gene encoding VEGF.

**[00104]** The untranslated regions may be obtained or derived from the genome of any virus utilizing any method well-known to one of skill in the art. The nucleotide sequence of an untranslated region for a genome of a virus can be obtained, *e.g.*, from the literature or a database such as GenBank. Examples of viruses from which the untranslated regions may be obtained or derived from include, but are not limited to, retroviruses (*e.g.*, human immunodeficiency virus (“HIV”) and human T cell leukemia virus (“HTLV”), herpesviruses (*e.g.*, herpes simplex virus, epstein barr virus and varicella zoster virus), reoviruses (*e.g.*, reovirus and rotavirus), picornaviruses (*e.g.*, poliovirus, rhinovirus and hepatitis A virus), togaviruses (*e.g.*, rubella virus), orthomyxovirus (*e.g.*, influenza virus), paramyxoviruses (*e.g.*, measles virus, mumps virus, respiratory syncytical virus and parainfluenza virus), filoviruses (*e.g.*, ebola virus and Marburg virus), rhabdoviruses (*e.g.*, rabies virus), coronaviruses (*e.g.*, coronavirus), rhinoviruses, hepatitis B virus, and hepatitis C virus.

[00105] The untranslated regions may be obtained or derived from the genome of any bacteria utilizing any method well-known to one of skill in the art. The nucleotide sequence of an untranslated region for a genome of a bacteria can be obtained, *e.g.*, from the literature or a database such as GenBank. Examples of bacteria from which the untranslated regions may be obtained or derived from include, but are not limited to, the *Aquaspirillum* family, *Azospirillum* family, *Azotobacteraceae* family, *Bacteroidaceae* family, *Bartonella* species, *Bdellovibrio* family, *Campylobacter* species, *Chlamydia* species (*e.g.*, *Chlamydia pneumoniae*), clostridium, *Enterobacteriaceae* family (*e.g.*, *Citrobacter* species, *Edwardsiella*, *Enterobacter aerogenes*, *Erwinia* species, *Escherichia coli*, *Hafnia* species, *Klebsiella* species, *Morganella* species, *Proteus vulgaris*, *Providencia*, *Salmonella* species, *Serratia marcescens*, and *Shigella flexneri*), *Gardinella* family, *Haemophilus influenzae*, *Halobacteriaceae* family, *Helicobacter* family, *Legionellaceae* family, *Listeria* species, *Methylococcaceae* family, mycobacteria (*e.g.*, *Mycobacterium tuberculosis*), *Neisseriaceae* family, *Oceanospirillum* family, *Pasteurellaceae* family, *Pneumococcus* species, *Pseudomonas* species, *Rhizobiaceae* family, *Spirillum* family, *Spirosomaceae* family, *Staphylococcus* (*e.g.*, methicillin resistant *Staphylococcus aureus* and *Staphylococcus pyrogenes*), *Streptococcus* (*e.g.*, *Streptococcus enteritidis*, *Streptococcus fasciae*, and *Streptococcus pneumoniae*), *VampirovibrHelicobacter* family, and *Vampirovibrio* family.

[00106] The untranslated regions may be obtained or derived from the genome of any fungus utilizing any method well-known to one of skill in the art. The nucleotide sequence of an untranslated region for a genome of a fungus can be obtained, *e.g.*, from the literature or a database such as GenBank. Examples of fungus from which the untranslated regions may be obtained or derived from include, but are not limited to, *Absidia* species (*e.g.*, *Absidia corymbifera* and *Absidia ramosa*), *Aspergillus* species, (*e.g.*, *Aspergillus flavus*, *Aspergillus fumigatus*, *Aspergillus nidulans*, *Aspergillus niger*, and *Aspergillus terreus*), *Basidiobolus ranarum*, *Blastomyces dermatitidis*, *Candida* species (*e.g.*, *Candida albicans*, *Candida glabrata*, *Candida kerr*, *Candida krusei*, *Candida parapsilosis*, *Candida pseudotropicalis*, *Candida quilliermondii*, *Candida rugosa*, *Candida stellatoidea*, and *Candida tropicalis*), *Coccidioides immitis*, *Conidiobolus* species, *Cryptococcus neoforms*, *Cunninghamella* species, dermatophytes, *Histoplasma capsulatum*, *Microsporium gypseum*, *Mucor pusillus*, *Paracoccidioides brasiliensis*, *Pseudallescheria boydii*, *Rhinosporidium seeberi*, *Pneumocystis carinii*, *Rhizopus* species (*e.g.*, *Rhizopus arrhizus*, *Rhizopus oryzae*, and *Rhizopus microsporus*), *Saccharomyces* species, *Sporothrix schenckii*, zygomycetes, and classes such as *Zygomycetes*, *Ascomycetes*, the *Basidiomycetes*, *Deuteromycetes*, and *Oomycetes*.

[00107] The untranslated regions may be obtained or derived from the genome of any plant utilizing any method well-known to one of skill in the art. The nucleotide sequence of an untranslated region for a genome of a plant can be obtained, *e.g.*, from the literature or a database such as GenBank, EMBL, DDBJ, rice genome database, cotton genome database and maize genome database. Examples of plants from which the untranslated regions may be obtained or derived from include, but are not limited to, soybean, canola, cotton, corn, wheat, rice, tomato, and potato. Specific examples of plant genes from which an untranslated region may be obtained or derived from include, but are not limited to, triose phosphate isomerase, fructose 1,6-bisphosphate adolase, fructose 1,6-bisphosphate, fructose 6-phosphate 2-kinase, phosphoglucosomerase, pyrophosphate-dependent fructose-6-phosphate phosphotransferase, vacuolar H<sup>+</sup> translocating-pyrophosphate, invertase, sucrose synthase, hexokinase, fructokinase, NDP-kinase, glucose-6-phosphate 1-dehydrogenase, phosphoglucosomutase, UDP-glucose pyrophosphorylase, glutenin genes, cis-prenyltransferase, lipoxygenase, and soybean vestitone reductase (see, *e.g.*, U.S. Patent Application Publication No. 2003/0135870 A1 and U.S. Patent Nos. 6,638,525, 6,645,747, 6,627,797, and 6,617,493, which are incorporated herein by reference in its entirety).

[00108] In particular, a 5' UTR of a target gene, a 3' UTR of a target gene, or a 5' UTR and a 3' UTR of a target gene may be utilized in a reporter construct. In a specific embodiment, a 5' UTR of a target gene with a stable hairpin secondary structure is utilized in a reporter construct. In another specific embodiment, a reporter gene in the reporter construct contains an intron. In a preferred embodiment, a 5' UTR and a 3' UTR of a target gene are utilized in a reporter construct. In another preferred embodiment, a 5' UTR and a 3' UTR of a target gene and an intron-containing reporter gene are utilized in a reporter construct.

#### **5.1.1. Elements of Untranslated Regions**

[00109] Any element of an untranslated region(s) of a target gene may be utilized in the reporter gene constructs described herein. Elements of an untranslated region(s) may be obtained and the nucleotide sequence of the elements determined by any method well-known to one of skill in the art. The nucleotide sequence of an element of an untranslated region for a target gene can be obtained, *e.g.*, from the literature or a database such as GenBank. Alternatively, the nucleotide sequence of an element of an untranslated region of a target gene may be generated from nucleic acid from a suitable source. If a clone containing a nucleic acid of an element of an untranslated region of a target gene is not available, but the sequence of the element is known, a nucleic acid of the element may be chemically synthesized or obtained from a suitable source (*e.g.*, a cDNA library) by PCR



amplification. Once the nucleotide sequence of an element is determined, the nucleotide sequence of the element may be manipulated using methods well-known in the art for the manipulation of nucleotide sequences, *e.g.*, recombinant DNA techniques, site directed mutagenesis, PCR, etc. (see, for example, the techniques described in Sambrook et al., 1990, Molecular Cloning, A Laboratory Manual, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY and Ausubel *et al.*, eds., 1998, Current Protocols in Molecular Biology, John Wiley & Sons, NY, which are both incorporated by reference herein in their entireties), to generate an element having a different nucleic acid sequence.

[00110] In one embodiment, an element(s) of an untranslated region comprises the full-length sequence of a UTR, *e.g.*, the 5' UTR or the 3' UTR. In a specific embodiment, an element(s) of an untranslated region that has been shown or has been suggested to be involved in the regulation of mRNA stability and/or translation is utilized in the reporter constructs described herein. Examples of elements of an untranslated region which may be utilized in the reporter constructs described herein include, but are not limited to, an IRE, IRES, uORF, MSL-2, G quartet element, 5'-terminal oligopyrimidine tract ("TOP"), ARE, SECIS, histone stem loop, CPE, nanos translational control element, APP, TGE/DRE, BRE, and a 15-LOX-DICE.

#### 5.1.1.1. Iron Response Element

[00111] The maintenance of cellular iron homeostasis occurs at the level of mRNA stability and translation. Two components of this regulatory system have been defined: a *cis*-acting mRNA sequence/structure motif called an iron-responsive element ("IRE") and a specific *trans*-acting cytoplasmic binding protein, referred to herein as IRE-binding protein ("IRE-BP") (reviewed in, *e.g.*, Mikulits et al., 1999, Mutat Res. 437(3):219-30; Harrison & Arosio, 1996, Biochim Biophys Acta. 1275(3):161-203; Kuhn & Hentze, 1992, J Inorg Biochem. 1992, 47(3-4):183-95; and Harford & Klausner, 1990, Enzyme 44(1-4):28-41, the disclosures of which are hereby incorporated by reference in their entireties). Iron scarcity induces binding of IRE-BPs to a single IRE in the 5' UTR of ferritin, eALAS, aconitase, erythroid 5-aminolevulinic acid synthase, and SDHb mRNAs, which specifically suppresses translation initiation. Simultaneous interaction of IRE-BPs with multiple IREs in the 3' UTR of transferrin receptor mRNA selectively causes its stabilization. The pattern is reverted under iron overload: IRE-BP mRNA binding affinity is reduced, which results in efficient protein synthesis of target transcripts harboring IREs in the 5' UTR and rapid degradation of transferrin mRNA. Any gene containing an IRE including, but not limited to, the IREs described in the references cited above, can be used in the present invention to identify compounds that modulate untranslated region-dependent gene expression.

### 5.1.1.2. Internal Ribosome Entry Site

[00112] The internal ribosome entry site (“IRES”) is one of the better characterized 5’ UTR-based *cis*-acting elements of post-transcriptional gene expression control. IRESes facilitate cap-independent translation initiation by recruiting ribosomes directly to the 5’ UTR of the mRNA. IRESes are commonly located in the 3’ region of 5’ UTR and are, as recent work has established, frequently composed of several discrete sequences. IRESes do not share significant primary structure homology, but do form distinct RNA tertiary structures. Some IRESes contain sequences complementary to 18S RNA and therefore may form stable complexes with 40S ribosomal subunit and initiate assembly of translationally competent complex. A classic example of an “RNA-only” IRES is the internal ribosome entry site from Hepatitis C virus. However, most known IRESes require protein co-factors for activity. More than 10 IRES trans-acting factors (“ITAFs”) have been identified so far. In addition, all canonical translation initiation factors, with the sole exception of 5’ end cap-binding eIF4E, have been shown to participate in IRES-mediated translation initiation (reviewed in Vagner et al., 2001, EMBO Reports 2:893 and Translational Control of Gene Expression, Sonenberg, Hershey, and Mathews, eds., 2000, CSHL Press, the disclosures of which are incorporated by reference in their entirety).

[00113] IRES were first identified in picornaviruses (see, *e.g.*, Pettetier & Sonenberg, 1988, Nature, 334:320-325). The 5’ UTRs of all picornaviruses are long and mediate translational initiation by directly recruiting and binding ribosomes, thereby circumventing the initial cap-binding step. Although IRES elements are frequently found in viral mRNAs, they are rarely found in non-viral mRNAs. The non-viral mRNAs shown to contain functional IRES elements in their respective 5’ UTRs include those encoding immunoglobulin heavy chain binding protein (“BiP”) (see, *e.g.*, Macejak et al., 1991, Nature, 353:90-4); *Drosophila* Antennapedia (see, *e.g.*, Oh et al., 1992, Genes Dev 6:1643-53) and Ultrabithorax (see, *e.g.*, Ye et al., 1997, Mol. Cell Biol. 17:1714-21); fibroblast growth factor 2 (see, *e.g.*, Vagner et al., 1995, Mol. Cell Biol. 15:35-44); initiation factor eIF4G (see, *e.g.*, Gan et al., 1998, J. Biol. Chem. 273:5006-12); proto-oncogene c-myc (see, *e.g.*, Nanbru et al., 1995, J. Biol. Chem. 272:32061-6 and Stoneley, 1998, Oncogene 16:423-8); vascular endothelial growth factor (“VEGF”) (see, *e.g.*, Stein et al., 1998, Mol. Cell Biol. 18:3112-9), and X-linked inhibitor of apoptosis protein (“XIAP”) (see, *e.g.*, U.S. Patent Nos. 6,159,709 and 6,171,821), the disclosures of which are incorporated by reference in their entirety. Any gene containing an IRES including, but not limited to, the IRESes described in the references cited above, can be used in the present invention to identify compounds that modulate untranslated region-dependent gene expression.

### 5.1.1.3. Male Specific Lethal Element

[00114] Male-specific expression of the protein male-specific-lethal 2 (“MSL-2”) controls dosage compensation in *Drosophila*. MSL-2 protein is not produced in females and sequences in both the 5’ and 3’ UTRs are important for this sex-specific regulation because *msl-2* gene expression is inhibited in females by Sex-lethal (“SXL”), an RNA binding protein known to regulate pre-mRNA splicing. An intron present in the 5’ untranslated region of *msl-2* mRNA contains putative SXL binding sites and is retained in female flies. The *msl-2* pre-mRNA is alternatively spliced in a Sex-lethal-dependent fashion (see, *e.g.*, Gebauer et al., 1998, RNA 4(2):142-50 and Bashaw & Baker, 1995, Development 121(10):3245-58, the disclosures of which are hereby incorporated by reference in their entireties). Any gene containing an MSL-2 element including, but not limited to, the MSL-2 elements described in the references cited above, can be used in the present invention to identify compounds that modulate untranslated region-dependent gene expression.

### 5.1.1.4. G-quartet Element

[00115] A symmetrical structure of two tetrads of guanosine base pairs connected by three loops is commonly referred to as a “G-quartet”, “G-quadruplex” or “G-tetraplex” structure (see, *e.g.*, Wang et al., 1993, Biochemistry 32:1899-1904; Macaya et al., 1993, Proc. Natl. Acad. Sci. 90:3745-3749; Schultze et al., 1994, J. Mol. Biol. 235:1532-1547; and Kelly et al., 1996, J. Mol. Biol. 256:417-422, the disclosures of which are incorporated by reference in their entireties). A G-quartet element was first identified as a conserved consensus sequence GGNTGGN<sub>2-5</sub>GGNTGG (SEQ ID NO: 1), which was present in single-stranded DNA aptamers that bind thrombin and inhibited thrombin-catalyzed fibrin-clot formation (see, *e.g.*, Bock et al., 1992, Nature 355:564-566, the disclosure of which is incorporated by reference in its entirety). A similar sequence in which the G-quartet structure is maintained when the length of the oligonucleotide between the G pairs is increased has been identified (see, *e.g.*, Dias et al., 1994, J. Am. Chem. Soc. 116:4479-4480, the disclosure of which is incorporated by reference in its entirety).

[0100] A G-quartet element has been identified in mRNAs associated with fragile X mental retardation syndrome (reviewed in, *e.g.*, Bardoni & Mandel, 2002, Curr Opin Genet Dev 12(3):284-93, the disclosure of which is incorporated by reference in its entirety). The fragile X mental retardation syndrome is caused by large methylated expansions of a CGG repeat in the FMR1 gene that lead to the loss of expression of FMRP, an RNA-binding protein. FMRP is proposed to act as a regulator of mRNA transport or translation that plays

a role in synaptic maturation and function and has been shown to interact preferentially with mRNAs containing a G quartet structure.

[0101] G-quartet oligonucleotides can have the sequence  $\text{GGN}_x\text{GGN}_y\text{GGN}_z\text{GG}$  (SEQ ID NO: 2), wherein x, y and z indicate a variable number of nucleotides (see, *e.g.*, U.S. Patent No. 5,691,145, the disclosure of which is incorporated by reference in its entirety). While x, y and z are each typically at least about 2, preferably about 2-10, these segments may be longer if desired. The regions of variable sequence (*i.e.*,  $\text{N}_x\text{N}_y\text{N}_z$ ) are not critical in the present invention and can be varied in length and sequence without disrupting the characteristic G-quartet structure. As a general rule, the variable N sequences should not be self-complementary and should not contain G residues which would result in alternative G-quartet structures within the molecule. Representative G-quartet oligonucleotides are 15-20 nucleotides in length, but G-quartet oligonucleotides of any length which conform to the general formula  $\text{GGN}_x\text{GGN}_y\text{GGN}_z\text{GG}$  (SEQ ID NO: 3) are also suitable. The G-quartet oligonucleotide is typically about 14-30 nucleotides in length. Any gene containing a G-quartet element including, but not limited to, the G-quartet elements described in the references cited above, can be used in the present invention to identify compounds that modulate untranslated region-dependent gene expression.

#### 5.1.1.5. 5'-terminal Oligopyrimidine Tract

[0102] Translation control can be mediated by a terminal oligopyrimidine element ("TOP") present in the 5' untranslated region of ribosomal protein-encoding mRNAs. TOP elements adopt a specific secondary structure that prevents ribosome-binding and translation-initiation of ribosomal protein-encoding mRNAs. However, binding of cellular nucleic acid binding protein ("CNBP") or La proteins to the TOP hairpin structure abolishes the TOP-mediated transcription block and induces ribosomal protein production (see, *e.g.*, Schlatter & Fussenegger, 2003, *Biotechnol Bioeng* 81(1):1-12; Zhu et al., 2001, *Biochim Biophys Acta* 1521(1-3):19-29; and Crosio et al., 2000, *Nucleic Acids Res.* 28(15):2927-34, the disclosures of which are incorporated by reference in their entireties).

[0103] The immunosuppressant rapamycin selectively suppresses the translation of mRNAs containing a TOP tract adjacent to the cap structure. *Trans*-acting factors, some of which are regulated by rapamycin-responsive signaling pathways, that bind to the 5' untranslated region of TOP mRNAs may be involved in selective translational repression (see, *e.g.*, Kakegawa et al., 2002, *Arch Biochem Biophys* 402(1):77-83, the disclosure of which is incorporated by reference in its entirety). Any gene containing a TOP element including, but not limited to, the TOP elements described in the references cited above, can

be used in the present invention to identify compounds that modulate untranslated region-dependent gene expression.

#### 5.1.1.6. Adenylate Uridylate-rich Element

[0104] AU-rich elements ("AREs") are the most extensively studied 3' UTR-based regulatory signals. AREs are the primary determinant of mRNA stability and one of the key determinants of mRNA translation initiation efficiency.

[0105] A typical ARE is 50 to 150 nt long and contains 3 to 6 copies of AU<sub>3</sub>A pentamer embedded in a generally A/U-enriched RNA region. The AU<sub>3</sub>A pentamers can be scattered within the region or can stagger or even overlap (Chen et al., 1995, Trends Biol. Sciences 20:465, the disclosure of which is incorporated by reference in its entirety). One or several AU<sub>3</sub>A pentamers can be replaced by expanded versions such as an AU<sub>4</sub>A hexamer or AU<sub>5</sub>A heptamer (see, *e.g.*, Wilkund et al., 2002, J. Biol. Chem. 277:40462 and Tholanikunnel & Malborn, 1997, J. Biol. Chem. 272:11471, the disclosures of which are incorporated by reference in their entirety). Single copies of the AU<sub>n</sub>A (where n = 3, 4, or 5) elements placed in a random sequence context are inactive. The minimal active ARE has been determined to have the sequence U<sub>2</sub>AU<sub>n</sub>A(U/A)(U/A) (where n = 3, 4, or 5) (see, *e.g.*, Worthington et al., 2002, J Biol Chem, 277:48558-64) the disclosure of which is incorporated by reference in its entirety). The activity of certain AU-rich elements in promoting mRNA degradation is enhanced in the presence of distal uridine-rich sequences. These U-rich elements do not affect mRNA stability when present alone and thus that have been termed "ARE enhancers" (see, *e.g.*, Chen et al., 1994, Mol. Cell. Biol. 14:416, the disclosure of which is incorporated by reference in its entirety).

[0106] Most AREs function in mRNA decay regulation and translation initiation regulation by interacting with specific ARE-binding proteins ("AUBPs"). There are at least 14 known cellular proteins that bind to AU-rich elements. AUBP functional properties determine ARE involvement in one or both pathways. For example, ELAV/HuR binding to c-fos ARE inhibits c-fos mRNA decay (see, *e.g.*, Brennan & Steitz, 2001, Cell Mol Life Sci. 58:266), association of tristetraprolin with TNF $\alpha$  ARE dramatically enhances TNF $\alpha$  mRNA hydrolysis (see, *e.g.*, Carballo et al., 1998, Science 281:1001), whereas interaction of TIA-1 with the TNF $\alpha$  ARE does not alter the TNF $\alpha$  mRNA stability but inhibits TNF $\alpha$  translation (see, *e.g.*, Piecyk et al., 2000, EMBO J. 19:4154).

[0107] Since AREs are clearly important in biological systems, including but not limited to a number of the early response genes that regulate cell proliferation and responses to exogenous agents, the identification of compounds that bind to one or more of the ARE clusters and potentially modulate the stability and translation of the target RNA can

potentially be of value as a therapeutic. Any gene containing an ARE including, but not limited to, the AREs described in the references cited above, can be used in the present invention to identify compounds that modulate untranslated region-dependent gene expression.

#### 5.1.1.7. Selenocysteine Insertion Sequence

[0108] Selenium is an essential micronutrient that is now known to be incorporated as selenocysteine in a number of selenoproteins, glutathione peroxidase being the prototypical example. Selenocysteine is specifically encoded by the UGA codon, and inserted in peptide chains by a cotranslational mechanism that is able to override the normal function of UGA as a termination codon. In eukaryotes, efficient selenocysteine incorporation at UGA codons requires a cellular protein factor and a *cis*-acting structural signal usually located in the mRNA 3' untranslated region, consisting of a selenocysteine insertion sequence ("SECIS") in a characteristic stem-loop structure (see, *e.g.*, Peterlin et al., 1993, "Tat Trans-Activator" In Human Retroviruses; Cullen, Ed.; Oxford University Press: New York; pp. 75-100; Le & Maizel, 1989, J. Theor. Biol. 138:495-510; and reviewed in Hubert et al., 1996, Biochimie 78(7):590-6, the disclosures of which are incorporated by reference in their entireties). The required protein factor is presumed to be present in certain cells types that express selenoproteins, such as liver cells, lymphocytes, macrophages, thrombocytes, and other blood cells. In such cell types, the presence of a SECIS element in an mRNA is necessary and sufficient for in-frame UGA codons to be translated as selenocysteine.

[0109] A SECIS element is usually UAAAG, although other SECIS elements have been identified or variants have been constructed (see, *e.g.*, U.S. Patent Nos. 6,303,295, 5,849,520, and 5,700,660, the disclosures of which are incorporated by reference in their entireties). Any gene containing a SECIS element including, but not limited to, the SECIS elements described in the references cited above, can be used in the present invention to identify compounds that modulate untranslated region-dependent gene expression.

#### 5.1.1.8. Histone Stem Loop

[0110] Replication-dependent histone mRNAs end with a conserved 26-nucleotide sequence that contains a 16-nucleotide stem-loop, *i.e.*, the histone stem loop, instead of a poly(A) tail. Formation of the 3' end of histone mRNA occurs by endonucleolytic cleavage of pre-mRNA releasing the mature mRNA from the chromatin template. Cleavage requires several *trans*-acting factors, including a protein, the stem-loop binding protein, which binds the 26-nucleotide sequence, and a small nuclear RNP, U7 snRNP (reviewed in, *e.g.*,

Dominski & Marzluff, 1999, Gene 239(1):1-14, the disclosure of which is incorporated by reference in its entirety).

[0111] Sequences of histone stem loops have been described in U.S. Patent Nos. 6,476,208; 6,455,280; 6,399,373; 6,346,381; 6,335,170; 6,331,396; 6,265,546; 6,265,167; 5,990,298; 5,908,779 and 5,843,770, the disclosures of which are incorporated by reference in their entireties. Any gene containing a histone stem loop including, but not limited to, the histone stem loops described in the references cited above, can be used in the present invention to identify compounds that modulate untranslated region-dependent gene expression.

#### **5.1.1.9. Cytoplasmic Polyadenylation Element**

[0112] Maturation-specific polyadenylation in *Xenopus* oocytes depends on the presence of a U-rich cytoplasmic polyadenylation element ("CPE") close to the 3' end of the RNA. RNAs that lack CPEs appear to be deadenylated by default when meiosis resumes. This default program also applies to maturing mouse oocytes (see, *e.g.*, Paynton & Bachvarova, 1994, Mol Reprod Dev 37(2):172-80, the disclosure of which is incorporated by reference in its entirety). CPEs have been identified in Wee1 protein tyrosine kinase mRNA (see, *e.g.*, Charlesworth et al., 2000, Dev Biol 227(2):706-19, the disclosure of which is incorporated by reference in its entirety), cyclin B1 mRNA (see, *e.g.*, Tay et al., 2000, Dev Biol 221(1):1-9 and Barkoff et al., 2000, Dev Biol 220(1):97-109, the disclosures of which are incorporated by reference in their entireties), and *Xenopus* Id3 mRNA (see, *e.g.*, Afouda et al., 1999, Mech Dev 88(1):15-31, the disclosure of which is incorporated by reference in its entirety).

[0113] A *Xenopus* oocyte CPE binding protein ("CPEB") binds the CPE and stimulates polyadenylation. CPEB is essential for the cytoplasmic polyadenylation of B4 RNA, G10, c-mos, cdk2, cyclins A1, B1 and B2 mRNAs which suggests that this protein is required for polyadenylation of most RNAs during oocyte maturation (see, *e.g.*, Stebbins-Boaz et al., 1996, EMBO J 15(10):2582-92, the disclosure of which is incorporated by reference in its entirety). Any gene containing a CPE including, but not limited to, the CPEs described in the references cited above, can be used in the present invention to identify compounds that modulate untranslated region-dependent gene expression.

#### **5.1.1.10. Nanos Translational Control Element**

[0114] The nanos translational control element is a discrete translational control element within the nanos 3' untranslated region that acts independently of the localization signal to mediate translational repression of unlocalized nanos RNA (see, *e.g.*, Clark et al.,

2002, Development 129(14):3325-34; Clark et al., 2000, Curr Biol 10(20):1311-4; Cruce et al., 2000, Mol Cell 5(3):457-67; Bergsten & Gavis, 1999, Development 126(4):659-69; Dahanukar & Wharton, 1996, Genes Dev (20):2610-20; and Gavis et al., 1996, Development 122(9):2791-800, the disclosures of which are incorporated by reference in their entireties).

[0115] During *Drosophila* embryogenesis, the Smaug protein represses translation of the nanos protein through an interaction with the nanos translational control element (see, *e.g.*, Green et al., 2002, Biochem Biophys Res Commun 297(5):1085-8, the disclosure of which is incorporated by reference in its entirety). Any gene containing a nanos translational control element including, but not limited to, the nanos translational control elements described in the references cited above, can be used in the present invention to identify compounds that modulate untranslated region-dependent gene expression.

#### **5.1.1.11. Amyloid Precursor Protein Element**

[0116] In one embodiment, the amyloid precursor protein element ("APP" element) refers to a novel iron-responsive element within the 5' untranslated region of the Alzheimer's amyloid precursor protein ("APP") transcript (+51 to +94 from the 5'-cap site) (see, *e.g.*, Rogers et al., 2002, J Biol Chem 277(47):45518-28). The APP mRNA IRE is located immediately upstream of an interleukin-1 responsive acute box domain (+101 to +146). The APP 5' UTR conferred translation was selectively down-regulated in response to intracellular iron chelation.

[0117] In another embodiment, the APP element refers to a 29 base instability element in the 3' UTR of the amyloid precursor protein involved in mRNA stability (see, *e.g.*, Westmark & Malter, 2001, Brain Res Mol Brain Res 90(2):193-201; Rajagopalan & Malter, 2000, J Neurochem 74(1):52-9; Amara et al., 1999, Brain Res Mol Brain Res 71(1):42-9; and Zaidi & Malter, 1995, J Biol Chem 270(29):17292-8, the disclosures of which are incorporated by reference in their entireties). Any gene containing a APP element including, but not limited to, the APP elements described in the references cited above, can be used in the present invention to identify compounds that modulate untranslated region-dependent gene expression.

#### **5.1.1.12. Translation Regulation Element**

[0118] Negative translational control elements in 3' UTRs regulate pattern formation, cell fate, and sex determination in a variety of organisms. *tra-2* mRNA in *Caenorhabditis elegans* is required for female development but must be repressed to permit spermatogenesis in hermaphrodites. Translational repression of *tra-2* mRNA in *C. elegans*



is mediated by tandemly repeated elements in its 3' UTR; these elements are called TGEs (for tra-2 and GLI element) (see, *e.g.*, Thompson et al., 2000, Mol Cell Biol 20(6):2129-37; Haag & Kimble, 2000, Genetics 155(1):105-16; and Jan et al., 1997, EMBO J 16(20):6301-13, the disclosures of which are incorporated by reference in their entireties). Any gene containing a TGE including, but not limited to, the TGEs described in the references cited above, can be used in the present invention to identify compounds that modulate untranslated region-dependent gene expression.

#### **5.1.1.13. Direct Repeat Element**

[0119] The direct repeat element ("DRE") is one control element in the 3' UTR of the tra-2 mRNA that causes repression of tra-2, *i.e.*, inhibits translation of tra-2 mRNA, which is responsible for the onset of hermaphrodite spermatogenesis in *C. elegans* (see, *e.g.*, Goodwin et al., 1993, Cell 75:329-339, the disclosure of which is incorporated by reference in its entirety). Three germline-specific regulators have been identified that mediate DRE regulation by the tra-2 3' UTR. These include DRFQ2/GLD-1, a protein that specifically binds the DRE (see, *e.g.*, Goodwin et al., 1993, Cell 75:329-339) and controls tra-2 translation (see, *e.g.*, Jan et al. 1999, EMBO J. 18:258-269); FOG-2, a protein that binds GLD-1 and is required for the onset of hermaphrodite spermatogenesis (see, *e.g.*, Schedl & Kimble, 1988, Genetics 119:43-61); and laf-1, a gene that has not yet been identified at the molecular level (see, *e.g.*, Goodwin et al., 1997, Development 124:749-758), the disclosures of which are incorporated by reference in their entireties. Any gene containing a DRE including, but not limited to, the DREs described in the references cited above, can be used in the present invention to identify compounds that modulate untranslated region-dependent gene expression.

#### **5.1.1.14. Bruno Response Element**

[0120] The Bruno Response Element ("BRE"), is located in the 3' untranslated region (UTR) of oskar mRNA (see, *e.g.*, Castagnetti et al., 2000, Development 127(5):1063-8, the disclosure of which is incorporated by reference in its entirety). The coupled regulation of oskar mRNA localization and translation in time and space is critical for correct anteroposterior patterning of the *Drosophila* embryo. Localization-dependent translation of oskar mRNA, a mechanism whereby oskar RNA localized at the posterior of the oocyte is selectively translated and the unlocalized RNA remains in a translationally repressed state, ensures that Oskar activity is present exclusively at the posterior pole. Genetic experiments indicate that translational repression involves the binding of Bruno protein to multiple sites, the BREs, in the 3' untranslated region of oskar mRNA. Any gene containing a BRE

including, but not limited to, the BREs described in the references cited above, can be used in the present invention to identify compounds that modulate untranslated region-dependent gene expression.

#### **5.1.1.15. 15-lipoxygenase Differentiation Control Element**

[0121] The translation of 15-lipoxygenase (“LOX”) mRNA in erythroid precursor cells and of the L2 mRNA of human papilloma virus type 16 (HPV-16) in squamous epithelial cells is silenced when either of these cells is immature and is activated in maturing cells by unknown mechanisms. It has been shown that hnRNP K and the c-Src kinase specifically interact with each other, leading to c-Src activation and tyrosine phosphorylation of hnRNP K in vivo and in vitro. c-Src-mediated phosphorylation reversibly inhibits the binding of hnRNP K to the differentiation control element (“DICE”) of the LOX mRNA 3’ untranslated region in vitro and specifically derepresses the translation of DICE-bearing mRNAs in vivo (see, *e.g.*, Ostareck-Lederer et al., 2002, Mol Cell Biol 22(13):4535-43, the disclosure of which is incorporated by reference in its entirety).

[0122] Cytidine-rich 15-lipoxygenase differentiation control element (“15-LOX-DICE”) is a multifunctional *cis*-acting element found in the 3’ untranslated region of numerous eukaryotic mRNAs. It binds KH domain proteins of the type hnRNP E and K, thus mediating mRNA stabilization and translational control. Translational silencing is caused by formation of a simple binary complex between DICE and recombinant hnRNP E1. Electromobility shift assays and sucrose gradient centrifugation demonstrate that rabbit 15-LOX-DICE, which is composed of ten subunits of the sequence (CCCCPuCCCUCUUCCCCAAG, SEQ ID NO: 4), is able to bind up to ten molecules of hnRNP E1 (see, *e.g.*, Reimann et al., 2002, J Mol Biol 315(5):965-74 and Thiele et al., 1999, Adv Exp Med Biol 447:45-61, the disclosures of which are incorporated by reference in their entireties). Any gene containing a 15-LOX-DICE including, but not limited to, the 15-LOX-DICES described in the references cited above, can be used in the present invention to identify compounds that modulate untranslated region-dependent gene expression.

#### **5.2. Reporter Gene Constructs, Transfected Cells, and Cell-Free Extracts**

[0123] The invention provides for specific vectors containing a reporter gene flanked by one or more UTRs of a target gene and host cells transfected with the vectors. The invention also provides for the *in vitro* translation of a reporter gene flanked by one or more UTRs of a target gene. Techniques for practicing this specific aspect of this invention will employ, unless otherwise indicated, conventional techniques of molecular biology,

microbiology, and recombinant DNA manipulation and production, which are routinely practiced by one of skill in the art. See, *e.g.*, Sambrook, 1989, *Molecular Cloning, A Laboratory Manual*, Second Edition; DNA Cloning, Volumes I and II (Glover, Ed. 1985); Oligonucleotide Synthesis (Gait, Ed. 1984); Nucleic Acid Hybridization (Hames & Higgins, Eds. 1984); Transcription and Translation (Hames & Higgins, Eds. 1984); Animal Cell Culture (Freshney, Ed. 1986); Immobilized Cells and Enzymes (IRL Press, 1986); Perbal, *A Practical Guide to Molecular Cloning* (1984); Gene Transfer Vectors for Mammalian Cells (Miller & Calos, Eds. 1987, Cold Spring Harbor Laboratory); Methods in Enzymology, Volumes 154 and 155 (Wu & Grossman, and Wu, Eds., respectively), (Mayer & Walker, Eds., 1987); Immunochemical Methods in Cell and Molecular Biology (Academic Press, London, Scopes, 1987), Expression of Proteins in Mammalian Cells Using Vaccinia Viral Vectors in *Current Protocols in Molecular Biology*, Volume 2 (Ausubel et al., Eds., 1991).

#### 5.2.1. Reporter Genes

[0124] Any reporter gene well-known to one of skill in the art may be used in reporter gene constructs to ascertain the effect of a compound on untranslated region-dependent expression of a target gene. Reporter genes refer to a nucleotide sequence encoding a protein that is readily detectable either by its presence or activity. Reporter genes may be obtained and the nucleotide sequence of the reporter gene determined by any method well-known to one of skill in the art. The nucleotide sequence of a reporter gene can be obtained, *e.g.*, from the literature or a database such as GenBank. Alternatively, a polynucleotide encoding a reporter gene may be generated from nucleic acid from a suitable source. If a clone containing a nucleic acid encoding a particular reporter gene is not available, but the sequence of the reporter gene is known, a nucleic acid encoding the reporter gene may be chemically synthesized or obtained from a suitable source (*e.g.*, a cDNA library, or a cDNA library generated from, or nucleic acid, preferably poly A<sup>+</sup> RNA, isolated from, any tissue or cells expressing the reporter gene) by PCR amplification. Once the nucleotide sequence of a reporter gene is determined, the nucleotide sequence of the reporter gene may be manipulated using methods well-known in the art for the manipulation of nucleotide sequences, *e.g.*, recombinant DNA techniques, site directed mutagenesis, PCR, (see, for example, the techniques described in Sambrook et al., 1990, *Molecular Cloning, A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY and Ausubel *et al.*, eds., 1998, *Current Protocols in Molecular Biology*, John Wiley & Sons, NY, which are both incorporated by reference herein in their

entireties), to generate reporter genes having a different amino acid sequence, for example to create amino acid substitutions, deletions, and/or insertions.

[0125] Examples of reporter genes include, but are not limited to, luciferase (*e.g.*, firefly luciferase, renilla luciferase, and click beetle luciferase), green fluorescent protein (“GFP”) (*e.g.*, green fluorescent protein, yellow fluorescent protein, red fluorescent protein, cyan fluorescent protein, and blue fluorescent protein), beta-galactosidase (“b-gal”), beta-glucoronidase, beta-lactamase, chloramphenicol acetyltransferase (“CAT”), and alkaline phosphatase (“AP”). In a preferred embodiment, a reporter gene utilized in the reporter constructs is easily assayed and has an activity which is not normally found in the cell or organism of interest.

#### 5.2.1.1. Luciferase

[0126] Luciferases are enzymes that emit light in the presence of oxygen and a substrate (luciferin) and which have been used for real-time, low-light imaging of gene expression in cell cultures, individual cells, whole organisms, and transgenic organisms (reviewed by Greer & Szalay, 2002, *Luminescence* 17(1):43-74).

[0127] As used herein, the term “luciferase” is intended to embrace all luciferases, or recombinant enzymes derived from luciferases which have luciferase activity. The luciferase genes from fireflies have been well characterized, for example, from the *Photinus* and *Luciola* species (see, *e.g.*, International Patent Publication No. WO 95/25798 for *Photinus pyralis*, European Patent Application No. EP 0 524 448 for *Luciola cruciata* and *Luciola lateralis*, and Devine et al., 1993, *Biochim. Biophys. Acta* 1173(2):121-132 for *Luciola mingrelica*). Other eucaryotic luciferase genes include, but are not limited to, the click beetle (*Photinus plagiophthalmus*, see, *e.g.*, Wood et al., 1989, *Science* 244:700-702), the sea panzy (*Renilla reniformis*, see, *e.g.*, Lorenz et al., 1991, *Proc Natl Acad Sci U S A* 88(10):4438-4442), and the glow worm (*Lampyrus noctiluca*, see *e.g.*, Sula-Newby et al., 1996, *Biochem J.* 313:761-767). The click beetle is unusual in that different members of the species emit bioluminescence of different colors, which emit light at 546 nm (green), 560 nm (yellow-green), 578 nm (yellow) and 593 nm (orange) (see, *e.g.*, U.S. Patent Nos. 6,475,719; 6,342,379; and 6,217,847, the disclosures of which are incorporated by reference in their entireties). Bacterial luciferin-luciferase systems include, but are not limited to, the bacterial lux genes of terrestrial *Photorhabdus luminescens* (see, *e.g.*, Manukhov et al., 2000, *Genetika* 36(3):322-30) and marine bacteria *Vibrio fischeri* and *Vibrio harveyi* (see, *e.g.*, Miyamoto et al., 1988, *J Biol Chem.* 263(26):13393-9, and Cohn et al., 1983, *Proc Natl Acad Sci USA.*, 80(1):120-3, respectively). The luciferases encompassed by the present

invention also includes the mutant luciferases described in U.S. Patent No. 6,265,177 to Squirrell et al., which is hereby incorporated by reference in its entirety.

[0128] In a preferred embodiment, the luciferase is a firefly luciferase, a renilla luciferase, or a click beetle luciferase, as described in any one of the references listed *supra*, the disclosures of which are incorporated by reference in their entireties.

#### **5.2.1.2. Green Fluorescent Protein**

[0129] Green fluorescent protein ("GFP") is a 238 amino acid protein with amino acid residues 65 to 67 involved in the formation of the chromophore which does not require additional substrates or cofactors to fluoresce (see, *e.g.*, Prasher et al., 1992, Gene 111:229-233; Yang et al., 1996, Nature Biotechnol. 14:1252-1256; and Cody et al., 1993, Biochemistry 32:1212-1218).

[0130] As used herein, the term "green fluorescent protein" or "GFP" is intended to embrace all GFPs (including the various forms of GFPs which exhibit colors other than green), or recombinant enzymes derived from GFPs which have GFP activity. In a preferred embodiment, GFP includes green fluorescent protein, yellow fluorescent protein, red fluorescent protein, cyan fluorescent protein, and blue fluorescent protein. The native gene for GFP was cloned from the bioluminescent jellyfish *Aequorea victoria* (see, *e.g.*, Morin et al., 1972, J. Cell Physiol. 77:313-318). Wild type GFP has a major excitation peak at 395 nm and a minor excitation peak at 470 nm. The absorption peak at 470 nm allows the monitoring of GFP levels using standard fluorescein isothiocyanate (FITC) filter sets. Mutants of the GFP gene have been found useful to enhance expression and to modify excitation and fluorescence. For example, mutant GFPs with alanine, glycine, isoleucine, or threonine substituted for serine at position 65 result in mutant GFPs with shifts in excitation maxima and greater fluorescence than wild type protein when excited at 488 nm (see, *e.g.*, Heim et al., 1995, Nature 373:663-664; U.S. Patent No. 5,625,048; Delagrave et al., 1995, Biotechnology 13:151-154; Cormack et al., 1996, Gene 173:33-38; and Cramer et al., 1996, Nature Biotechnol. 14:315-319). The ability to excite GFP at 488 nm permits the use of GFP with standard fluorescence activated cell sorting ("FACS") equipment. In another embodiment, GFPs are isolated from organisms other than the jellyfish, such as, but not limited to, the sea pansy, *Renilla reriformis*.

[0131] Techniques for labeling cells with GFP in general are described in U.S. Patent Nos. 5,491,084 and 5,804,387, which are incorporated by reference in their entireties; Chalfie et al., 1994, Science 263:802-805; Heim et al., 1994, Proc. Natl. Acad. Sci. USA 91:12501-12504; Morise et al., 1974, Biochemistry 13:2656-2662; Ward et al., 1980, Photochem. Photobiol. 31:611-615; Rizzuto et al., 1995, Curr. Biology 5:635-642; and

Kaether & Gerdes, 1995, FEBS Lett 369:267-271. The expression of GFPs in *E. coli* and *C. elegans* are described in U.S. Patent No. 6,251,384 to Tan et al., which is incorporated by reference in its entirety. The expression of GFP in plant cells is discussed in Hu & Cheng, 1995, FEBS Lett 369:331-33, and GFP expression in *Drosophila* is described in Davis et al., 1995, Dev. Biology 170:726-729.

#### **5.2.1.3. Beta-galactosidase**

[0132] Beta galactosidase (“b-gal”) is an enzyme that catalyzes the hydrolysis of b-galactosides, including lactose, and the galactoside analogs o-nitrophenyl-b-D-galactopyranoside (“ONPG”) and chlorophenol red-b-D-galactopyranoside (“CPRG”) (see, e.g., Nielsen et al., 1983 Proc Natl Acad Sci USA 80(17):5198-5202; Eustice et al., 1991, Biotechniques 11:739-742; and Henderson et al., 1986, Clin. Chem. 32:1637-1641). The b-gal gene functions well as a reporter gene because the protein product is extremely stable, resistant to proteolytic degradation in cellular lysates, and easily assayed. When ONPG is used as the substrate, b-gal activity can be quantitated with a spectrophotometer or microplate reader.

[0133] As used herein, the term “beta galactosidase” or “b-gal” is intended to embrace all b-gals, including *lacZ* gene products, or recombinant enzymes derived from b-gals which have b-gal activity. The b-gal gene functions well as a reporter gene because the protein product is extremely stable, resistant to proteolytic degradation in cellular lysates, and easily assayed. In an embodiment where ONPG is the substrate, b-gal activity can be quantitated with a spectrophotometer or microplate reader to determine the amount of ONPG converted at 420 nm. In an embodiment when CPRG is the substrate, b-gal activity can be quantitated with a spectrophotometer or microplate reader to determine the amount of CPRG converted at 570 to 595 nm. In yet another embodiment, the b-gal activity can be visually ascertained by plating bacterial cells transformed with a b-gal construct onto plates containing Xgal and IPTG. Bacterial colonies that are dark blue indicate the presence of high b-gal activity and colonies that are varying shades of blue indicate varying levels of b-gal activity.

#### **5.2.1.4. Beta-glucuronidase**

[0134] Beta-glucuronidase (“GUS”) catalyzes the hydrolysis of a very wide variety of b-glucuronides, and, with much lower efficiency, hydrolyzes some b-galacturonides. GUS is very stable, will tolerate many detergents and widely varying ionic conditions, has no cofactors, nor any ionic requirements, can be assayed at any physiological pH, with an

optimum between 5.0 and 7.8, and is reasonably resistant to thermal inactivation (see, *e.g.*, U.S. Patent No. 5,268,463, which is incorporated by reference in its entirety).

[0135] In one embodiment, the GUS is derived from the *Escherichia coli* b-glucuronidase gene. In alternate embodiments of the invention, the b-glucuronidase encoding nucleic acid is homologous to the *E. coli* b-glucuronidase gene and/or may be derived from another organism or species.

[0136] GUS activity can be assayed either by fluorescence or spectrometry, or any other method described in U.S. Patent No. 5,268,463, the disclosure of which is incorporated by reference in its entirety. For a fluorescent assay, 4-trifluoromethylumbelliferyl b-D-glucuronide is a very sensitive substrate for GUS. The fluorescence maximum is close to 500 nm--bluish green, where very few plant compounds fluoresce or absorb. 4-trifluoromethylumbelliferyl b-D-glucuronide also fluoresces much more strongly near neutral pH, allowing continuous assays to be performed more readily than with MUG. 4-trifluoromethylumbelliferyl b-D-glucuronide can be used as a fluorescent indicator *in vivo*. The spectrophotometric assay is very straightforward and moderately sensitive (Jefferson et al., 1986, Proc. Natl. Acad. Sci. USA 86:8447-8451). A preferred substrate for spectrophotometric measurement is p-nitrophenyl b-D-glucuronide, which when cleaved by GUS releases the chromophore p-nitrophenol. At a pH greater than its pK<sub>a</sub> (around 7.15) the ionized chromophore absorbs light at 400-420 nm, giving a yellow color.

#### 5.2.1.5. Beta-lactamase

[0137] Beta-lactamases are nearly optimal enzymes in respect to their almost diffusion-controlled catalysis of b-lactam hydrolysis, making them suited to the task of an intracellular reporter enzyme (see, *e.g.*, Christensen et al., 1990, Biochem. J. 266: 853-861). They cleave the b-lactam ring of b-lactam antibiotics, such as penicillins and cephalosporins, generating new charged moieties in the process (see, *e.g.*, O'Callaghan et al., 1968, Antimicrob. Agents. Chemother. 8: 57-63 and Stratton, 1988, J. Antimicrob. Chemother. 22, Suppl. A: 23-35). A large number of b-lactamases have been isolated and characterized, all of which would be suitable for use in accordance with the present invention (see, *e.g.*, Richmond & Sykes, 1978, Adv. Microb. Physiol. 9:31-88 and Ambler, 1980, Phil. Trans. R. Soc. Lond. [Ser.B.] 289: 321-331, the disclosures of which are incorporated by reference in their entireties).

[0138] The coding region of an exemplary b-lactamase employed has been described in U.S. Patent No. 6,472,205, Kadonaga et al., 1984, J. Biol. Chem. 259: 2149-2154, and Sutcliffe, 1978, Proc. Natl. Acad. Sci. USA 75: 3737-3741, the disclosures of which are incorporated by reference in their entireties. As would be readily apparent to those skilled

in the field, this and other comparable sequences for peptides having b-lactamase activity would be equally suitable for use in accordance with the present invention. The combination of a fluorogenic substrate described in U.S. Patent Nos. 6,472,205, 5,955,604, and 5,741,657, the disclosures of which are incorporated by reference in their entireties, and a suitable b-lactamase can be employed in a wide variety of different assay systems, such as are described in U.S. Patent No. 4,740,459, which is hereby incorporated by reference in its entirety.

#### **5.2.1.6. Chloramphenicol Acetyltransferase**

[0139] Chloramphenicol acetyl transferase (“CAT”) is commonly used as a reporter gene in mammalian cell systems because mammalian cells do not have detectable levels of CAT activity. The assay for CAT involves incubating cellular extracts with radiolabeled chloramphenicol and appropriate co-factors, separating the starting materials from the product by, for example, thin layer chromatography (“TLC”), followed by scintillation counting (see, *e.g.*, U.S. Patent No. 5,726,041, which is hereby incorporated by reference in its entirety).

[0140] As used herein, the term “chloramphenicol acetyltransferase” or “CAT” is intended to embrace all CATs, or recombinant enzymes derived from CAT which have CAT activity. While it is preferable that a reporter system which does not require cell processing, radioisotopes, and chromatographic separations would be more amenable to high through-put screening, CAT as a reporter gene may be preferable in situations when stability of the reporter gene is important. For example, the CAT reporter protein has an *in vivo* half life of about 50 hours, which is advantageous when an accumulative versus a dynamic change type of result is desired.

#### **5.2.1.7. Secreted Alkaline Phosphatase**

[0141] The secreted alkaline phosphatase (“SEAP”) enzyme is a truncated form of alkaline phosphatase, in which the cleavage of the transmembrane domain of the protein allows it to be secreted from the cells into the surrounding media. In a preferred embodiment, the alkaline phosphatase is isolated from human placenta.

[0142] As used herein, the term “secreted alkaline phosphatase” or “SEAP” is intended to embrace all SEAP or recombinant enzymes derived from SEAP which have alkaline phosphatase activity. SEAP activity can be detected by a variety of methods including, but not limited to, measurement of catalysis of a fluorescent substrate, immunoprecipitation, HPLC, and radiometric detection. The luminescent method is preferred due to its increased sensitivity over calorimetric detection methods. The advantages of using SEAP is that a



cell lysis step is not required since the SEAP protein is secreted out of the cell, which facilitates the automation of sampling and assay procedures. A cell-based assay using SEAP for use in cell-based assessment of inhibitors of the Hepatitis C virus protease is described in U.S. Patent No. 6,280,940 to Potts et al. which is hereby incorporated by reference in its entirety.

### 5.2.2. Reporter Gene Constructs

[0143] The invention provides reporter gene constructs for use in the reporter gene-based assays described herein for the identification of compounds that modulate untranslated region-dependent expression of a target gene. The reporter gene constructs of the invention comprise one or more reporter genes fused to one or more untranslated regions. For example, specific RNA sequences, RNA structural motifs, and/or RNA structural elements that are known or suspected to modulate untranslated region-dependent expression of a target gene may be fused to the reporter gene.

[0144] The present invention provides for a reporter gene flanked by one or more untranslated regions (*e.g.*, the 5' UTR, 3' UTR, or both the 5' UTR and 3' UTR of the target gene). The present invention also provides for a reporter gene flanked by one or more UTRs of a target gene, said UTRs containing one or more mutations (*e.g.*, one or more substitutions, deletions and/or additions). In a preferred embodiment, the reporter gene is flanked by both 5' and 3' UTRs so that compounds that interfere with an interaction between the 5' and 3' UTRs can be identified. In another preferred embodiment, a stable hairpin secondary structure is inserted into the UTR, preferably the 5' UTR of the target gene. For example, in cases where the 5' UTR possesses IRES activity, the addition of a stable hairpin secondary structure in the 5' UTR can be used to separate cap-dependent from cap-independent translation (see, *e.g.*, Muhlrads et al., 1995, *Mol. Cell. Biol.* 15(4):2145-56, the disclosure of which is incorporated by reference in its entirety). In another embodiment, an intron is inserted into a UTR (preferably, the 5' UTR) or at the 5' end of an ORF of a target gene. For example, but not by limitation, in cases where an RNA possesses instability elements, an intron, *e.g.*, the human elongation factor one alpha (EF-1 alpha) first intron, can be cloned into a UTR (preferably, the 5' UTR) or a 5' end of the ORF to increase expression (see, *e.g.*, Kim et al., 2002, *J Biotechnol* 93(2):183-7, the disclosure of which is incorporated by reference in its entirety). In a preferred embodiment, both a stable hairpin secondary structure and an intron are added to the reporter gene construct. In a more preferred embodiment, the stable hairpin secondary structure is cloned into the 5' UTR and the intron is added at the 5' end of the ORF of the reporter gene.

[0145] The reporter gene can be positioned such that the translation of that reporter gene is dependent upon the mode of translation initiation, such as, but not limited to, cap-dependent translation or cap-independent translation (*i.e.*, translation via an internal ribosome entry site). Alternatively, where the UTR contains an upstream open reading frame, the reporter gene can be positioned such that the reporter protein is translated only in the presence of a compound that shifts the reading frame of the UTR so that the formerly untranslated open reading frame is then translated.

[0146] The reporter gene constructs can be monocistronic or multicistronic. A multicistronic reporter gene construct may encode 2, 3, 4, 5, 6, 7, 8, 9, 10 or more, or in the range of 2-5, 5-10 or 10-20 reporter genes. For example, a dicistronic reporter gene construct comprising in the following order a promoter, a first reporter gene, a 5' UTR of a target gene, a second reporter gene and optionally, a 3' UTR of a target gene. In such a reporter construct, the transcription of both reporter genes is driven by the promoter, whereas the translation of the mRNA from the first reporter gene is by a cap-dependent scanning mechanism and the translation of the mRNA from the second reporter gene is by a cap-independent mechanism by an IRES. The IRES-dependent translation of the mRNA of the second reporter gene can be normalized against cap-dependent translation.

### **5.2.3. Expression of Reporter Gene Constructs in Cells**

#### **5.2.3.1. Vectors**

[0147] The nucleotide sequence coding for a reporter gene can be inserted into an appropriate expression vector, *i.e.*, a vector which contains the necessary elements for the transcription and translation of the inserted protein-coding sequence. The necessary transcriptional and translational signals can also be supplied by the target gene or the reporter gene. A variety of host-vector systems may be utilized to express the reporter gene. These include, but are not limited to, mammalian cell systems infected with virus (*e.g.*, vaccinia virus, adenovirus, *etc.*); insect cell systems infected with virus (*e.g.*, baculovirus); microorganisms such as yeast containing yeast vectors, or bacteria transformed with bacteriophage, DNA, plasmid DNA, or cosmid DNA; and stable cell lines generated by transformation using a selectable marker. The expression elements of vectors vary in their strengths and specificities. Depending on the host-vector system utilized, any one of a number of suitable transcription and translation elements may be used. In specific embodiments, the reporter gene is expressed, or a fusion protein comprising the reporter gene and ORF of a fragment thereof, of the target gene is expressed.

[0148] Any of the methods previously described for the insertion of DNA fragments into a vector may be used to construct expression vectors containing a chimeric nucleic acid consisting of appropriate transcriptional/translational control signals and the protein coding sequences. These methods may include *in vitro* recombinant DNA and synthetic techniques and *in vivo* recombinants (genetic recombination). Expression of the reporter gene construct may be regulated by a second nucleic acid sequence so that the reporter gene is expressed in a host transformed with the recombinant DNA molecule. For example, expression of a reporter gene construct may be controlled by any promoter/enhancer element known in the art, such as a constitutive promoter, a tissue-specific promoter, or an inducible promoter. Specific examples of promoters which may be used to control gene expression include, but are not limited to, the SV40 early promoter region (Bernoist & Chambon, 1981, Nature 290:304-310), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., 1980, Cell 22:787-797), the herpes thymidine kinase promoter (Wagner et al., 1981, Proc. Natl. Acad. Sci. U.S.A. 78:1441-1445), the regulatory sequences of the metallothionein gene (Brinster et al., 1982, Nature 296:39-42); prokaryotic expression vectors such as the  $\beta$ -lactamase promoter (Villa-Kamaroff et al., 1978, Proc. Natl. Acad. Sci. U.S.A. 75:3727-3731), or the *tac* promoter (DeBoer et al., 1983, Proc. Natl. Acad. Sci. U.S.A. 80:21-25); see also "Useful proteins from recombinant bacteria" in Scientific American, 1980, 242:74-94; plant expression vectors comprising the nopaline synthetase promoter region (Herrera-Estrella et al., Nature 303:209-213) or the cauliflower mosaic virus 35S RNA promoter (Gardner, et al., 1981, Nucl. Acids Res. 9:2871), and the promoter of the photosynthetic enzyme ribulose biphosphate carboxylase (Herrera-Estrella et al., 1984, Nature 310:115-120); promoter elements from yeast or other fungi such as the Gal 4 promoter, the ADC (alcohol dehydrogenase) promoter, PGK (phosphoglycerol kinase) promoter, alkaline phosphatase promoter, and the following animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: elastase I gene control region which is active in pancreatic acinar cells (Swift et al., 1984, Cell 38:639-646; Ornitz et al., 1986, Cold Spring Harbor Symp. Quant. Biol. 50:399-409; MacDonald, 1987, Hepatology 7:425-515); insulin gene control region which is active in pancreatic beta cells (Hanahan, 1985, Nature 315:115-122), immunoglobulin gene control region which is active in lymphoid cells (Grosschedl et al., 1984, Cell 38:647-658; Adames et al., 1985, Nature 318:533-538; Alexander et al., 1987, Mol. Cell. Biol. 7:1436-1444), mouse mammary tumor virus control region which is active in testicular, breast, lymphoid and mast cells (Leder et al., 1986, Cell 45:485-495), albumin gene control region which is active in liver

(Pinkert et al., 1987, Genes and Devel. 1:268-276), alpha-fetoprotein gene control region which is active in liver (Krumlauf et al., 1985, Mol. Cell. Biol. 5:1639-1648; Hammer et al., 1987, Science 235:53-58; alpha 1-antitrypsin gene control region which is active in the liver (Kelsey et al., 1987, Genes and Devel. 1:161-171), beta-globin gene control region which is active in myeloid cells (Mogam et al., 1985, Nature 315:338-340; Kollias et al., 1986, Cell 46:89-94; myelin basic protein gene control region which is active in oligodendrocyte cells in the brain (Readhead et al., 1987, Cell 48:703-712); myosin light chain-2 gene control region which is active in skeletal muscle (Sani, 1985, Nature 314:283-286), and gonadotropic releasing hormone gene control region which is active in the hypothalamus (Mason et al., 1986, Science 234:1372-1378).

[0149] In a specific embodiment, a vector is used that comprises a promoter operably linked to a reporter gene flanked by one or more UTRs of a target gene, origins of replication from one or more species, and, optionally, one or more selectable markers (*e.g.*, an antibiotic resistance gene). In a preferred embodiment, the vectors are CMV vectors, T7 vectors, lac vectors, pCEP4 vectors, or 5.0/FRT vectors.

[0150] In a specific embodiment, an expression construct is made by amplifying the 5' and/or 3' UTRs of a target gene and ligating the UTRs to a reporter gene such as luciferase, and subcloning them into a pT-Adv vector (Clontech Laboratories, Palo Alto, California). It is understood by one of skill in the art that the construction of the reporter plasmid may require the construction of intermediate plasmids if several ligations are involved.

[0151] Expression vectors containing the reporter gene construct of the present invention can be identified by four general approaches: (a) nucleic acid sequencing, (b) nucleic acid hybridization, (c) presence or absence of "marker" nucleic acid functions, and (d) expression of inserted sequences. In the first approach, the presence of the UTRs and/or the reporter gene inserted in an expression vector can be detected by sequencing. In the second approach, the presence of the UTRs and/or the reporter gene inserted in an expression vector can be detected by nucleic acid hybridization using probes comprising sequences that are homologous to the inserted UTRs and/or reporter gene. In the third approach, the recombinant vector/host system can be identified and selected based upon the presence or absence of certain "marker" nucleic acid functions (*e.g.*, thymidine kinase activity, resistance to antibiotics, transformation phenotype, occlusion body formation in baculovirus, *etc.*) caused by the insertion of the nucleic acid of interest, *i.e.*, the reporter gene construct, in the vector. For example, if the nucleic acid of interest is inserted within the marker nucleic acid sequence of the vector, recombinants containing the insert can be identified by the absence of the marker nucleic acid function. In the fourth approach,

recombinant expression vectors can be identified by assaying the reporter gene product expressed by the recombinant. Such assays can be based, for example, on the physical or functional properties of the particular reporter gene.

[0152] In a preferred embodiment, the reporter gene constructs are cloned into stable cell line expression vectors. In a preferred embodiment, the stable cell line expression vector contains a site specific genomic integration site, such as but not limited to, pCMP1 (see, *e.g.*, FIG. 8C in Example 10). In another preferred embodiment, the reporter gene construct is cloned into an episomal mammalian expression vector, such as, but not limited to, pCMR2 (see, *e.g.*, FIG. 8B in Example 10).

#### 5.2.3.2. Transfection

[0153] Once a vector encoding the appropriate gene has been synthesized, a host cell is transformed or transfected with the vector of interest. The use of stable transformants is preferred. In a preferred embodiment, the host cell is a mammalian cell. In a more preferred embodiment, the host cell is a human cell. In another embodiment, the host cells are primary cells isolated from a tissue or other biological sample of interest. Host cells that can be used in the methods of the present invention include, but are not limited to, hybridomas, pre-B cells, 293 cells, 293T cells, HeLa cells, HepG2 cells, K562 cells, 3T3 cells, MCF7 cells, SkBr3 cells, or BT474 cells. In another preferred embodiment, the host cells are derived from tissue specific to the target gene. In yet another preferred embodiment, the host cells are immortalized cell lines derived from a source, *e.g.*, a tissue, specific to the target gene. Other host cells that can be used in the present invention include, but are not limited to, bacterial cells, yeast cells, virally-infected cells, or plant cells.

[0154] Transformation may be by any known method for introducing polynucleotides into a host cell, for example by packaging the polynucleotide in a virus and transducing a host cell with the virus, and by direct uptake of the polynucleotide. The transformation procedure used depends upon the host to be transformed. Bacterial transformation by direct uptake generally employs treatment with calcium or rubidium chloride (see, *e.g.*, Cohen, 1972, Proc. Nat. Acad. Sci. USA 69:2110 and Maniatis et al., 1982, "Molecular Cloning; A Laboratory Manual" (Cold Spring Harbor Press, Cold Spring Harbor, N.Y.). Yeast transformation by direct uptake may be carried out using the method of Schiestl & Gietz, 1989, Current Genetics 16:339-346 or Hinnen et al., 1978, Proc. Nat. Acad. Sci. USA 75:1929. Mammalian transformations (*i.e.*, transfections) by direct uptake may be conducted using the calcium phosphate precipitation method of Graham & Van der Eb, 1978, Virol. 52:546, or the various known modifications thereof. Other methods for

introducing recombinant polynucleotides into cells, particularly into mammalian cells, include dextran-mediated transfection, calcium phosphate mediated transfection, polybrene mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the polynucleotides into nuclei. Such methods are well-known to one of skill in the art.

**[0155]** In a preferred embodiment, stable cell lines containing the constructs of interest are generated for high throughput screening. Such stable cells lines may be generated by introducing a reporter gene construct comprising a selectable marker, allowing the cells to grow for 1-2 days in an enriched medium, and then growing the cells on a selective medium. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines.

**[0156]** A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (see, *e.g.*, Wigler et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (see, *e.g.*, Szybalska & Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (see, *e.g.*, Lowy et al., 1980, Cell 22:817) genes can be employed in tk-, hgp<sup>rt</sup>- or ap<sup>rt</sup>- cells, respectively. Also, anti-metabolite resistance can be used as the basis of selection for dhfr, which confers resistance to methotrexate (see, *e.g.*, Wigler et al., 1980, Natl. Acad. Sci. USA 77:3567 and O'Hare et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to mycophenolic acid (see, *e.g.*, Mulligan & Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 gene (see, *e.g.*, Colberre-Garapin et al., 1981, J. Mol. Biol. 150:1); and hyg<sup>ro</sup>, which confers resistance to hygromycin genes (see, *e.g.*, Santerre et al., 1984, Gene 30:147).

#### **5.2.4. Cell-Free Extracts**

**[0157]** The invention provides for the translation of the reporter gene constructs in a cell-free system. Techniques for practicing this specific aspect of this invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, and recombinant DNA manipulation and production, which are routinely practiced by one of skill in the art. See, *e.g.*, Sambrook, 1989, Molecular Cloning, A Laboratory Manual, Second Edition; DNA Cloning, Volumes I and II (Glover, Ed. 1985); and Transcription and Translation (Hames & Higgins, Eds. 1984).

**[0158]** Any technique well-known to one of skill in the art may be used to generate cell-free extracts for translation *in vitro* (otherwise referred to herein as cell-free translation mixtures). For example, the cell-free extracts for *in vitro* translation reactions can be

generated by centrifuging cells and clarifying the supernatant. The cell extracts for the present invention is about a S1 (*i.e.*, the supernatant from a 1,000 x g spin) to about a S500 extract (*i.e.*, the supernatant from a 500,000 x g spin), preferably about a S10 (*i.e.*, the supernatant from a 10,000 x g spin) to S250 (*i.e.*, the supernatant from a 250,000 x g spin) extract. In some embodiments, about a S50 (*i.e.*, the supernatant from a 50,000 x g spin) to S100 (*i.e.*, the supernatant from a 100,000 x g spin) extract is preferred.

[0159] The cell-free translation extract may be isolated from cells of any species origin. For example, the cell-free translation extract may be isolated from human cells (*e.g.*, HeLa cells), 293 cells, Vero cells, yeast, mouse cells (*e.g.*, cultured mouse cells), rat cells (*e.g.*, cultured rat cells), Chinese hamster ovary (CHO) cells, *Xenopus* oocytes, rabbit reticulocytes, primary cells, cancer cells (*e.g.*, undifferentiated cancer cells), cell lines, wheat germ, rye embryo, or bacterial cell extract (see, *e.g.*, Krieg & Melton, 1984, *Nature* 308:203 and Dignam et al., 1990 *Methods Enzymol.* 182:194-203). Alternatively, the cell-free translation extract, *e.g.*, rabbit reticulocyte lysates and wheat germ extract, can be purchased from, *e.g.*, Promega, (Madison, WI). It is preferred that the cells from which the cell-free extract is obtained do not endogenously express a target gene of interest. In a preferred embodiment, the cell-free extract is an extract isolated from human cells. In a more preferred embodiment, the human cells are HeLa cells.

### 5.3. Libraries of Compounds

[0160] Libraries screened using the methods of the present invention can comprise a variety of types of compounds. Examples of libraries that can be screened in accordance with the methods of the invention include, but are not limited to, peptoids; random biooligomers; diversomers such as hydantoins, benzodiazepines and dipeptides; vinylogous polypeptides; nonpeptidal peptidomimetics; oligocarbamates; peptidyl phosphonates; peptide nucleic acid libraries; antibody libraries; carbohydrate libraries; and small molecule libraries (preferably small organic molecules). In some embodiments, the compounds in the libraries screened are nucleic acid or peptide molecules. In a non-limiting example, peptide molecules can exist in a phage display library. In other embodiments, the types of compounds include, but are not limited to, peptide analogs including peptides comprising non-naturally occurring amino acids, *e.g.*, D-amino acids, phosphorous analogs of amino acids, such as  $\alpha$ -amino phosphoric acids and  $\alpha$ -amino phosphonic acids, or amino acids having non-peptide linkages, nucleic acid analogs such as phosphorothioates and PNAs, hormones, antigens, synthetic or naturally occurring drugs, opiates, dopamine, serotonin, catecholamines, thrombin, acetylcholine, prostaglandins, organic molecules, pheromones,

adenosine, sucrose, glucose, lactose and galactose. Libraries of polypeptides or proteins can also be used in the assays of the invention.

[0161] In a preferred embodiment, the combinatorial libraries are small organic molecule libraries including, but not limited to, benzodiazepines, isoprenoids, beta carbalines, thiazolidinones, metathiazanones, pyrrolidines, morpholino compounds, and benzodiazepines. In another embodiment, the combinatorial libraries comprise peptoids; random bio-oligomers; benzodiazepines; diversomers such as hydantoins, benzodiazepines and dipeptides, vinylogous polypeptides; nonpeptidal peptidomimetics; oligocarbamates; peptidyl phosphonates; peptide nucleic acid libraries; antibody libraries; or carbohydrate libraries. Combinatorial libraries are themselves commercially available (see, *e.g.*, ComGenex, Princeton, New Jersey; Asinex, Moscow, Ru, Tripos, Inc., St. Louis, Missouri; ChemStar, Ltd, Moscow, Russia; 3D Pharmaceuticals, Exton, Pennsylvania; Martek Biosciences, Columbia, Maryland; *etc.*).

[0162] In a preferred embodiment, the library is preselected so that the compounds of the library are more amenable for cellular uptake. For example, compounds are selected based on specific parameters such as, but not limited to, size, lipophilicity, hydrophilicity, and hydrogen bonding, which enhance the likelihood of compounds getting into the cells. In another embodiment, the compounds are analyzed by three-dimensional or four-dimensional computer computation programs.

[0163] The combinatorial compound library for use in accordance with the methods of the present invention may be synthesized. There is a great interest in synthetic methods directed toward the creation of large collections of small organic compounds, or libraries, which could be screened for pharmacological, biological or other activity. The synthetic methods applied to create vast combinatorial libraries are performed in solution or in the solid phase, *i.e.*, on a solid support. Solid-phase synthesis makes it easier to conduct multi-step reactions and to drive reactions to completion with high yields because excess reagents can be easily added and washed away after each reaction step. Solid-phase combinatorial synthesis also tends to improve isolation, purification and screening. However, the more traditional solution phase chemistry supports a wider variety of organic reactions than solid-phase chemistry.

[0164] Combinatorial compound libraries of the present invention may be synthesized using the apparatus described in U.S. Patent No. 6,190,619 to Kilcoin et al., which is hereby incorporated by reference in its entirety. U.S. Patent No. 6,190,619 discloses a synthesis apparatus capable of holding a plurality of reaction vessels for parallel synthesis of multiple discrete compounds or for combinatorial libraries of compounds.



**[0165]** In one embodiment, the combinatorial compound library can be synthesized in solution. The method disclosed in U.S. Patent No. 6,194,612 to Boger et al., which is hereby incorporated by reference in its entirety, features compounds useful as templates for solution phase synthesis of combinatorial libraries. The template is designed to permit reaction products to be easily purified from unreacted reactants using liquid/liquid or solid/liquid extractions. The compounds produced by combinatorial synthesis using the template will preferably be small organic molecules. Some compounds in the library may mimic the effects of non-peptides or peptides. In contrast to solid phase synthesis of combinatorial compound libraries, liquid phase synthesis does not require the use of specialized protocols for monitoring the individual steps of a multistep solid phase synthesis (Egner et al., 1995, J. Org. Chem. 60:2652; Anderson et al., 1995, J. Org. Chem. 60:2650; Fitch et al., 1994, J. Org. Chem. 59:7955; Look et al., 1994, J. Org. Chem. 49:7588; Metzger et al., 1993, Angew. Chem., Int. Ed. Engl. 32:894; Youngquist et al., 1994, Rapid Commun. Mass Spect. 8:77; Chu et al., 1995, J. Am. Chem. Soc. 117:5419; Brummel et al., 1994, Science 264:399; and Stevanovic et al., 1993, Bioorg. Med. Chem. Lett. 3:431).

**[0166]** Combinatorial compound libraries useful for the methods of the present invention can be synthesized on solid supports. In one embodiment, a split synthesis method, a protocol of separating and mixing solid supports during the synthesis, is used to synthesize a library of compounds on solid supports (see *e.g.*, Lam et al., 1997, Chem. Rev. 97:41-448; Ohlmeyer et al., 1993, Proc. Natl. Acad. Sci. USA 90:10922-10926 and references cited therein). Each solid support in the final library has substantially one type of compound attached to its surface. Other methods for synthesizing combinatorial libraries on solid supports, wherein one product is attached to each support, will be known to those of skill in the art (see, *e.g.*, Nefzi et al., 1997, Chem. Rev. 97:449-472).

**[0167]** As used herein, the term "solid support" is not limited to a specific type of solid support. Rather a large number of supports are available and are known to one skilled in the art. Solid supports include silica gels, resins, derivatized plastic films, glass beads, cotton, plastic beads, polystyrene beads, alumina gels, and polysaccharides. A suitable solid support may be selected on the basis of desired end use and suitability for various synthetic protocols. For example, for peptide synthesis, a solid support can be a resin such as p-methylbenzhydrylamine (pMBHA) resin (Peptides International, Louisville, KY), polystyrenes (*e.g.*, PAM-resin obtained from Bachem Inc., Peninsula Laboratories, *etc.*), including chloromethylpolystyrene, hydroxymethylpolystyrene and aminomethylpolystyrene, poly (dimethylacrylamide)-grafted styrene co-divinyl-benzene (*e.g.*, POLYHIPE resin, obtained from Aminotech, Canada), polyamide resin (obtained

from Peninsula Laboratories), polystyrene resin grafted with polyethylene glycol (*e.g.*, TENTAGEL or ARGOGEL, Bayer, Tübingen, Germany) polydimethylacrylamide resin (obtained from Milligen/Biosearch, California), or Sepharose (Pharmacia, Sweden).

[0168] In some embodiments of the present invention, compounds can be attached to solid supports via linkers. Linkers can be integral and part of the solid support, or they may be nonintegral that are either synthesized on the solid support or attached thereto after synthesis. Linkers are useful not only for providing points of compound attachment to the solid support, but also for allowing different groups of molecules to be cleaved from the solid support under different conditions, depending on the nature of the linker. For example, linkers can be, *inter alia*, electrophilically cleaved, nucleophilically cleaved, photocleavable, enzymatically cleaved, cleaved by metals, cleaved under reductive conditions or cleaved under oxidative conditions. In a preferred embodiment, the compounds are cleaved from the solid support prior to high throughput screening of the compounds.

#### 5.4. Reporter Gene-Based Screening Assays

##### 5.4.1. Cell-Based Assays

[0169] After a vector containing the reporter gene construct is transformed or transfected into a host cell and a compound library is synthesized or purchased or both, the cells are used to screen the library to identify compounds that modulate untranslated region-dependent expression of a target gene. In a preferred embodiment, the cells are stably transfected with the reporter gene construct. The reporter gene-based assays may be conducted by contacting a compound or a member of a library of compounds with a cell genetically engineered to express a nucleic acid comprising a reporter gene operably linked to one or more untranslated regions of a target gene, and measuring the expression of said reporter gene. The alteration in reporter gene expression relative to a previously determined reference range, the absence of the compound or a control in such reporter-gene based assays indicates that a particular compound modulates untranslated region-dependent expression of a target gene. In a preferred embodiment, a negative control (*e.g.*, PBS or another agent that is known to have no effect on the expression of the reporter gene) and a positive control (*e.g.*, an agent that is known to have an effect on the expression of the reporter gene, preferably an agent that effects untranslated region-dependent expression) are included in the cell-based assays described herein.

[0170] The step of contacting a compound or a member of a library of compounds with a cell genetically engineered to express a reporter gene operably linked to one or more

untranslated regions may be conducted under physiologic conditions. In specific embodiment, a compound or a member of a library of compounds is added to the cells in the presence of an aqueous solution. In accordance with this embodiment, the aqueous solution may comprise a buffer and a combination of salts, preferably approximating or mimicking physiologic conditions. Alternatively, the aqueous solution may comprise a buffer, a combination of salts, and a detergent or a surfactant. Examples of salts which may be used in the aqueous solution include, but not limited to, KCl, NaCl, and/or MgCl<sub>2</sub>. The optimal concentration of each salt used in the aqueous solution is dependent on the cells and compounds used and can be determined using routine experimentation.

[0171] The invention provides for contacting a compound or a member of a library of compounds with a cell genetically engineering to express a reporter gene operably linked to one or more untranslated regions for a specific period of time. For example, the contacting can take place for about 1 minute, 2 minutes, 3 minutes, 4, minutes, 5, minutes, 10 minutes, 15 minutes, 20 minutes, 30 minutes, 1 hour, 2 hours, 3 hours, 4 hours, 5 hours, 10 hours, 15 hours, 20 hours, 1 day, 2 days, 3 days, 4 days, 5 days, or 1 week. In a preferred embodiment, the contacting is about 15 hours, *i.e.*, overnight. The contacting can take place for about 1 minute to 1 week, preferably about 5 minutes to 5 days, more preferably about 10 minutes to 2 days, and even more preferably about 1 hour to 1 day.

[0172] In one embodiment, the invention provides a method for identifying a compound that modulates untranslated region-dependent expression of a target gene, said method comprising: (a) expressing a nucleic acid comprising a reporter gene operably linked to one or more untranslated regions of said target gene in a cell; (b) contacting said cell with a member of a library of compounds; and (c) detecting the expression of said reporter gene, wherein a compound that modulates untranslated region-dependent regulation of expression is identified if the expression of said reporter gene in the presence of a compound is altered relative to a previously determined reference range or the expression of said reporter gene in the absence of said compound or the presence of a control (*e.g.*, phosphate buffered saline ("PBS")). In another embodiment, the invention provides a method for identifying a compound that modulates untranslated region-dependent expression of a target gene, said method comprising: (a) contacting a member of a library of compounds with a cell containing a nucleic acid comprising a reporter gene operably linked to one or more untranslated regions of said target gene; and (b) detecting a reporter protein translated from said reporter gene, wherein detecting the expression of said reporter gene, wherein a compound that modulates untranslated region-dependent expression is identified if the expression of said reporter gene in the presence of a compound is altered relative to a

previously determined reference range or the expression of said reporter gene in the absence of said compound or the presence of a control (*e.g.*, PBS).

[0173] The invention also provides methods of identifying compounds that upregulate or down-regulate untranslated region-dependent expression of a target gene utilizing the cell-based reporter gene assays described herein. In a specific embodiment, the invention provides a method of upregulating untranslated region-dependent expression of a target gene, said method comprising (a) contacting a compound with a cell containing a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of said target gene; and (b) detecting a reporter gene protein translated from said reporter gene, wherein a compound that upregulates untranslated region dependent expression is identified if the expression of said reporter gene in the presence of a compound is increased relative a previously determined reference range, or the expression in the absence of the compound or the presence of a control (*e.g.*, PBS). In another embodiment, the invention provides a method of down-regulating untranslated region-dependent expression of a target gene, said method comprising (a) contacting a compound with a cell containing a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of said target gene; and (b) detecting a reporter gene protein translated from said reporter gene, wherein a compound that down-regulates untranslated region dependent expression is identified if the expression of said reporter gene in the presence of a compound is decreased relative a previously determined reference range, or the expression in the absence of the compound or the presence of a control (*e.g.*, PBS).

[0174] The present invention provides methods of identifying environmental stimuli (*e.g.*, exposure to different concentrations of CO<sub>2</sub> and/or O<sub>2</sub>, stress and different pHs) that modulate untranslated region-dependent expression of a target gene utilizing the cell-based reporter gene assays described herein. In particular, the invention provides a method of identifying an environmental stimulus, said method comprising (a) contacting a cell containing a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of said target gene with an environmental stimulus; and (b) detecting a reporter gene protein translated from said reporter gene, wherein a compound that modulates untranslated region dependent expression is identified if the expression of said reporter gene in the presence of an environmental stimuli is altered relative to a previously determined reference range, or the expression in the absence of the compound or the presence of a control (*e.g.*, PBS). In a specific embodiment, the environmental stimuli

is not hypoxia. In another embodiment, the environmental stimuli does not include a compound.

[0175] The expression of a reporter gene in the cell-based reporter-gene assays may be detected by any technique well-known to one of skill in the art. Methods for detecting the expression of a reporter gene will vary with the reporter gene used. Assays for the various reporter genes are well-known to one of skill in the art. For example, as described in Section 5.2.1., luciferase, beta-galactosidase (“b-gal”), beta-glucoronidase (“GUS”), beta-lactamase, chloramphenicol acetyltransferase (“CAT”), and alkaline phosphatase (“AP”) are enzymes that can be analyzed in the presence of a substrate and could be amenable to high throughput screening. For example, the reaction products of luciferase, beta-galactosidase (“b-gal”), and alkaline phosphatase (“AP”) are assayed by changes in light imaging (*e.g.*, luciferase), spectrophotometric absorbance (*e.g.*, b-gal), or fluorescence (*e.g.*, AP). Assays for changes in light output, absorbance, and/or fluorescence are easily adapted for high throughput screening. For example, b-gal activity can be measured with a microplate reader. Green fluorescent protein (“GFP”) activity can be measured by changes in fluorescence. For example, in the case of mutant GFPs that fluoresce at 488 nm, standard fluorescence activated cell sorting (“FACS”) equipment can be used to separate cells based upon GFP activity.

[0176] Alterations in the expression of a reporter gene may be determined by comparing the level of expression of the reporter gene to a negative control (*e.g.*, PBS or another agent that is known to have no effect on the expression of the reporter gene) and optionally, a positive control (*e.g.*, an agent that is known to have an effect on the expression of the reporter gene, preferably an agent that effects untranslated region-dependent expression). Alternatively, alterations in the expression of a reporter gene may be determined by comparing the level of expression of the reporter gene to a previously determined reference range.

#### **5.4.2. Cell-Free Assays**

[0177] After a vector containing the reporter gene construct is produced, a cell-free translation extract is generated or purchased, and a compound library is synthesized or purchased or both, the cell-free translation extract and nucleic acid are used to screen the library to identify compounds that modulate untranslated region-dependent expression of a target gene. The reporter gene-based assays may be conducted in a cell-free manner by contacting a compound or a member of a library of compounds with a cell-free translation mixture and a nucleic acid comprising a reporter gene operably linked to one or more untranslated regions of a target gene, and measuring the expression of said reporter gene.

The alteration in reporter gene expression relative to a previously determined reference range, the absence of a compound or a control in such reporter-gene based assays indicates that a particular compound modulates untranslated region-dependent expression of a target gene. In a preferred embodiment, a negative control (*e.g.*, PBS or another agent that is known to have no effect on the expression of the reporter gene) and a positive control (*e.g.*, an agent that is known to have an effect on the expression of the reporter gene, preferably an agent that effects untranslated region-dependent expression) are included in the cell-free assays described herein.

[0178] The step of contacting a compound or a member of a library of compounds with a cell-free translation mixture containing a nucleic acid comprising a reporter gene operably linked to one or more untranslated regions may be conducted under conditions approximating or mimicking physiologic conditions. In specific embodiment, a compound or a member of a library of compounds is added to the cells in the presence of an aqueous solution. In accordance with this embodiment, the aqueous solution may comprise a buffer and a combination of salts, preferably approximating or mimicking physiologic conditions. Alternatively, the aqueous solution may comprise a buffer, a combination of salts, and a detergent or a surfactant. Examples of salts which may be used in the aqueous solution include, but not limited to, KCl, NaCl, and/or MgCl<sub>2</sub>. The optimal concentration of each salt used in the aqueous solution is dependent on the cells and compounds used and can be determined using routine experimentation.

[0179] The invention provides for contacting a compound or a member of a library of compounds with a cell genetically engineering to express a reporter gene operably linked to one or more untranslated regions for a specific period of time. For example, the contacting can take place for about 1 minute, 2 minutes, 3 minutes, 4, minutes, 5, minutes, 10 minutes, 15 minutes, 20 minutes, 30 minutes, 1 hour, 2 hours, 3 hours, 4 hours, 5 hours, 10 hours, 15 hours, 20 hours, 1 day, 2 days, 3 days, 4 days, 5 days, or 1 week. In a preferred embodiment, the contacting is about 15 hours, *i.e.*, overnight. The contacting can take place for about 1 minute to 1 week, preferably about 5 minutes to 5 days, more preferably about 10 minutes to 2 days, and even more preferably about 1 hour to 1 day.

[0180] In a specific embodiment, the invention provides a method for identifying a compound that modulates untranslated region-dependent expression of a target gene, said method comprising: (a) contacting a member of a library of compounds with a cell-free translation mixture and a nucleic acid comprising a reporter gene operably linked to one or more untranslated regions of said target gene; and (b) detecting the expression of said reporter gene, wherein a compound that modulates untranslated region-dependent

expression is identified if the expression of said reporter gene in the presence of a compound is altered relative to a previously determined reference range or the expression of said reporter gene in the absence of said compound or the presence of a control (*e.g.*, PBS).

[0181] The invention also provides methods of identifying compounds that upregulate or down-regulate untranslated region-dependent expression of a target gene utilizing the cell-free reporter gene assays described herein. In a specific embodiment, the invention provides a method of upregulating untranslated region-dependent expression of a target gene, said method comprising (a) contacting a compound with a cell-free translation mixture and a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of said target gene; and (b) detecting a reporter gene protein translated from said reporter gene, wherein a compound that upregulates untranslated region dependent expression is identified if the expression of said reporter gene in the presence of a compound is increased relative a previously determined reference range, or the expression in the absence of the compound or the presence of a control (*e.g.*, PBS). In another embodiment, the invention provides a method of down-regulating untranslated region-dependent expression of a target gene, said method comprising (a) contacting a compound with a cell-free translation mixture and a nucleic acid comprising a reporter gene operably linked to one, two, three or more untranslated regions of said target gene; and (b) detecting a reporter gene protein translated from said reporter gene, wherein a compound that down-regulates untranslated region dependent expression is identified if the expression of said reporter gene in the presence of a compound is decreased relative to a previously determined reference range, or the expression in the absence of the compound or the presence of a control (*e.g.*, PBS).

[0182] The activity of a compound in the *in vitro* translation mixture can be determined by assaying the activity of a reporter protein encoded by a reporter gene, or alternatively, by quantifying the expression of the reporter gene by, for example, labeling the *in vitro* translated protein (*e.g.*, with <sup>35</sup>S-labeled methionine), northern blot analysis, RT-PCR or by immunological methods, such as western blot analysis or immunoprecipitation. Such methods are well-known to one of skill in the art.

#### **5.4.3. Direct Binding Assays**

[0183] Compounds that modulate untranslated region-dependent expression of a target gene can be identified by direct binding assays. In this embodiment, the target RNA comprises one or more untranslated regions, and preferably contains at least one element of an untranslated region. Such assays are described in International Patent Publication Nos. WO 02/083837 and WO 02/083953, the disclosures of which are hereby incorporated by

reference in their entireties. Briefly, direct binding assays may be conducted by attaching a library of compounds to solid supports, *e.g.*, polymer beads, with each solid support having substantially one type of compound attached to its surface. The plurality of solid supports of the library is exposed in aqueous solution to target RNA having a detectable label, forming a dye-labeled target RNA:support-attached compound complex. Binding of a target RNA molecule to a particular compound labels the solid support, *e.g.*, bead, comprising the compound, which can be physically separated from other, unlabeled solid supports. Once labeled solid supports are identified, the chemical structures of the compounds thereon can be determined by, *e.g.*, by reading a code on the solid support that correlates with the structure of the attached compound.

[0184] Alternatively, direct binding assays may be conducted by contacting a target RNA having a detectable label with a member of a library of compounds free in solution, in labeled tubes or microtiter wells, or a microarray. Compounds in the library that bind to the labeled target RNA will form a detectably labeled complex that can be identified and removed from the uncomplexed, unlabeled compounds in the library, and from uncomplexed, labeled target RNA, by a variety of methods including, but not limited to, methods that differentiate changes in the electrophoretic, chromatographic, or thermostable properties of the complexed target RNA.

#### 5.4.3.1. Electrophoresis

[0185] Methods for separation of the complex of a target RNA bound to a compound from the unbound RNA comprises any method of electrophoretic separation, including but not limited to, denaturing and non-denaturing polyacrylamide gel electrophoresis, urea gel electrophoresis, gel filtration, pulsed field gel electrophoresis, two dimensional gel electrophoresis, continuous flow electrophoresis, zone electrophoresis, agarose gel electrophoresis, and capillary electrophoresis.

[0186] In a preferred embodiment, an automated electrophoretic system comprising a capillary cartridge having a plurality of capillary tubes is used for high-throughput screening of compounds bound to target RNA. Such an apparatus for performing automated capillary gel electrophoresis is disclosed in U.S. Patent Nos. 5,885,430; 5,916,428; 6,027,627; and 6,063,251, the disclosures of which are incorporated by reference in their entireties.

[0187] The device disclosed in U.S. Patent No. 5,885,430, which is incorporated by reference in its entirety, allows one to simultaneously introduce samples into a plurality of capillary tubes directly from microtiter trays having a standard size. U.S. Patent No. 5,885,430 discloses a disposable capillary cartridge which can be cleaned between



electrophoresis runs, the cartridge having a plurality of capillary tubes. A first end of each capillary tube is retained in a mounting plate, the first ends collectively forming an array in the mounting plate. The spacing between the first ends corresponds to the spacing between the centers of the wells of a microtiter tray having a standard size. Thus, the first ends of the capillary tubes can simultaneously be dipped into the samples present in the tray's wells. The cartridge is provided with a second mounting plate in which the second ends of the capillary tubes are retained. The second ends of the capillary tubes are arranged in an array which corresponds to the wells in the microtiter tray, which allows for each capillary tube to be isolated from its neighbors and therefore free from cross-contamination, as each end is dipped into an individual well.

**[0188]** Plate holes may be provided in each mounting plate and the capillary tubes inserted through these plate holes. In such a case, the plate holes are sealed airtight so that the side of the mounting plate having the exposed capillary ends can be pressurized. Application of a positive pressure in the vicinity of the capillary openings in this mounting plate allows for the introduction of air and fluids during electrophoretic operations and also can be used to force out gel and other materials from the capillary tubes during reconditioning. The capillary tubes may be protected from damage using a needle comprising a cannula and/or plastic tubes, and the like when they are placed in these plate holes. When metallic cannula or the like are used, they can serve as electrical contacts for current flow during electrophoresis. In the presence of a second mounting plate, the second mounting plate is provided with plate holes through which the second ends of the capillary tubes project. In this instance, the second mounting plate serves as a pressure containment member of a pressure cell and the second ends of the capillary tubes communicate with an internal cavity of the pressure cell. The pressure cell is also formed with an inlet and an outlet. Gels, buffer solutions, cleaning agents, and the like may be introduced into the internal cavity through the inlet, and each of these can simultaneously enter the second ends of the capillaries.

**[0189]** In another preferred embodiment, the automated electrophoretic system can comprise a chip system consisting of complex designs of interconnected channels that perform and analyze enzyme reactions using part of a channel design as a tiny, continuously operating electrophoresis material, where reactions with one sample are going on in one area of the chip while electrophoretic separation of the products of another sample is taking place in a different part of the chip. Such a system is disclosed in U.S. Patent Nos. 5,699,157; 5,842,787; 5,869,004; 5,876,675; 5,942,443; 5,948,227; 6,042,709; 6,042,710; 6,046,056; 6,048,498; 6,086,740; 6,132,685; 6,150,119; 6,150,180; 6,153,073; 6,167,910;

6,171,850; and 6,186,660, the disclosures of which are incorporated by reference in their entireties.

[0190] The system disclosed in U.S. Patent No. 5,699,157, which is hereby incorporated by reference in its entirety, provides for a microfluidic system for high-speed electrophoretic analysis of subject materials for applications in the fields of chemistry, biochemistry, biotechnology, molecular biology and numerous other areas. The system has a channel in a substrate, a light source and a photoreceptor. The channel holds subject materials in solution in an electric field so that the materials move through the channel and separate into bands according to species. The light source excites fluorescent light in the species bands and the photoreceptor is arranged to receive the fluorescent light from the bands. The system further has a means for masking the channel so that the photoreceptor can receive the fluorescent light only at periodically spaced regions along the channel. The system also has an unit connected to analyze the modulation frequencies of light intensity received by the photoreceptor so that velocities of the bands along the channel are determined, which allows the materials to be analyzed.

[0191] The system disclosed in U.S. Patent No. 5,699,157 also provides for a method of performing high-speed electrophoretic analysis of subject materials, which comprises the steps of holding the subject materials in solution in a channel of a microfluidic system; subjecting the materials to an electric field so that the subject materials move through the channel and separate into species bands; directing light toward the channel; receiving light from periodically spaced regions along the channel simultaneously; and analyzing the frequencies of light intensity of the received light so that velocities of the bands along the channel can be determined for analysis of said materials. The determination of the velocity of a species band determines the electrophoretic mobility of the species and its identification.

[0192] U.S. Patent No. 5,842,787, which is hereby incorporated by reference in its entirety, is generally directed to devices and systems employ channels having, at least in part, depths that are varied over those which have been previously described (such as the device disclosed in U.S. Patent No. 5,699,157), wherein said channel depths provide numerous beneficial and unexpected results such as but not limited to, a reduction in sample perturbation, reduced non-specific sample mixture by diffusion, and increased resolution.

[0193] In another embodiment, the electrophoretic method of separation comprises polyacrylamide gel electrophoresis. In a preferred embodiment, the polyacrylamide gel electrophoresis is non-denaturing, so as to differentiate the mobilities of the target RNA bound to a compound from free target RNA. If the polyacrylamide gel electrophoresis is

denaturing, then the target RNA:compound complex must be cross-linked prior to electrophoresis to prevent the disassociation of the target RNA from the compound during electrophoresis. Such techniques are well known to one of skill in the art.

[0194] In one embodiment of the method, the binding of compounds to target nucleic acid can be detected, preferably in an automated fashion, by gel electrophoretic analysis of interference footprinting. RNA can be degraded at specific base sites by enzymatic methods such as ribonucleases A, U<sub>2</sub>, CL<sub>3</sub>, T<sub>1</sub>, Phy M, and *B. cereus* or chemical methods such as diethylpyrocarbonate, sodium hydroxide, hydrazine, piperidine formate, dimethyl sulfate, [2,12-dimethyl-3,7,11,17-tetraazacyclo[11.3.1]heptadeca-1(17),2,11,13,15-pentaenato] nickel(II) (NiCR), cobalt(II)chloride, or iron(II) ethylenediaminetetraacetate (Fe-EDTA) as described for example in Zheng et al., 1999, *Biochem.* 37:2207-2214; Latham & Cech, 1989, *Science* 245:276-282; and Sambrook et al., 2001, in *Molecular Cloning: A Laboratory Manual*, pp 12.61-12.73, Cold Spring Harbor Laboratory Press, and the references cited therein, which are hereby incorporated by reference in their entireties. The specific pattern of cleavage sites is determined by the accessibility of particular bases to the reagent employed to initiate cleavage and, as such, is therefore determined by the three-dimensional structure of the RNA.

[0195] The interaction of small molecules with a target nucleic acid can change the accessibility of bases to these cleavage reagents both by causing conformational changes in the target nucleic acid or by covering a base at the binding interface. When a compound binds to the nucleic acid and changes the accessibility of bases to cleavage reagents, the observed cleavage pattern will change. This method can be used to identify and characterize the binding of small molecules to RNA as described, for example, by Prudent et al., 1995, *J. Am. Chem. Soc.* 117:10145-10146 and Mei et al., 1998, *Biochem.* 37:14204-14212.

[0196] In the preferred embodiment of this technique, the detectably labeled target nucleic acid is incubated with an individual compound and then subjected to treatment with a cleavage reagent, either enzymatic or chemical. The reaction mixture can be preferably be examined directly, or treated further to isolate and concentrate the nucleic acid. The fragments produced are separated by electrophoresis and the pattern of cleavage can be compared to a cleavage reaction performed in the absence of compound. A change in the cleavage pattern directly indicates that the compound binds to the target nucleic acid. Multiple compounds can be examined both in parallel and serially.

[0197] Other embodiments of electrophoretic separation include, but are not limited to urea gel electrophoresis, gel filtration, pulsed field gel electrophoresis, two dimensional gel

electrophoresis, continuous flow electrophoresis, zone electrophoresis, and agarose gel electrophoresis.

#### **5.4.3.2. Size Exclusion Chromatography**

[0198] In another embodiment of the present invention, size-exclusion chromatography is used to purify compounds that are bound to a target nucleic acid from a complex mixture of compounds. Size-exclusion chromatography separates molecules based on their size and uses gel-based media comprised of beads with specific size distributions. When applied to a column, this media settles into a tightly packed matrix and forms a complex array of pores. Separation is accomplished by the inclusion or exclusion of molecules by these pores based on molecular size. Small molecules are included into the pores and, consequently, their migration through the matrix is retarded due to the added distance they must travel before elution. Large molecules are excluded from the pores and migrate with the void volume when applied to the matrix. In the present invention, a target nucleic acid is incubated with a mixture of compounds while free in solution and allowed to reach equilibrium. When applied to a size exclusion column, compounds free in solution are retained by the column, and compounds bound to the target nucleic acid are passed through the column. In a preferred embodiment, spin columns commonly used for gel filtration of nucleic acids will be employed to separate bound from unbound compounds (*e.g.*, Bio-Spin columns manufactured by BIO-RAD). In another embodiment, the size exclusion matrix is packed into multiwell plates to allow high throughput separation of mixtures (*e.g.*, PLASMID 96-well SEC plates manufactured by Millipore).

#### **5.4.3.3. Affinity Chromatography**

[0199] In one embodiment of the present invention, affinity capture is used to purify compounds that are bound to a target nucleic acid labeled with an affinity tag from a complex mixture of compounds. To accomplish this, a target nucleic acid labeled with an affinity tag is incubated with a mixture of compounds while free in solution and then captured to a solid support once equilibrium has been established; alternatively, target nucleic acids labeled with an affinity tag can be captured to a solid support first and then allowed to reach equilibrium with a mixture of compounds.

[0200] The solid support is typically comprised of, but not limited to, cross-linked agarose beads that are coupled with a ligand for the affinity tag. Alternatively, the solid support may be a glass, silicon, metal, or carbon, plastic (polystyrene, polypropylene) surface with or without a self-assembled monolayer (SAM) either with a covalently

attached ligand for the affinity tag, or with inherent affinity for the tag on the target nucleic acid.

[0201] Once the complex between the target nucleic acid and compound has reached equilibrium and has been captured, one skilled in the art will appreciate that the retention of bound compounds and removal of unbound compounds is facilitated by washing the solid support with large excesses of binding reaction buffer. Furthermore, retention of high affinity compounds and removal of low affinity compounds can be accomplished by a number of means that increase the stringency of washing; these means include, but are not limited to, increasing the number and duration of washes, raising the salt concentration of the wash buffer, addition of detergent or surfactant to the wash buffer, and addition of non-specific competitor to the wash buffer.

[0202] In one embodiment, the compounds themselves are detectably labeled with fluorescent dyes, radioactive isotopes, or nanoparticles. When the compounds are applied to the captured target nucleic acid in a spatially addressed fashion (*e.g.*, in separate wells of a 96-well microplate), binding between the compounds and the target nucleic acid can be determined by the presence of the detectable label on the compound using fluorescence.

[0203] Following the removal of unbound compounds, bound compounds with high affinity for the target nucleic acid can be eluted from the immobilized target nucleic acids and analyzed. The elution of compounds can be accomplished by any means that break the non-covalent interactions between the target nucleic acid and compound. Means for elution include, but are not limited to, changing the pH, changing the salt concentration, the application of organic solvents, and the application of molecules that compete with the bound ligand. In a preferred embodiment, the means employed for elution will release the compound from the target RNA, but will not effect the interaction between the affinity tag and the solid support, thereby achieving selective elution of compound. Moreover, a preferred embodiment will employ an elution buffer that is volatile to allow for subsequent concentration by lyophilization of the eluted compound (*e.g.*, 0 M to 5 M ammonium acetate).

#### **5.5. Methods for Confirming that a Compound Modulates Untranslated Region-Dependent Expression**

[0204] In order to exclude the possibility that a particular compound is functioning solely by modulating the expression of a target gene in an untranslated region-independent manner, one or more mutations may be introduced into the untranslated regions operably linked to a reporter gene and the effect on the expression of the reporter gene in a reporter gene-based assay described herein can be determined. For example, a reporter gene

construct comprising the 5' UTR of a target gene may be mutated by deleting a fragment of the 5' UTR of the target gene or substituting a fragment of the 5' UTR of the target gene with a fragment of the 5' UTR of another gene and measuring the expression of the reporter gene in the presence and absence of a compound that has been identified in a screening assays described *supra* (See Section 5.4). If the deletion of a fragment of the 5' UTR of the target gene or the substitution of a fragment of the 5' UTR of the target gene with a fragment of the 5' UTR of another gene affects the ability of the compound to modulate the expression of the reporter gene, then the fragment of the 5' UTR deleted or substituted plays a role in the regulation of the reporter gene expression and the regulation, at least in part, in an untranslated region-dependent manner.

**[0205]** The possibility that a particular compound is functioning solely by modulating the expression of a target gene in an untranslated region-independent manner may be also determined by changing the vector utilized as a reporter construct. The untranslated regions flanked by a reporter gene from the first reporter construct in which an effect on reporter gene expression was detected following exposure to a compound may be inserted into a new reporter construct that has, *e.g.*, different transcriptional regulation elements (*e.g.*, a different promoter) and a different selectable marker. The level of reporter gene expression in the presence of the compound can be compared to the level of reporter gene expression in the absence of the compound or in the presence of a control (*e.g.*, PBS). If there is no change in the level of expression of the reporter gene in the presence of the compound relative to the absence of the compound or in the presence of a control, then the compound probably is functioning in an untranslated region-independent manner.

**[0206]** The specificity of a particular compound's effect on untranslated region-dependent expression of a target gene can also be determined. In particular, the effect of a particular compound on the expression of one or more genes (preferably, a plurality of genes) can be determined utilizing assays well-known to one of skill in the art or described herein. In a specific embodiment, the specificity of a particular compound for an untranslated region of a target gene is determined by (a) contacting the compound of interest with a cell containing a nucleic acid comprising a reporter gene operably linked to an UTR of a different gene (*i.e.*, a gene different from the target gene which has a UTR different from the target gene); and (b) detecting a reporter gene protein translated from the reporter gene, wherein the compound is specific for the untranslated region of the target gene if the expression of said reporter gene in the presence of the compound is not altered or is not substantially altered relative to a previously determined reference range, or the expression in the absence of the compound or the presence of a control (*e.g.*, PBS). In

another embodiment, the specificity of a particular compound for an untranslated region of a target gene is determined by (a) contacting the compound of interest with a panel of cells, each cell in a different well of a container (*e.g.*, a 48 or 96 well microtiter plate) and each cell containing a nucleic acid comprising a reporter gene operably linked to an UTR of a different gene; and (b) detecting a reporter gene protein translated from the reporter gene, wherein the compound is specific for the untranslated region of the target gene if the expression of said reporter gene in the presence of the compound is not altered or is not substantially altered relative to a previously determined reference range, or the expression in the absence of the compound or the presence of a control (*e.g.*, PBS). In accordance with this embodiment, the panel may comprise 5, 7, 10, 15, 20, 25, 50, 75, 100 or more cells. In another embodiment, the specificity of a particular compound for an untranslated region of a target gene is determined by (a) contacting the compound of interest with a cell-free translation mixture and a nucleic acid comprising a reporter gene operably linked to an UTR of a different gene; and (b) detecting a reporter gene protein translated from the reporter gene, wherein the compound is specific for the untranslated region of the target gene if the expression of said reporter gene in the presence of the compound is not altered or is not substantially altered relative to a previously determined reference range, or the expression in the absence of the compound or the presence of a control (*e.g.*, PBS). As used herein, the term “not substantially altered” means that the compound alters the expression of the reporter gene or target gene by less than 20%, less than 15%, less than 10%, less than 5%, or less than 2% relative to a negative control such as PBS.

[0207] The compounds identified in the assays described *supra* that modulate untranslated region-dependent expression of a target gene (for convenience referred to herein as a “lead” compound) can be further tested for untranslated region-dependent binding to the target RNA (which contains at least one untranslated region, and preferably at least one element of an untranslated region). Furthermore, by assessing the effect of a compound on target gene expression, *cis*-acting elements, *i.e.*, specific nucleotide sequences, that are involved in untranslated region-dependent expression may be identified.

#### **5.5.1. RNA Binding Assays**

[0208] The compounds that modulate untranslated region-dependent expression of a target gene can be tested for binding to the target RNA (which contains at least one untranslated region, and preferably at least one element of an untranslated region) by any method known in the art. See Section 5.4.3 *supra*.

### 5.5.1. Subtraction Assay

[0209] The element(s) of an untranslated region(s) that is necessary for a compound identified in accordance with the methods of the invention to modulate untranslated region-dependent expression of a target gene can be determined utilizing standard mutagenesis techniques well-known to one of skill in the art. One or more mutations (*e.g.*, deletions, additions and/or substitutions) may be introduced into the untranslated regions operably linked to a reporter gene and the effect on the expression of the reporter gene in a reporter gene-based assay described herein can be determined. For example, a reporter gene construct comprising the 5' UTR of a target gene may be mutated by deleting a fragment or all of the 5' UTR of the target gene or substituting a fragment of the 5' UTR of the target gene with a fragment of the 5' UTR of another gene and measuring the expression of the reporter gene in the presence and absence of a compound that has been identified in a screening assays described *supra* (See Section 5.4). If the deletion of a fragment of the 5' UTR of the target gene or the substitution of a fragment of the 5' UTR of the target gene with a fragment of the 5' UTR of another gene affects the ability of the compound to modulate the expression of the reporter gene, then the fragment of the 5' UTR deleted or substituted plays a role in the regulation of the reporter gene expression.

[0210] Standard techniques known to those of skill in the art can be used to introduce mutations in the nucleotide sequence of an untranslated region of a target gene, including, for example, site-directed mutagenesis and PCR-mediated mutagenesis. In a specific embodiment, less than 75 nucleic acid residue substitutions, less than 50 nucleic acid residue substitutions, less than 45 nucleic acid residue substitutions, less than 40 nucleic acid residue substitutions, less than 35 nucleic acid residue substitutions, less than 30 nucleic acid residue substitutions, less than 25 nucleic acid residue substitutions, less than 20 nucleic acid residue substitutions, less than 15 nucleic acid residue substitutions, less than 10 nucleic acid residue substitutions, or less than 5 nucleic acid residue substitutions are introduced into the nucleotide sequence of an untranslated region of a target gene. In another embodiment, less than 10 elements of an untranslated region of a target gene, less than 9 of an untranslated region of a target gene, less than 8 elements of an untranslated region of a target gene, less than 7 elements of an untranslated region of a target gene, less than 6 elements of an untranslated region of a target gene, less than 5 elements of an untranslated region of a target gene, less than 4 elements of an untranslated region of a target gene, less than 3 elements of an untranslated region of a target gene, or less than 2 elements of an untranslated region of a target gene are mutated at one time.

### 5.5.3. Expressed Protein Concentration and Activity Assays



[0211] The compounds identified in the reporter gene-based assays described herein that modulate untranslated region-dependent expression may be tested in *in vitro* assays (*e.g.*, cell-free assays) or *in vivo* assays (*e.g.*, cell-based assays) well-known to one of skill in the art or described herein for the effect of said compounds on the expression of the target gene from which the untranslated regions of the reporter gene construct were derived. The specificity of a particular compound's effect on untranslated region-dependent expression of one or more other genes (preferably, a plurality of genes) can also be determined utilizing assays well-known to one of skill in the art or described herein. In a preferred embodiment, a compound identified utilizing the reporter gene-based assays described herein has a specific effect on the expression of only one gene or a group of genes within the same signaling pathway.

[0212] The expression of a gene can be readily detected, *e.g.*, by quantifying the protein and/or RNA encoded by said gene. Many methods standard in the art can be thus employed, including, but not limited to, immunoassays to detect and/or visualize gene expression (*e.g.*, western blot, immunoprecipitation followed by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE), immunocytochemistry, *etc.*) and/or hybridization assays to detect gene expression by detecting and/or visualizing respectively mRNA encoding a gene (*e.g.*, northern assays, dot blots, *in situ* hybridization, *etc.*). Such assays are routine and well known in the art (see, *e.g.*, Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York, which is incorporated by reference herein in its entirety). Exemplary immunoassays are described briefly below (but are not intended by way of limitation).

[0213] Immunoprecipitation protocols generally comprise lysing a population of cells in a lysis buffer such as RIPA buffer (1% NP-40 or Triton X-100, 1% sodium deoxycholate, 0.1% SDS, 0.15 M NaCl, 0.01 M sodium phosphate at pH 7.2, 1% Trasyolol) supplemented with protein phosphatase and/or protease inhibitors (*e.g.*, EDTA, PMSF, aprotinin, sodium vanadate), adding the antibody of interest to the cell lysate, incubating for a period of time (*e.g.*, 1 to 4 hours) at 40° C, adding protein A and/or protein G sepharose beads to the cell lysate, incubating for about an hour or more at 40° C, washing the beads in lysis buffer and resuspending the beads in SDS/sample buffer. The ability of the antibody of interest to immunoprecipitate a particular antigen can be assessed by, *e.g.*, western blot analysis. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the binding of the antibody to an antigen and decrease the background (*e.g.*, pre-clearing the cell lysate with sepharose beads). For further discussion regarding

immunoprecipitation protocols see, *e.g.*, Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.16.1.

**[0214]** Western blot analysis generally comprises preparing protein samples, electrophoresis of the protein samples in a polyacrylamide gel (*e.g.*, 8%- 20% SDS-PAGE depending on the molecular weight of the antigen), transferring the protein sample from the polyacrylamide gel to a membrane such as nitrocellulose, PVDF or nylon, blocking the membrane in blocking solution (*e.g.*, PBS with 3% BSA or non-fat milk), washing the membrane in washing buffer (*e.g.*, PBS-Tween 20), incubating the membrane with primary antibody (the antibody of interest) diluted in blocking buffer, washing the membrane in washing buffer, incubating the membrane with a secondary antibody (which recognizes the primary antibody, *e.g.*, an anti-human antibody) conjugated to an enzymatic substrate (*e.g.*, horseradish peroxidase or alkaline phosphatase) or radioactive molecule (*e.g.*,  $^{32}\text{P}$  or  $^{125}\text{I}$ ) diluted in blocking buffer, washing the membrane in wash buffer, and detecting the presence of the antigen. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected and to reduce the background noise. For further discussion regarding western blot protocols see, *e.g.*, Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.8.1.

**[0215]** ELISAs comprise preparing antigen, coating the well of a 96 well microtiter plate with the antigen, adding the antibody of interest conjugated to a detectable compound such as an enzymatic substrate (*e.g.*, horseradish peroxidase or alkaline phosphatase) to the well and incubating for a period of time, and detecting the presence of the antigen. In ELISAs the antibody of interest does not have to be conjugated to a detectable compound; instead, a second antibody (which recognizes the antibody of interest) conjugated to a detectable compound may be added to the well. Further, instead of coating the well with the antigen, the antibody may be coated to the well. In this case, a second antibody conjugated to a detectable compound may be added following the addition of the antigen of interest to the coated well. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected as well as other variations of ELISAs known in the art. For further discussion regarding ELISAs see, *e.g.*, Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 11.2.1.

**[0216]** Another antibody based separation that can be used to detect the protein of interest is the use of flow cytometry such as by a fluorescence activated cell sorter ("FACS"). Typically, separation by flow cytometry is performed as follows. The suspended mixture of

cells are centrifuged and resuspended in media. Antibodies which are conjugated to fluorochrome are added to allow the binding of the antibodies to specific proteins. In another embodiment, the secondary antibodies that are conjugated to fluorochromes can be used to detect primary antibodies specific to the protein of interest. The cell mixture is then washed by one or more centrifugation and resuspension steps. The mixture is run through a FACS which separates the cells based on different fluorescence characteristics. FACS systems are available in varying levels of performance and ability, including multi-color analysis. The facilitating cell can be identified by a characteristic profile of forward and side scatter which is influenced by size and granularity, as well as by positive and/or negative expression of certain cell surface markers.

[0217] In addition to measuring the effect of a compound identified in the reporter gene-based assays described herein on the expression of the target gene from which the untranslated regions of the reporter gene construct were derived, the activity of the protein encoded by the target gene can be assessed utilizing techniques well-known to one of skill in the art. For example, the activity of a protein encoded by a target gene can be determined by detecting induction of a cellular second messenger (*e.g.*, intracellular  $\text{Ca}^{2+}$ , diacylglycerol,  $\text{IP}_3$ , *etc.*), detecting the phosphorylation of a protein, detecting the activation of a transcription factor, or detecting a cellular response, for example, cellular differentiation, or cell proliferation. The induction of a cellular second messenger or phosphorylation of a protein can be determined by, *e.g.*, immunoassays well-known to one of skill in the art and described herein. The activation of a transcription factor can be detected by, *e.g.*, electromobility shift assays, and a cellular response such as cellular proliferation can be detected by, *e.g.*, trypan blue cell counts,  $^3\text{H}$ -thymidine incorporation, and flow cytometry.

#### **5.6. Methods for Characterizing the Compounds That Modulate Untranslated Region-Dependent Expression of a Target Gene**

[0218] If the library comprises arrays or microarrays of compounds, wherein each compound has an address or identifier, the compound can be deconvoluted, *e.g.*, by cross-referencing the positive sample to original compound list that was applied to the individual test assays.

[0219] If the library is a peptide or nucleic acid library, the sequence of the compound can be determined by direct sequencing of the peptide or nucleic acid. Such methods are well known to one of skill in the art.

[0220] A number of physico-chemical techniques can be used for the *de novo* characterization of compounds bound to the target RNA. Examples of such techniques

include, but are not limited to, mass spectrometry, NMR spectroscopy, X-ray crystallography and vibrational spectroscopy.

#### **5.6.1. Mass Spectrometry**

[0221] Mass spectrometry (*e.g.*, electrospray ionization (“ESI”), matrix-assisted laser desorption-ionization (“MALDI”), and Fourier-transform ion cyclotron resonance (“FT-ICR”) can be used for elucidating the structure of a compound.

[0222] MALDI uses a pulsed laser for desorption of the ions and a time-of-flight analyzer, and has been used for the detection of noncovalent tRNA:amino-acyl-tRNA synthetase complexes (Gruic-Sovulj et al., 1997, J. Biol. Chem. 272:32084-32091). However, covalent cross-linking between the target nucleic acid and the compound is required for detection, since a non-covalently bound complex may dissociate during the MALDI process.

[0223] ESI mass spectrometry (“ESI-MS”) has been of greater utility for studying non-covalent molecular interactions because, unlike the MALDI process, ESI-MS generates molecular ions with little to no fragmentation (Xavier et al., 2000, Trends Biotechnol. 18(8):349-356). ESI-MS has been used to study the complexes formed by HIV Tat peptide and protein with the TAR RNA (Sannes-Lowery et al., 1997, Anal. Chem. 69:5130-5135).

[0224] Fourier-transform ion cyclotron resonance (“FT-ICR”) mass spectrometry provides high-resolution spectra, isotope-resolved precursor ion selection, and accurate mass assignments (Xavier et al., 2000, Trends Biotechnol. 18(8):349-356). FT-ICR has been used to study the interaction of aminoglycoside antibiotics with cognate and non-cognate RNAs (Hofstadler et al., 1999, Anal. Chem. 71:3436-3440; and Griffey et al., 1999, Proc. Natl. Acad. Sci. USA 96:10129-10133). As true for all of the mass spectrometry methods discussed herein, FT-ICR does not require labeling of the target RNA or a compound.

[0225] An advantage of mass spectroscopy is not only the elucidation of the structure of the compound, but also the determination of the structure of the compound bound to a target RNA. Such information can enable the discovery of a consensus structure of a compound that specifically binds to a target RNA.

#### **5.6.2. NMR Spectroscopy**

[0226] NMR spectroscopy is a valuable technique for identifying complexed target nucleic acids by qualitatively determining changes in chemical shift, specifically from distances measured using relaxation effects, and NMR-based approaches have been used in the identification of small molecule binders of protein drug targets (Xavier et al., 2000,

Trends Biotechnol. 18(8):349-356). The determination of structure-activity relationships ("SAR") by NMR is the first method for NMR described in which small molecules that bind adjacent subsites are identified by two-dimensional  $^1\text{H}$ - $^{15}\text{N}$  spectra of the target protein (Shuker et al., 1996, Science 274:1531-1534). The signal from the bound molecule is monitored by employing line broadening, transferred NOEs and pulsed field gradient diffusion measurements (Moore, 1999, Curr. Opin. Biotechnol. 10:54-58). A strategy for lead generation by NMR using a library of small molecules has been recently described (Fejzo et al., 1999, Chem. Biol. 6:755-769).

[0227] In one embodiment of the present invention, the target nucleic acid complexed to a compound can be determined by SAR by NMR. Furthermore, SAR by NMR can also be used to elucidate the structure of a compound.

[0228] As described above, NMR spectroscopy is a technique for identifying binding sites in target nucleic acids by qualitatively determining changes in chemical shift, specifically from distances measured using relaxation effects. Examples of NMR that can be used for the invention include, but are not limited to, one-dimensional NMR, two-dimensional NMR, correlation spectroscopy ("COSY"), and nuclear Overhauser effect ("NOE") spectroscopy. Such methods of structure determination of compounds are well-known to one of skill in the art.

[0229] Similar to mass spectroscopy, an advantage of NMR is the not only the elucidation of the structure of the compound, but also the determination of the structure of the compound bound to the target RNA. Such information can enable the discovery of a consensus structure of a compound that specifically binds to a target RNA.

### 5.6.3. X ray Crystallography

[0230] X-ray crystallography can be used to elucidate the structure of a compound. For a review of x-ray crystallography see, *e.g.*, Blundell et al., 2002, Nat Rev Drug Discov 1(1):45-54. The first step in x-ray crystallography is the formation of crystals. The formation of crystals begins with the preparation of highly purified and soluble samples. The conditions for crystallization are then determined by optimizing several solution variables known to induce nucleation, such as pH, ionic strength, temperature, and specific concentrations of organic additives, salts and detergent. Techniques for automating the crystallization process have been developed for the production of high-quality protein crystals. Once crystals have been formed, the crystals are harvested and prepared for data collection. The crystals are then analyzed by diffraction (such as multi-circle diffractometers, high-speed CCD detectors, and detector off-set). Generally, multiple crystals must be screened for structure determinations.

#### **5.6.4. Vibrational Spectroscopy**

[0231] Vibrational spectroscopy (*e.g.* infrared (IR) spectroscopy or Raman spectroscopy) can be used for elucidating the structure of a compound.

[0232] Infrared spectroscopy measures the frequencies of infrared light (wavelengths from 100 to 10,000 nm) absorbed by the compound as a result of excitation of vibrational modes according to quantum mechanical selection rules which require that absorption of light cause a change in the electric dipole moment of the molecule. The infrared spectrum of any molecule is a unique pattern of absorption wavelengths of varying intensity that can be considered as a molecular fingerprint to identify any compound.

[0233] Infrared spectra can be measured in a scanning mode by measuring the absorption of individual frequencies of light, produced by a grating which separates frequencies from a mixed-frequency infrared light source, by the compound relative to a standard intensity (double-beam instrument) or pre-measured ("blank") intensity (single-beam instrument). In a preferred embodiment, infrared spectra are measured in a pulsed mode ("FT-IR") where a mixed beam, produced by an interferometer, of all infrared light frequencies is passed through or reflected off the compound. The resulting interferogram, which may or may not be added with the resulting interferograms from subsequent pulses to increase the signal strength while averaging random noise in the electronic signal, is mathematically transformed into a spectrum using Fourier Transform or Fast Fourier Transform algorithms.

[0234] Raman spectroscopy measures the difference in frequency due to absorption of infrared frequencies of scattered visible or ultraviolet light relative to the incident beam. The incident monochromatic light beam, usually a single laser frequency, is not truly absorbed by the compound but interacts with the electric field transiently. Most of the light scattered off the sample will be unchanged (Rayleigh scattering) but a portion of the scatter light will have frequencies that are the sum or difference of the incident and molecular vibrational frequencies. The selection rules for Raman (inelastic) scattering require a change in polarizability of the molecule. While some vibrational transitions are observable in both infrared and Raman spectrometry, must are observable only with one or the other technique. The Raman spectrum of any molecule is a unique pattern of absorption wavelengths of varying intensity that can be considered as a molecular fingerprint to identify any compound.

[0235] Raman spectra are measured by submitting monochromatic light to the sample, either passed through or preferably reflected off, filtering the Rayleigh scattered light, and detecting the frequency of the Raman scattered light. An improved Raman spectrometer is

described in US Patent No. 5,786,893 to Fink et al., which is hereby incorporated by reference.

[0236] Vibrational microscopy can be measured in a spatially resolved fashion to address single beads by integration of a visible microscope and spectrometer. A microscopic infrared spectrometer is described in U.S. Patent No. 5,581,085 to Reffner et al., which is hereby incorporated by reference in its entirety. An instrument that simultaneously performs a microscopic infrared and microscopic Raman analysis on a sample is described in U.S. Patent No. 5,841,139 to Sostek et al., which is hereby incorporated by reference in its entirety.

[0237] In one embodiment of the method, compounds are synthesized on polystyrene beads doped with chemically modified styrene monomers such that each resulting bead has a characteristic pattern of absorption lines in the vibrational (IR or Raman) spectrum, by methods including but not limited to those described by Fenniri et al., 2000, J. Am. Chem. Soc. 123:8151-8152. Using methods of split-pool synthesis familiar to one of skill in the art, the library of compounds is prepared so that the spectroscopic pattern of the bead identifies one of the components of the compound on the bead. Beads that have been separated according to their ability to bind target RNA can be identified by their vibrational spectrum. In one embodiment of the method, appropriate sorting and binning of the beads during synthesis then allows identification of one or more further components of the compound on any one bead. In another embodiment of the method, partial identification of the compound on a bead is possible through use of the spectroscopic pattern of the bead with or without the aid of further sorting during synthesis, followed by partial resynthesis of the possible compounds aided by doped beads and appropriate sorting during synthesis.

[0238] In another embodiment, the IR or Raman spectra of compounds are examined while the compound is still on a bead, preferably, or after cleavage from bead, using methods including but not limited to photochemical, acid, or heat treatment. The compound can be identified by comparison of the IR or Raman spectral pattern to spectra previously acquired for each compound in the combinatorial library.

### **5.7. Secondary Screens of Compounds**

[0239] Once a compound has been identified to modulate untranslated region-dependent expression of a target gene and preferably, the structure of the compound has been identified by the methods described in Section 5.6, the compounds are tested for biological activity in further assays and/or animal models (see, *e.g.*, Sections 5.7.1 and 5.7.2). Further, a lead compound may be used to design congeners or analogs (see, *e.g.*, Section 5.7.3).

### 5.7.1. Cell-based Screens

[0240] The compounds identified in the assays described *supra* (for convenience referred to herein as a “lead” compound) can be tested for biological activity using host cells containing or engineered to contain the target RNA element involved in untranslated region-dependent gene expression coupled to a functional readout system. For example, a phenotypic or physiological readout can be used to assess untranslated region-dependent activity of the target RNA in the presence and absence of the lead compound.

[0241] In one embodiment, a phenotypic or physiological readout can be used to assess untranslated region-dependent activity of the target RNA in the presence and absence of the lead compound. For example, the target RNA may be overexpressed in a cell in which the target RNA is endogenously expressed. Where the target RNA controls untranslated region-dependent expression of a gene product involved in cell growth or viability, the *in vivo* effect of the lead compound can be assayed by measuring the cell growth or viability of the target cell. Such assays can be carried out with representative cells of cell types involved in a particular disease or disorder (*e.g.*, leukocytes such as T cells, B cells, natural killer cells, macrophages, neutrophils and eosinophils). A lower level of proliferation or survival of the contacted cells indicates that the lead compound is effective to treat a condition in the patient characterized by uncontrolled cell growth. Alternatively, instead of culturing cells from a patient, a lead compound may be screened using cells of a tumor or malignant cell line or an endothelial cell line. Specific examples of cell culture models include, but are not limited to, for lung cancer, primary rat lung tumor cells (see, *e.g.*, Swafford et al., 1997, Mol. Cell. Biol., 17:1366-1374) and large-cell undifferentiated cancer cell lines (see, *e.g.*, Mabry et al., 1991, Cancer Cells, 3:53-58); colorectal cell lines for colon cancer (see, *e.g.*, Park & Gazdar, 1996, J. Cell Biochem. Suppl. 24:131-141); multiple established cell lines for breast cancer (see, *e.g.*, Hambly et al., 1997, Breast Cancer Res. Treat. 43:247-258; Gierthy et al., 1997, Chemosphere 34:1495-1505; and Prasad & Church, 1997, Biochem. Biophys. Res. Commun. 232:14-19); a number of well-characterized cell models for prostate cancer (see, *e.g.*, Webber et al., 1996, Prostate, Part 1, 29:386-394; Part 2, 30:58-64; and Part 3, 30:136-142 and Boulikas, 1997, Anticancer Res. 17:1471-1505); for genitourinary cancers, continuous human bladder cancer cell lines (see, *e.g.*, Ribeiro et al., 1997, Int. J. Radiat. Biol. 72:11-20); organ cultures of transitional cell carcinomas (see, *e.g.*, Booth et al., 1997, Lab Invest. 76:843-857) and rat progression models (see, *e.g.*, Vet et al., 1997, Biochim. Biophys Acta 1360:39-44); and established cell lines for leukemias and lymphomas (see, *e.g.*, Drexler, 1994, Leuk. Res. 18:919-927 and Tohyama, 1997, Int. J. Hematol. 65:309-317).



[0242] Many assays well-known in the art can be used to assess the survival and/or growth of a patient cell or cell line following exposure to a lead compound; for example, cell proliferation can be assayed by measuring bromodeoxyuridine (BrdU) incorporation (see, *e.g.*, Hoshino et al., 1986, *Int. J. Cancer* 38:369 and Campana et al., 1988, *J. Immunol. Meth.* 107:79) or (<sup>3</sup>H)-thymidine incorporation (see, *e.g.*, Chen, 1996, *Oncogene* 13:1395-403 and Jeoung, 1995, *J. Biol. Chem.* 270:18367-73), by direct cell count, by detecting changes in transcription, translation or activity of known genes such as proto-oncogenes (*e.g.*, *fos*, *myc*) or cell cycle markers (Rb, *cdc2*, cyclin A, D1, D2, D3, E, *etc.*). The levels of such protein and mRNA and activity can be determined by any method well known in the art. For example, protein can be quantitated by known immunodiagnostic methods such as western blotting or immunoprecipitation using commercially available antibodies. mRNA can be quantitated using methods that are well known and routine in the art, for example, using northern analysis, RNase protection, the polymerase chain reaction in connection with reverse transcription ("RT-PCR"). Cell viability can be assessed by using trypan-blue staining or other cell death or viability markers known in the art. In a specific embodiment, the level of cellular ATP is measured to determine cell viability.

Differentiation can be assessed, for example, visually based on changes in morphology.

[0243] The lead compound can also be assessed for its ability to inhibit cell transformation (or progression to malignant phenotype) *in vitro*. In this embodiment, cells with a transformed cell phenotype are contacted with a lead compound, and examined for change in characteristics associated with a transformed phenotype (a set of *in vitro* characteristics associated with a tumorigenic ability *in vivo*), for example, but not limited to, colony formation in soft agar, a more rounded cell morphology, looser substratum attachment, loss of contact inhibition, loss of anchorage dependence, release of proteases such as plasminogen activator, increased sugar transport, decreased serum requirement, or expression of fetal antigens, *etc.* (see, *e.g.*, Luria et al., 1978, *General Virology*, 3d Ed., John Wiley & Sons, New York, pp. 436-446).

[0244] Loss of invasiveness or decreased adhesion can also be assessed to demonstrate the anti-cancer effects of a lead compound. For example, an aspect of the formation of a metastatic cancer is the ability of a precancerous or cancerous cell to detach from primary site of disease and establish a novel colony of growth at a secondary site. The ability of a cell to invade peripheral sites reflects its potential for a cancerous state. Loss of invasiveness can be measured by a variety of techniques known in the art including, for example, induction of E-cadherin-mediated cell-cell adhesion. Such E-cadherin-mediated

adhesion can result in phenotypic reversion and loss of invasiveness (see, *e.g.*, Hordijk et al., 1997, Science 278:1464-66).

[0245] Loss of invasiveness can further be examined by inhibition of cell migration. A variety of 2-dimensional and 3-dimensional cellular matrices are commercially available (Calbiochem-Novabiochem Corp. San Diego, CA). Cell migration across or into a matrix can be examined using microscopy, time-lapsed photography or videography, or by any method in the art allowing measurement of cellular migration. In a related embodiment, loss of invasiveness is examined by response to hepatocyte growth factor ("HGF"). HGF-induced cell scattering is correlated with invasiveness of cells such as Madin-Darby canine kidney ("MDCK") cells. This assay identifies a cell population that has lost cell scattering activity in response to HGF (see, *e.g.*, Hordijk et al., 1997, Science 278:1464-66).

[0246] Alternatively, loss of invasiveness can be measured by cell migration through a chemotaxis chamber (Neuroprobe/ Precision Biochemicals Inc. Vancouver, BC). In such assay, a chemo-attractant agent is incubated on one side of the chamber (*e.g.*, the bottom chamber) and cells are plated on a filter separating the opposite side (*e.g.*, the top chamber). In order for cells to pass from the top chamber to the bottom chamber, the cells must actively migrate through small pores in the filter. Checkerboard analysis of the number of cells that have migrated can then be correlated with invasiveness (see *e.g.*, Ohnishi, 1993, Biochem. Biophys. Res. Commun. 193:518-25).

[0247] The effect of a compound of the invention on cell adhesion can be measured using HUVECS. HUVECs are seeded on 24 well culture plates and incubated for 2 days to allow formation of a confluent monolayer. Cancerous cells or a cancer cell line such as LS-180 human colon adenocarcinoma cells are labeled with 5  $\mu$ M Calcein-AM for 30 min. Calcein-AM labeled LS180 cells are added into each well of the HUVEC culture; and incubated for 10 min at 37°C. TNF- $\alpha$  (80 ng/ml) is then added and the culture incubated for an additional 110 min. Non-adherent cells are removed by washing with PBS. The fluorescence intensity of adherent LS-180 cell in each individual well is measured by a fluorescent plate reader set at excitation 485/20 nm and emission at 530/25 nm.

[0248] The effect of a compound of the invention on cell migration and invasion can also be determined using an assay based on the BD BioCoast Angiogenesis System (BD Biosciences, Bedford, MA). The fluorescence blocking membrane of the insert is a 3 micron pore size PET filter which has been coated either with BD Matrigel basement matrix (for invasion assay) or without Matrigel matrix (for migration assay). HUVECs (250  $\mu$ l/well) in culture medium without serum are added to the top chamber; a compound added to bottom wells containing medium (750  $\mu$ l/well) with VEGF as a chemo-attractant. Cells

are then incubated for 22 h at 37°C. After incubation, cells are stained with Calcein AM for measurement of fluorescence.

[0249] Further, a lead compound can be assessed for its ability to alter viral replication (as determined, *e.g.*, by plaque formation) or the production of viral proteins (as determined, *e.g.*, by western blot analysis) or viral RNAs (as determined, *e.g.*, by RT-PCR or northern blot analysis) in cultured cells *in vitro* using methods which are well known in the art. A lead compound can also be assessed for its ability to alter bacterial replication (as determined, *e.g.*, by measuring bacterial growth rates) or the production of bacterial proteins (as determined, *e.g.*, by western blot analysis) or bacterial RNAs (as determined, *e.g.*, by RT-PCR or northern blot analysis) in cultured cells *in vitro* using methods which are well known in the art. Finally, a lead compound can be assessed for its ability to alter fungal replication (as determined, *e.g.*, by fungal growth rates, such as macrodilution methods and/or microdilution methods using protocols known to those skilled in the art (see, *e.g.*, Clancy et al., 1997, Journal of Clinical Microbiology, 35(11): 2878-82; Ryder et al., 1998, Antimicrobial Agents and Chemotherapy, 42(5): 1057-61; or U.S. Patent Nos. 5,521,153; U.S. 5,883,120, U.S. 5,521,169)) or the production of fungal proteins (as determined, *e.g.*, by western blot analysis) or fungal RNAs (as determined, *e.g.*, by RT-PCR or northern blot analysis) in cultured cells *in vitro* using methods which are well known in the art.

### 5.7.2. Animal Model-based Screens

[0250] The lead compounds identified in the reporter gene-based assay described herein can be tested for biological activity using animal models for a disease, disorder, condition, or syndrome of interest. These include animals engineered to contain a target gene coupled to a functional readout system, such as a transgenic mouse. Such animal model systems include, but are not limited to, rats, mice, chicken, cows, monkeys, pigs, dogs, rabbits, *etc.* In a specific embodiment of the invention, a compound identified in accordance with the methods of the invention is tested in a mouse model system. Such model systems are widely used and well-known to the skilled artisan such as the SCID mouse model or transgenic mice.

[0251] The anti-angiogenic activity of a compound identified in accordance with the invention can be determined by using various experimental animal models of vascularized tumors. The anti-tumor activity of a compound identified in accordance with the invention can be determined by administering the compound to an animal model and verifying that the compound is effective in reducing the proliferation or spread of cancer cells in said animal model. An example of an animal model for human cancer in general includes, but is

not limited to, spontaneously occurring tumors of companion animals (see, *e.g.*, Vail & MacEwen, 2000, *Cancer Invest* 18(8):781-92).

**[0252]** Examples of animal models for lung cancer include, but are not limited to, lung cancer animal models described by Zhang & Roth (1994, *In Vivo* 8(5):755-69) and a transgenic mouse model with disrupted p53 function (see, *e.g.*, Morris et al., 1998, *J La State Med Soc* 150(4):179-85). An example of an animal model for breast cancer includes, but is not limited to, a transgenic mouse that overexpresses cyclin D1 (see, *e.g.*, Hosokawa et al., 2001, *Transgenic Res* 10(5):471-8). An example of an animal model for colon cancer includes, but is not limited to, a TCR b and p53 double knockout mouse (see, *e.g.*, Kado et al., 2001, *Cancer Res* 61(6):2395-8). Examples of animal models for pancreatic cancer include, but are not limited to, a metastatic model of Panc02 murine pancreatic adenocarcinoma (see, *e.g.*, Wang et al., 2001, *Int J Pancreatol* 29(1):37-46) and nu-nu mice generated in subcutaneous pancreatic tumours (see, *e.g.*, Ghaneh et al., 2001, *Gene Ther* 8(3):199-208). Examples of animal models for non-Hodgkin's lymphoma include, but are not limited to, a severe combined immunodeficiency ("SCID") mouse (see, *e.g.*, Bryant et al., 2000, *Lab Invest* 80(4):553-73) and an IgHmu-HOX11 transgenic mouse (see, *e.g.*, Hough et al., 1998, *Proc Natl Acad Sci USA* 95(23):13853-8). An example of an animal model for esophageal cancer includes, but is not limited to, a mouse transgenic for the human papillomavirus type 16 E7 oncogene (see, *e.g.*, Herber et al., 1996, *J Virol* 70(3):1873-81). Examples of animal models for colorectal carcinomas include, but are not limited to, Apc mouse models (see, *e.g.*, Fodde & Smits, 2001, *Trends Mol Med* 7(8):369-73 and Kuraguchi et al., 2000, *Oncogene* 19(50):5755-63).

**[0253]** The anti-inflammatory activity of a compound identified in accordance with the invention can be determined by using various experimental animal models of inflammatory arthritis known in the art and described in Crofford & Wilder, "Arthritis and Autoimmunity in Animals", in *Arthritis and Allied Conditions: A Textbook of Rheumatology*, McCarty et al.(eds.), Chapter 30 (Lee & Febiger, 1993). Experimental and spontaneous animal models of inflammatory arthritis and autoimmune rheumatic diseases can also be used to assess the anti-inflammatory activity of a compound identified in accordance with the invention. The principle animal models for arthritis or inflammatory disease known in the art and widely used include: adjuvant-induced arthritis rat models, collagen-induced arthritis rat and mouse models and antigen-induced arthritis rat, rabbit and hamster models, all described in Crofford & Wilder, "Arthritis and Autoimmunity in Animals", in *Arthritis and Allied Conditions: A Textbook of Rheumatology*, McCarty et al.(eds.), Chapter 30 (Lee & Febiger, 1993), incorporated herein by reference in its entirety.

[0254] The anti-inflammatory activity of a compound identified in accordance with the invention can be assessed using a carrageenan-induced arthritis rat model. Carrageenan-induced arthritis has also been used in rabbit, dog and pig in studies of chronic arthritis or inflammation. Quantitative histomorphometric assessment is used to determine therapeutic efficacy. The methods for using such a carrageenan-induced arthritis model is described in Hansra et al., 2000, *Inflammation*, 24(2): 141-155. Also commonly used are zymosan-induced inflammation animal models as known and described in the art.

[0255] The anti-inflammatory activity of a compound identified in accordance with the invention can also be assessed by measuring the inhibition of carrageenan-induced paw edema in the rat, using a modification of the method described in Winter et al., 1962, *Proc. Soc. Exp. Biol Med.* 111, 544-547. This assay has been used as a primary *in vivo* screen for the anti-inflammatory activity of most NSAIDs, and is considered predictive of human efficacy. The anti-inflammatory activity of a compound identified in accordance with the invention is expressed as the percent inhibition of the increase in hind paw weight of the test group relative to the vehicle dosed control group.

[0256] In a specific embodiment of the invention where the experimental animal model used is adjuvant-induced arthritis rat model, body weight can be measured relative to a control group to determine the anti-inflammatory activity of a compound identified in accordance with the invention. Alternatively, the efficacy of a compound identified in accordance with the invention can be assessed using assays that determine bone loss. Animal models such as ovariectomy-induced bone resorption mice, rat and rabbit models are known in the art for obtaining dynamic parameters for bone formation. Using methods such as those described by Yositate et al. or Yamamoto et al., bone volume is measured *in vivo* by microcomputed tomography analysis and bone histomorphometry analysis (see, e.g., Yositate et al., 1999, *Proc. Natl. Acad. Sci.* 96:8156-8160 and Yamamoto et al., 1998, *Endocrinology* 139(3):1411-1419, both incorporated herein by reference in their entireties).

[0257] Additionally, animal models for inflammatory bowel disease can also be used to assess the efficacy of a compound identified in accordance with the invention (see, e.g., Kim et al., 1992, *Scand. J. Gastroentrol.* 27:529-537 and Strober, 1985, *Dig. Dis. Sci.* 30(12 Suppl):3S-10S). Ulcerative colitis and Crohn's disease are human inflammatory bowel diseases that can be induced in animals. Sulfated polysaccharides including, but not limited to, amylopectin, carrageen, amylopectin sulfate, and dextran sulfate or chemical irritants including, but not limited to, trinitrobenzenesulphonic acid (TNBS) and acetic acid can be administered to animals orally to induce inflammatory bowel diseases.

[0258] Animal models for asthma can also be used to assess the efficacy of a compound identified in accordance with the invention. An example of one such model is the murine adoptive transfer model in which aeroallergen provocation of TH1 or TH2 recipient mice results in TH effector cell migration to the airways and is associated with an intense neutrophilic (TH1) and eosinophilic (TH2) lung mucosal inflammatory response (see, *e.g.*, Cohn et al., 1997, J. Exp. Med. 186:1737-1747).

[0259] Animal models for autoimmune disorders can also be used to assess the efficacy of a compound identified in accordance with the invention. Animal models for autoimmune disorders such as type 1 diabetes, thyroid autoimmunity, systemic lupus erythematosus, and glomerulonephritis have been developed (see, *e.g.*, Flanders et al., 1999, Autoimmunity 29:235-246; Krogh et al., 1999, Biochimie 81:511-515; and Foster, 1999, Semin. Nephrol. 19:12-24).

[0260] Animal models for viral infections can also be used to assess the efficacy of a compound identified in accordance with the invention. Animal models for viral infections such as EBV-associated diseases, gammaherpesviruses, infectious mononucleosis, simian immunodeficiency virus ("SIV"), Borna disease virus infection, hepatitis, varicella virus infection, viral pneumonitis, Epstein-Barr virus pathogenesis, feline immunodeficiency virus ("FIV"), HTLV type 1 infection, human rotaviruses, and genital herpes have been developed (see, *e.g.*, Hayashi et al., 2002, Histol Histopathol 17(4):1293-310; Arico et al., 2002, J Interferon Cytokine Res 22(11):1081-8; Flano et al., 2002, Immunol Res 25(3):201-17; Sauermann, 2001, Curr Mol Med 1(4):515-22; Pletnikov et al., 2002, Front Biosci 7:d593-607; Engler et al., 2001, Mol Immunol 38(6):457-65; White et al., 2001, Brain Pathol 11(4):475-9; Davis & Matalon, 2001, News Physiol Sci 16:185-90; Wang, 2001, Curr Top Microbiol Immunol. 258:201-19; Phillips et al., 2000, J Psychopharmacol. 14(3):244-50; Kazanji, 2000, AIDS Res Hum Retroviruses. 16(16):1741-6; Saif et al., 1996, Arch Virol Suppl. 12:153-61; and Hsiung et al., 1984, Rev Infect Dis. 6(1):33-50).

[0261] Animal models for bacterial infections can also be used to assess the efficacy of a compound identified in accordance with the invention. Animal models for bacterial infections such as *H. pylori*-infection, genital mycoplasmosis, primary sclerosing cholangitis, cholera, chronic lung infection with *Pseudomonas aeruginosa*, Legionnaires' disease, gastroduodenal ulcer disease, bacterial meningitis, gastric *Helicobacter* infection, pneumococcal otitis media, experimental allergic neuritis, leprous neuropathy, mycobacterial infection, endocarditis, Aeromonas-associated enteritis, *Bacteroides fragilis* infection, syphilis, streptococcal endocarditis, acute hematogenous osteomyelitis, human scrub typhus, toxic shock syndrome, anaerobic infections, *Escherichia coli* infections, and

"*Mycoplasma pneumoniae* infections have been developed (see, e.g., Sugiyama et al., 2002, J Gastroenterol. 37 Suppl 13:6-9; Brown et al., 2001, Am J Reprod Immunol. 46(3):232-41; Vierling, 2001, Best Pract Res Clin Gastroenterol. 15(4):591-610; Klose, 2000, Trends Microbiol. 8(4):189-91; Stotland et al., 2000, Pediatr Pulmonol. 30(5):413-24; Brieland et al., 2000, Immunopharmacology 48(3):249-52; Lee, 2000, Baillieres Best Pract Res Clin Gastroenterol. 14(1):75-96; Koedel & Pfister, 1999, Infect Dis Clin North Am. 13(3):549-77; Nedrud, 1999, FEMS Immunol Med Microbiol. 24(2):243-50; Prellner et al., 1999, Microb Drug Resist. 5(1):73-82; Vriesendorp, 1997, J Infect Dis. 176 Suppl 2:S164-8; Shetty & Antia, 1996, Indian J Lepr. 68(1):95-104; Balasubramanian et al., 1994, Immunobiology 191(4-5):395-401; Carbon et al., 1994, Int J Biomed Comput. 36(1-2):59-67; Haberberger et al., 1991, Experientia. 47(5):426-9; Onderdonk et al., 1990, Rev Infect Dis. 12 Suppl 2:S169-77; Wicher & Wicher, 1989, Crit Rev Microbiol. 16(3):181-234; Scheld, 1987, J Antimicrob Chemother. 20 Suppl A:71-85; Emslie & Nade, 1986, Rev Infect Dis. 8(6):841-9; Ridgway et al., 1986, Lab Anim Sci. 36(5):481-5; Quimby & Nguyen, 1985, Crit Rev Microbiol. 12(1):1-44; Onderdonk et al., 1979, Rev Infect Dis. 1(2):291-301; Smith, 1976, Ciba Found Symp. (42):45-72, and Taylor-Robinson, 1976, Infection. 4(1 Suppl):4-8).

[0262] Animal models for fungal infections can also be used to assess the efficacy of a compound identified in accordance with the invention. Animal models for fungal infections such as *Candida* infections, zygomycosis, *Candida* mastitis, progressive disseminated trichosporonosis with latent trichosporonemia, disseminated candidiasis, pulmonary paracoccidioidomycosis, pulmonary aspergillosis, *Pneumocystis carinii* pneumonia, cryptococcal meningitis, coccidioidal meningoencephalitis and cerebrospinal vasculitis, *Aspergillus niger* infection, *Fusarium* keratitis, paranasal sinus mycoses, *Aspergillus fumigatus* endocarditis, tibial dyschondroplasia, *Candida glabrata* vaginitis, oropharyngeal candidiasis, X-linked chronic granulomatous disease, tinea pedis, cutaneous candidiasis, mycotic placentitis, disseminated trichosporonosis, allergic bronchopulmonary aspergillosis, mycotic keratitis, *Cryptococcus neoformans* infection, fungal peritonitis, *Curvularia geniculata* infection, staphylococcal endophthalmitis, sporotrichosis, and dermatophytosis have been developed (see, e.g., Arendrup et al., 2002, Infection 30(5):286-91; Kamei, 2001, Mycopathologia 152(1):5-13; Guhad et al., 2000, FEMS Microbiol Lett. 192(1):27-31; Yamagata et al., 2000, J Clin Microbiol. 38(9):32606; Andrutis et al., 2000, J Clin Microbiol. 38(6):2317-23; Cock et al., 2000, Rev Inst Med Trop Sao Paulo 42(2):59-66; Shibuya et al., 1999, Microb Pathog. 27(3):123-31; Beers et al., 1999, J Lab Clin Med. 133(5):423-33; Najvar et al., 1999, Antimicrob Agents Chemother. 43(2):413-4;

Williams et al., 1988, J Infect Dis. 178(4):1217-21; Yoshida, 1988, Kansenshogaku Zasshi. 1998 Jun;72(6):621-30; Alexandrakis et al., 1998, Br J Ophthalmol. 82(3):306-11; Chakrabarti et al., 1997, J Med Vet Mycol. 35(4):295-7; Martin et al., 1997, Antimicrob Agents Chemother. 41(1):13-6; Chu et al., 1996, Avian Dis. 40(3):715-9; Fidel et al., 1996, J Infect Dis. 173(2):425-31; Cole et al., 1995, FEMS Microbiol Lett. 15;126(2):177-80; Pollock et al., 1995, Nat Genet. 9(2):202-9; Uchida et al., 1994, Jpn J Antibiot. 47(10):1407-12; : Maebashi et al., 1994, J Med Vet Mycol. 32(5):349-59; Jensen & Schonheyder, 1993, J Exp Anim Sci. 35(4):155-60; Gokaslan & Anaissie, 1992, Infect Immun. 60(8):3339-44; Kurup et al., 1992, J Immunol. 148(12):3783-8; Singh et al., 1990, Mycopathologia. 112(3):127-37; Salkowski & Balish, 1990, Infect Immun. 58(10):3300-6; Ahmad et al., 1986, Am J Kidney Dis. 7(2):153-6; Altire-Werber E, Edberg SC, 1985, Mycopathologia. 89(2):69-73; Kane et al., 1981, Antimicrob Agents Chemother. 20(5):595-9; Barbee et al., 1977, Am J Pathol. 86(1):281-4; and Maestrone et al., 1973, Am J Vet Res. 34(6):833-6).

**[0263]** The toxicity and/or efficacy of a compound identified in accordance with the invention can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD<sub>50</sub> (the dose lethal to 50% of the population) and the ED<sub>50</sub> (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD<sub>50</sub>/ED<sub>50</sub>. A compound identified in accordance with the invention that exhibits large therapeutic indices is preferred. While a compound identified in accordance with the invention that exhibits toxic side effects may be used, care should be taken to design a delivery system that targets such agents to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

**[0264]** The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage of a compound identified in accordance with the invention for use in humans. The dosage of such agents lies preferably within a range of circulating concentrations that include the ED<sub>50</sub> with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any agent used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC<sub>50</sub> (*i.e.*, the concentration of the compound that achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately



determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

### **5.7.3. Design of Congeners or Analogs**

[0265] The compounds which display the desired biological activity can be used as lead compounds for the development or design of congeners or analogs having useful pharmacological activity. For example, once a lead compound is identified, molecular modeling techniques can be used to design variants of the compound that can be more effective. Examples of molecular modeling systems are the CHARM and QUANTA programs (Polygen Corporation, Waltham, MA). CHARM performs the energy minimization and molecular dynamics functions. QUANTA performs the construction, graphic modelling and analysis of molecular structure. QUANTA allows interactive construction, modification, visualization, and analysis of the behavior of molecules with each other.

[0266] A number of articles review computer modeling of drugs interactive with specific proteins, such as Rotivinen et al., 1988, Acta Pharmaceutica Fennica 97:159-166; Ripka, 1998, New Scientist 54-57; McKinaly & Rossmann, 1989, Annu. Rev. Pharmacol. Toxicol. 29:111-122; Perry & Davies, OSAR: Quantitative Structure-Activity Relationships in Drug Design pp. 189-193 (Alan R. Liss, Inc. 1989); Lewis & Dean, 1989, Proc. R. Soc. Lond. 236:125-140 and 141-162; Askew et al., 1989, J. Am. Chem. Soc. 111:1082-1090. Other computer programs that screen and graphically depict chemicals are available from companies such as BioDesign, Inc. (Pasadena, California), Allelix, Inc. (Mississauga, Ontario, Canada), and Hypercube, Inc. (Cambridge, Ontario). Although these are primarily designed for application to drugs specific to particular proteins, they can be adapted to design of drugs specific to any identified region. The analogs and congeners can be tested for binding to the target RNA using the above-described secondary screens for biologic activity. Alternatively, lead compounds with little or no biologic activity, as ascertained in the secondary screen, can also be used to design analogs and congeners of the compound that have biologic activity.

### **5.8. Use of Identified Compounds That Modulate Untranslated Region-Dependent Gene Expression to Treat/Prevent Disease**

[0267] Biologically active compounds identified using the methods of the invention or a pharmaceutically acceptable salt thereof can be administered to a patient, preferably a mammal, more preferably a human, suffering from a disease or disorder whose onset, progression, development and/or severity is associated with the expression of a target gene.

Alternatively, biologically active compounds identified using the methods of the invention or a pharmaceutically acceptable salt thereof that are beneficial for the treatment of a disease or disorder can be administered to a patient, preferably a mammal, more preferably a human, suffering from such a disease or disorder. In a specific embodiment, a compound or a pharmaceutically acceptable salt thereof is administered to a patient, preferably a mammal, more preferably a human, as a preventative measure against a disease or disorder associated with an RNA:host cell factor interaction *in vivo*.

[0268] When administered to a patient, the compound or a pharmaceutically acceptable salt thereof is preferably administered as component of a composition that optionally comprises a pharmaceutically acceptable vehicle. The composition can be administered orally, or by any other convenient route, for example, by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (*e.g.*, oral mucosa, rectal, and intestinal mucosa, *etc.*) and may be administered together with another biologically active agent. Administration can be systemic or local. Various delivery systems are known, *e.g.*, encapsulation in liposomes, microparticles, microcapsules, capsules, *etc.*, and can be used to administer the compound and pharmaceutically acceptable salts thereof.

[0269] Methods of administration include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, oral, sublingual, intranasal, intracerebral, intravaginal, transdermal, rectally, by inhalation, or topically, particularly to the ears, nose, eyes, or skin. The mode of administration is left to the discretion of the practitioner. In most instances, administration will result in the release of the compound or a pharmaceutically acceptable salt thereof into the bloodstream.

[0270] In specific embodiments, it may be desirable to administer the compound or a pharmaceutically acceptable salt thereof locally. This may be achieved, for example, and not by way of limitation, by local infusion during surgery, topical application, *e.g.*, in conjunction with a wound dressing after surgery, by injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers.

[0271] In certain embodiments, it may be desirable to introduce the compound or a pharmaceutically acceptable salt thereof into the central nervous system by any suitable route, including intraventricular, intrathecal and epidural injection. Intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir.

[0272] Pulmonary administration can also be employed, *e.g.*, by use of an inhaler or nebulizer, and formulation with an aerosolizing agent, or via perfusion in a fluorocarbon or

synthetic pulmonary surfactant. In certain embodiments, the compound and pharmaceutically acceptable salts thereof can be formulated as a suppository, with traditional binders and vehicles such as triglycerides.

[0273] In another embodiment, the compound and pharmaceutically acceptable salts thereof can be delivered in a vesicle, in particular a liposome (see Langer, 1990, *Science* 249:1527-1533; Treat et al., 1989, in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353-365; and Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*).

[0274] In yet another embodiment, the compound and pharmaceutically acceptable salts thereof can be delivered in a controlled release system (see, e.g., Goodson, 1984, in *Medical Applications of Controlled Release*, supra, vol. 2, pp. 115-138). Other controlled-release systems discussed in the review by Langer, 1990, *Science* 249:1527-1533 may be used. In one embodiment, a pump may be used (see Langer, *supra*; Sefton, 1987, *CRC Crit. Ref. Biomed. Eng.* 14:201; Buchwald et al., 1980, *Surgery* 88:507; Saudek et al., 1989, *N. Engl. J. Med.* 321:574). In another embodiment, polymeric materials can be used (see *Medical Applications of Controlled Release*, Langer and Wise (eds.), 1974, CRC Pres., Boca Raton, Florida; *Controlled Drug Bioavailability, Drug Product Design and Performance*, Smolen and Ball (eds.), 1984, Wiley, New York; Ranger & Peppas, 1983, *J. Macromol. Sci. Rev. Macromol. Chem.* 23:61; see also Levy et al., 1985, *Science* 228:190; During et al., 1989, *Ann. Neurol.* 25:351; Howard et al., 1989, *J. Neurosurg.* 71:105). In yet another embodiment, a controlled-release system can be placed in proximity of a target RNA of the compound or a pharmaceutically acceptable salt thereof, thus requiring only a fraction of the systemic dose.

[0275] Compositions comprising the compound or a pharmaceutically acceptable salt thereof ("compound compositions") can additionally comprise a suitable amount of a pharmaceutically acceptable vehicle so as to provide the form for proper administration to the patient.

[0276] In a specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, mammals, and more particularly in humans. The term "vehicle" refers to a diluent, adjuvant, excipient, or carrier with which a compound of the invention is administered. Such pharmaceutical vehicles can be liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. The pharmaceutical vehicles can be saline, gum acacia, gelatin,

starch paste, talc, keratin, colloidal silica, urea, and the like. In addition, auxiliary, stabilizing, thickening, lubricating and coloring agents may be used. When administered to a patient, the pharmaceutically acceptable vehicles are preferably sterile. Water is a preferred vehicle when the compound of the invention is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid vehicles, particularly for injectable solutions. Suitable pharmaceutical vehicles also include excipients such as starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. Compound compositions, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents.

[0277] Compound compositions can take the form of solutions, suspensions, emulsion, tablets, pills, pellets, capsules, capsules containing liquids, powders, sustained-release formulations, suppositories, emulsions, aerosols, sprays, suspensions, or any other form suitable for use. In one embodiment, the pharmaceutically acceptable vehicle is a capsule (see *e.g.*, U.S. Patent No. 5,698,155). Other examples of suitable pharmaceutical vehicles are described in Remington's Pharmaceutical Sciences, Alfonso R. Gennaro, ed., Mack Publishing Co. Easton, PA, 19th ed., 1995, pp. 1447 to 1676, incorporated herein by reference.

[0278] In a preferred embodiment, the compound or a pharmaceutically acceptable salt thereof is formulated in accordance with routine procedures as a pharmaceutical composition adapted for oral administration to human beings. Compositions for oral delivery may be in the form of tablets, lozenges, aqueous or oily suspensions, granules, powders, emulsions, capsules, syrups, or elixirs, for example. Orally administered compositions may contain one or more agents, for example, sweetening agents such as fructose, aspartame or saccharin; flavoring agents such as peppermint, oil of wintergreen, or cherry; coloring agents; and preserving agents, to provide a pharmaceutically palatable preparation. Moreover, where in tablet or pill form, the compositions can be coated to delay disintegration and absorption in the gastrointestinal tract thereby providing a sustained action over an extended period of time. Selectively permeable membranes surrounding an osmotically active driving compound are also suitable for orally administered compositions. In these later platforms, fluid from the environment surrounding the capsule is imbibed by the driving compound, which swells to displace the agent or agent composition through an aperture. These delivery platforms can provide an essentially zero order delivery profile as opposed to the spiked profiles of immediate release

formulations. A time delay material such as glycerol monostearate or glycerol stearate may also be used. Oral compositions can include standard vehicles such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like. Such vehicles are preferably of pharmaceutical grade. Typically, compositions for intravenous administration comprise sterile isotonic aqueous buffer. Where necessary, the compositions may also include a solubilizing agent.

[0279] In another embodiment, the compound or a pharmaceutically acceptable salt thereof can be formulated for intravenous administration. Compositions for intravenous administration may optionally include a local anesthetic such as lignocaine to lessen pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water-free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the compound or a pharmaceutically acceptable salt thereof is to be administered by infusion, it can be dispensed, for example, with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the compound or a pharmaceutically acceptable salt thereof is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

[0280] The amount of a compound or a pharmaceutically acceptable salt thereof that will be effective in the treatment of a particular disease will depend on the nature of the disease, and can be determined by standard clinical techniques. In addition, *in vitro* or *in vivo* assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed will also depend on the route of administration, and the seriousness of the disease, and should be decided according to the judgment of the practitioner and each patient's circumstances. However, suitable dosage ranges for oral administration are generally about 0.001 milligram to about 500 milligrams of a compound or a pharmaceutically acceptable salt thereof per kilogram body weight per day. In specific preferred embodiments of the invention, the oral dose is about 0.01 milligram to about 500 milligrams per kilogram body weight per day, about 0.01 milligram to about 250 milligram per kilogram body weight per day, about 0.01 milligram to about 100 milligrams per kilogram body weight per day, more preferably about 0.1 milligram to about 75 milligrams per kilogram body weight per day, more preferably about 0.5 milligram to 5 milligrams per kilogram body weight per day. The dosage amounts described herein refer to total amounts administered; that is, if more than one compound is administered, or if a compound is administered with a therapeutic agent, then the preferred dosages correspond to the total

amount administered. Oral compositions preferably contain about 10% to about 95% active ingredient by weight.

[0281] Suitable dosage ranges for intravenous (i.v.) administration are about 0.01 milligram to about 100 milligrams per kilogram body weight per day, about 0.1 milligram to about 35 milligrams per kilogram body weight per day, and about 1 milligram to about 10 milligrams per kilogram body weight per day. Suitable dosage ranges for intranasal administration are generally about 0.01 pg/kg body weight per day to about 1 mg/kg body weight per day. Suppositories generally contain about 0.01 milligram to about 50 milligrams of a compound of the invention per kilogram body weight per day and comprise active ingredient in the range of about 0.5% to about 10% by weight.

[0282] Recommended dosages for intradermal, intramuscular, intraperitoneal, subcutaneous, epidural, sublingual, intracerebral, intravaginal, transdermal administration or administration by inhalation are in the range of about 0.001 milligram to about 200 milligrams per kilogram of body weight per day. Suitable doses for topical administration are in the range of about 0.001 milligram to about 1 milligram, depending on the area of administration. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model test systems. Such animal models and systems are well known in the art.

[0283] The compound and pharmaceutically acceptable salts thereof are preferably assayed *in vitro* and *in vivo*, for the desired therapeutic or prophylactic activity, prior to use in humans. For example, *in vitro* assays can be used to determine whether it is preferable to administer the compound, a pharmaceutically acceptable salt thereof, and/or another therapeutic agent. Animal model systems can be used to demonstrate safety and efficacy.

### 5.9. Target Diseases or Disorders

[0284] The present invention provides methods for preventing, treating or ameliorating a disease or disorder or one or more symptoms thereof, said methods comprising administering to a subject in need thereof one or more compounds identified in accordance with the methods of the invention. In one embodiment, the invention provides a method of preventing, treating or ameliorating a disease or disorder or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of one or more compounds identified in accordance with the methods of the invention. In another embodiment, the invention provides a method of preventing, treating or ameliorating a disease or disorder or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of one or more compounds

identified in the assays described herein, said compounds increasing untranslated region-dependent expression of a target gene whose expression useful in the prevention or treatment of said disease or disorder. In another embodiment, the invention provides a method of preventing, treating or ameliorating a disease or disorder or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of one or more compounds identified in the assays described, said compounds decreasing untranslated region-dependent expression of a target gene whose expression is associated with the onset, progression, development and/or severity of said disease or disorder. In a specific embodiment, a compound identified in accordance with the methods of the invention is not administered to prevent, treat, or ameliorate a disease or disorder or one or more symptoms thereof, if such compound has been used previously to prevent, treat or ameliorate said disease or disorder.

**[0285]** The invention also provides methods of preventing, treating or ameliorating a disease or disorder or one or more symptoms thereof, said methods comprising administering to a subject in need thereof one or more of the compounds identified utilizing the screening methods described herein, and one or more other therapies (*e.g.*, prophylactic or therapeutic agents). Preferably, such therapies (*e.g.*, prophylactic or therapeutic agents) are currently being used, have been used or are known to be useful in the prevention, treatment or amelioration of one or more symptoms associated with said disease or disorder. The therapies (*e.g.*, prophylactic or therapeutic agents) of the combination therapies of the invention can be administered sequentially or concurrently. In a specific embodiment, the combination therapies of the invention comprise a compound of the invention and at least one other therapy (*e.g.*, prophylactic or therapeutic agent) which has a different mechanism of action than said compound. The combination therapies of the present invention improve the prophylactic or therapeutic effect of a compound of the invention by functioning together with the compound to have an additive or synergistic effect. The combination therapies of the present invention reduce the side effects associated with the therapies (*e.g.*, prophylactic or therapeutic agents).

**[0286]** The prophylactic or therapeutic agents of the combination therapies can be administered to a subject in the same pharmaceutical composition. Alternatively, the prophylactic or therapeutic agents of the combination therapies can be administered concurrently to a subject in separate pharmaceutical compositions. The prophylactic or therapeutic agents may be administered to a subject by the same or different routes of administration.

[0287] In a specific embodiment, a pharmaceutical composition comprising one or more compounds identified in a screening assay described herein is administered to a subject, preferably a human, to prevent, treat or ameliorate a disease or disorder or a symptom thereof. In accordance with the invention, pharmaceutical compositions of the invention may also comprise one or more prophylactic or therapeutic agents which are currently being used, have been used or are known to be useful in the prevention, treatment or amelioration of one or more symptoms associated with a disease or disorder.

#### **5.9.1. Proliferative Disorders**

[0288] A compound identified in accordance with the methods of the invention may be administered to a subject in need thereof to prevent, treat or ameliorate a cancer or one or more symptoms thereof. A compound identified in accordance with the methods of the invention may also be administered in combination with one or more other therapies (*e.g.*, prophylactic or therapeutic agents) to a subject in need thereof to prevent, treat or ameliorate a cancer or one or more symptoms thereof. Preferably, such therapies are useful for the prevention or treatment of cancer. Examples of such therapies include, but are not limited to chemotherapeutic agents (*e.g.*, acivicin, anthramycin, bleomycin sulfate, carbetimer, carboplatin, cisplatin, cyclophosphamide, daunorubicin hydrochloride, docetaxel, doxorubicin, doxorubicin hydrochloride, epipropidine, etoposide, etoposide phosphate, etoprine fluorouracil, gemcitabine, gemcitabine hydrochloride, hydroxyurea, idarubicin hydrochloride, ifosfamide, ilmofofosine, methotrexate, methotrexate sodium, paclitaxel, trimetrexate, trimetrexate glucuronate, vinblastine sulfate, vincristine sulfate, vindesine, vindesine sulfate, and vinepidine sulfate) and anti-angiogenic agents (*e.g.*, angiostatin (plasminogen fragment), antiangiogenic antithrombin III, angiozyme, combretastatin A-4, endostatin (collagen XVIII fragment), and fibronectin fragment). In a specific embodiment, the invention provides a method of preventing, treating or ameliorating cancer or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention. In another embodiment, the invention provides a method of preventing, treating or ameliorating cancer or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention and a dose of a prophylactically or therapeutically effective amount of one or more other therapies (*e.g.*, prophylactic or therapeutic agents).



**[0289]** A compound identified in accordance with the methods of the invention may be used as a first, second, third or fourth line of therapy for the treatment of cancer. The invention provides methods for treating or ameliorating cancer or a symptom thereof in a subject refractory to conventional therapies for such a cancer, said methods comprising administering to said subject a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention. A cancer may be determined to be refractory to a therapy means when at least some significant portion of the cancer cells are not killed or their cell division arrested in response to the therapy. Such a determination can be made either *in vivo* or *in vitro* by any method known in the art for assaying the effectiveness of treatment on cancer cells, using the art-accepted meanings of “refractory” in such a context. In a specific embodiment, a cancer is refractory where the number of cancer cells has not been significantly reduced, or has increased.

**[0290]** The invention provides methods for treating or ameliorating cancer or a symptom thereof in a subject refractory to existing single agent therapies for such a cancer, said methods comprising administering to said subject a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention and a dose of a prophylactically or therapeutically effective amount of one or more other therapies (*e.g.*, prophylactic or therapeutic agents). The invention also provides methods for treating cancer by administering a compound identified in accordance with the methods of the invention in combination with any other therapy (*e.g.*, radiation therapy, chemotherapy or surgery) to patients who have proven refractory to other therapies but are no longer on these therapies. The invention also provides methods for the treatment of a patient having cancer and immunosuppressed by reason of having previously undergone other cancer therapies. The invention also provides alternative methods for the treatment of cancer where chemotherapy, radiation therapy, hormonal therapy, and/or biological therapy/immunotherapy has proven or may prove too toxic, *i.e.*, results in unacceptable or unbearable side effects, for the subject being treated. Further, the invention provides methods for preventing the recurrence of cancer in patients that have been treated and have no disease activity by administering a compound identified in accordance with the methods of the invention.

**[0291]** In this embodiment, target genes encoding proteins include, but are not limited to, angiogenin; angiopoietin1; angiopoietin2; antigen CD82; aryl hydrocarbon receptor nuclear translocator; B cell lymphoma 2; beta-catenin; cadherin-1; CLCA homolog; connective tissue growth factor, cysteine-rich 61; cyclin D1; cyclin-dependent kinase

inhibitor 2A, CDKN2 CDK4 inhibitor multiple tumor suppressor 1, SOR 1, MTS1TP16 p16(INK4) p16(INK4A) p14(ARF); cyclin-dependent kinase inhibitor 1A (p21, Cip1); dihydrofolate reductase; DNA methyltransferase; effector cell protease receptor; EMMPRIN; epithelial growth factor receptor; fibroblast growth factor 2; fibroblast growth factor 1; FMS-related tyrosine kinase 1; heparanase; hepsin; Her-2; histone acetyltransferase; histone deacetylase3; histone deacetylase 1; Hu Antigen R, a member of the Elav (embryonic lethal abnormal vision) family of RNA-binding proteins; hypoxia-inducible factor 1-alpha inhibitor; hypoxia-inducible factor 1; insulin like growth factor 1 receptor, IGF-1R; insulin-like growth factor 1; insulin-like growth factor binding protein-2; interleukin 2; interleukin-8 precursor (il-8) (monocyte-derived neutrophilchemotactic factor) (mdnsf) (T-cell chemotactic factor) (neutrophil-activating protein 1) (nap-1) (lymphocyte-derived neutrophil-activating factor) (lynaf) (protein 3-10c) (neutrophil-activating factor) (naf) (granulocyte chemotactic protein 1) (gsp-1); kit ligand, stem cell factor; large tumor suppressor; leucine amino peptidase-3; livin; major histocompatibility complex class I chain-related gene B; major histocompatibility complex class I chain-related gene A; matrix metalloproteinase 9; matrix metalloproteinase 12; max interacting protein 1 (mx1 protein); methyl-CpG-binding endonuclease; NF-Kappa-B; oncoprotein MDM2; oncoprotein fos; P-glycoprotein-1 (PGY1); placental growth factor; plasminogen activator inhibitor protein; platelet derived growth factor, beta chain; pleiotrophin; progranulin; proliferating cell nuclear antigen; protein kinase B/Akt (PBK), v-akt murine thymoma viral oncogene homolog 1, oncogene akt1 protein kinase b, pkb rac serine/threonine protein kinase; protein-tyrosine phosphatase, 4A, 3, PTP4A3; ras; retinoblastoma-binding protein 1-like 1; ribonuclease/angiogenin inhibitor; soluble-type polypeptide FZD4S; src, oncogen src protooncogene src src oncogene avian sarcoma; TEK tyrosine kinase; thrombopoietin (TPO); TIAM1: T-cell lymphoma invasion and metastasis 1; tissue inhibitor of metalloprotease 1; tissue inhibitor of metalloprotease 2; tissue inhibitor of metalloprotease 4; transforming growth factor, beta-1; tumor necrosis factor receptor superfamily, member 5, TNFRSF5; urokinase plasminogen activator; and v-myc myelocytomatosis viral oncogene homolog.

[0292] In a specific embodiment, a compound identified in the assays described herein to down-regulate untranslated region-dependent VEGF expression may be used to prevent, treat or ameliorate a vascularized tumor or one or more symptoms thereof. In another embodiment, a compound not previously known to affect VEGF expression which was identified in the assays described herein to down-regulate untranslated region-dependent

VEGF may be used to prevent, treat or ameliorate a vascularized tumor or one or more symptoms thereof.

**[0293]** In another embodiment, a compound identified in the assays described herein to down-regulate untranslated region-dependent survivin expression may be used to prevent, treat or ameliorate cancer (in particular, cancer in which survivin is highly expressed) or one or more symptoms thereof. In another embodiment, a compound not previously known to affect survivin expression which was identified in the assays described herein to down-regulate untranslated region-dependent survivin may be used to prevent, treat or ameliorate cancer or one or more symptoms thereof.

**[0294]** In another embodiment, a compound identified in the assays described herein to down-regulate untranslated region-dependent Her-2 expression in breast cancer cells may be used to prevent, treat or ameliorate breast cancer (in particular, Her-2 positive breast cancer) or one or more symptoms thereof. In another embodiment, a compound not previously known to affect Her-2 expression which was identified in the assays described herein to down-regulate untranslated region-dependent Her-2 expression may be used to prevent, treat or ameliorate breast cancers or one or more symptoms thereof.

**[0295]** Cancers that can be treated by the methods encompassed by the invention include, but are not limited to, neoplasms, tumors, metastases, or any disease or disorder characterized by uncontrolled cell growth. The cancer may be a primary or metastatic cancer. Specific examples of cancers that can be treated by the methods encompassed by the invention include, but are not limited to, cancer of the head, neck, eye, mouth, throat, esophagus, chest, bone, lung, colon, rectum, stomach, prostate, breast, ovaries, kidney, liver, pancreas, and brain. Additional cancers include, but are not limited to, the following: leukemias such as but not limited to, acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemias such as myeloblastic, promyelocytic, myelomonocytic, monocytic, erythroleukemia leukemias and myelodysplastic syndrome, chronic leukemias such as but not limited to, chronic myelocytic (granulocytic) leukemia, chronic lymphocytic leukemia, hairy cell leukemia; polycythemia vera; lymphomas such as but not limited to Hodgkin's disease, non-Hodgkin's disease; multiple myelomas such as but not limited to smoldering multiple myeloma, nonsecretory myeloma, osteosclerotic myeloma, plasma cell leukemia, solitary plasmacytoma and extramedullary plasmacytoma; Waldenström's macroglobulinemia; monoclonal gammopathy of undetermined significance; benign monoclonal gammopathy; heavy chain disease; bone and connective tissue sarcomas such as but not limited to bone sarcoma, osteosarcoma, chondrosarcoma, Ewing's sarcoma, malignant giant cell tumor, fibrosarcoma of bone, chordoma, periosteal sarcoma, soft-tissue

sarcomas, angiosarcoma (hemangiosarcoma), fibrosarcoma, Kaposi's sarcoma, leiomyosarcoma, liposarcoma, lymphangiosarcoma, neurilemmoma, rhabdomyosarcoma, synovial sarcoma; brain tumors such as but not limited to, glioma, astrocytoma, brain stem glioma, ependymoma, oligodendroglioma, nonglial tumor, acoustic neurinoma, craniopharyngioma, medulloblastoma, meningioma, pineocytoma, pineoblastoma, primary brain lymphoma; breast cancer including but not limited to adenocarcinoma, lobular (small cell) carcinoma, intraductal carcinoma, medullary breast cancer, mucinous breast cancer, tubular breast cancer, papillary breast cancer, Paget's disease, and inflammatory breast cancer; adrenal cancer such as but not limited to pheochromocytoma and adrenocortical carcinoma; thyroid cancer such as but not limited to papillary or follicular thyroid cancer, medullary thyroid cancer and anaplastic thyroid cancer; pancreatic cancer such as but not limited to, insulinoma, gastrinoma, glucagonoma, vipoma, somatostatin-secreting tumor, and carcinoid or islet cell tumor; pituitary cancers such as but limited to Cushing's disease, prolactin-secreting tumor, acromegaly, and diabetes insipidus; eye cancers such as but not limited to ocular melanoma such as iris melanoma, choroidal melanoma, and ciliary body melanoma, and retinoblastoma; vaginal cancers such as squamous cell carcinoma, adenocarcinoma, and melanoma; vulvar cancer such as squamous cell carcinoma, melanoma, adenocarcinoma, basal cell carcinoma, sarcoma, and Paget's disease; cervical cancers such as but not limited to, squamous cell carcinoma, and adenocarcinoma; uterine cancers such as but not limited to endometrial carcinoma and uterine sarcoma; ovarian cancers such as but not limited to, ovarian epithelial carcinoma, borderline tumor, germ cell tumor, and stromal tumor; esophageal cancers such as but not limited to, squamous cancer, adenocarcinoma, adenoid cystic carcinoma, mucoepidermoid carcinoma, adenosquamous carcinoma, sarcoma, melanoma, plasmacytoma, verrucous carcinoma, and oat cell (small cell) carcinoma; stomach cancers such as but not limited to, adenocarcinoma, fungating (polypoid), ulcerating, superficial spreading, diffusely spreading, malignant lymphoma, liposarcoma, fibrosarcoma, and carcinosarcoma; colon cancers; rectal cancers; liver cancers such as but not limited to hepatocellular carcinoma and hepatoblastoma, gallbladder cancers such as adenocarcinoma; cholangiocarcinomas such as but not limited to papillary, nodular, and diffuse; lung cancers such as non-small cell lung cancer, squamous cell carcinoma (epidermoid carcinoma), adenocarcinoma, large-cell carcinoma and small-cell lung cancer; testicular cancers such as but not limited to germinal tumor, seminoma, anaplastic, classic (typical), spermatocytic, nonseminoma, embryonal carcinoma, teratoma carcinoma, choriocarcinoma (yolk-sac tumor), prostate cancers such as but not limited to, adenocarcinoma, leiomyosarcoma, and rhabdomyosarcoma; penile cancers; oral cancers

such as but not limited to squamous cell carcinoma; basal cancers; salivary gland cancers such as but not limited to adenocarcinoma, mucoepidermoid carcinoma, and adenoidcystic carcinoma; pharynx cancers such as but not limited to squamous cell cancer, and verrucous; skin cancers such as but not limited to, basal cell carcinoma, squamous cell carcinoma and melanoma, superficial spreading melanoma, nodular melanoma, lentigo malignant melanoma, acral lentiginous melanoma; kidney cancers such as but not limited to renal cell cancer, adenocarcinoma, hypernephroma, fibrosarcoma, transitional cell cancer (renal pelvis and/or uterine); Wilms' tumor; bladder cancers such as but not limited to transitional cell carcinoma, squamous cell cancer, adenocarcinoma, carcinosarcoma. In addition, cancers include myxosarcoma, osteogenic sarcoma, endotheliosarcoma, lymphangioendotheliosarcoma, mesothelioma, synovioma, hemangioblastoma, epithelial carcinoma, cystadenocarcinoma, bronchogenic carcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma and papillary adenocarcinomas (for a review of such disorders, see Fishman et al., 1985, *Medicine*, 2d Ed., J.B. Lippincott Co., Philadelphia and Murphy et al., 1997, *Informed Decisions: The Complete Book of Cancer Diagnosis, Treatment, and Recovery*, Viking Penguin, Penguin Books U.S.A., Inc., United States of America). It is also contemplated that cancers caused by aberrations in apoptosis can also be treated by the methods and compositions of the invention. Such cancers may include, but not be limited to, follicular lymphomas, carcinomas with p53 mutations, hormone dependent tumors of the breast, prostate and ovary, and precancerous lesions such as familial adenomatous polyposis, and myelodysplastic syndromes.

[0296] Anti-cancer therapies and their dosages, routes of administration and recommended usage are known in the art and have been described in such literature as the *Physician's Desk Reference* (56<sup>th</sup> ed., 2002).

### **5.9.2. Inflammatory Disorders**

[0297] A compound identified in accordance with the methods of the invention may be administered to a subject in need thereof to prevent, treat or ameliorate an inflammatory disorder or one or more symptoms thereof. A compound identified in accordance with the methods of the invention may also be administered in combination with one or more other therapies (e.g., prophylactic or therapeutic agents) to a subject in need thereof to prevent, treat or ameliorate an inflammatory disorder or one or more symptoms thereof. Preferably, such therapies are useful for the prevention or treatment of an inflammatory disorder. Examples of such therapies include, but are not limited to, immunomodulatory agents (e.g., methothrexate, leflunomide, cyclophosphamide, cytoxan, Immuran, cyclosporine A, minocycline, azathioprine, and antibiotics (e.g., FK506 (tacrolimus)), anti-angiogenic

agents (*e.g.*, endostatin, angiostatin, apomigren, anti-angiogenic antithrombin III, the 29 kDa N-terminal and a 40 kDa C-terminal proteolytic fragments of fibronectin, the anti-angiogenic factor designated 13.40, the anti-angiogenic 22 amino acid peptide fragment of thrombospondin I, the anti-angiogenic 20 amino acid peptide fragment of SPARC, RGD and NGR containing peptides, the small anti-angiogenic peptides of laminin, fibronectin, procollagen and EGF, acid fibroblast growth factor (“aFGF”) antagonists, basic fibroblast growth factor (“bFGF”) antagonists, vascular endothelial growth factor (“VEGF”) antagonists, and VEGF receptor (“VEGFR”) antagonists (*e.g.*, anti-VEGFR antibodies), TNF- $\alpha$  antagonists (*e.g.*, infliximab (REMICADE™; Centacor), D2E7 (Abbott Laboratories/Knoll Pharmaceuticals Co., Mt. Olive, N.J.), CDP571 which is also known as HUMICADE™ and CDP-870 (both of Celltech/Pharmacia, Slough, U.K.), and TN3-19.12 (Williams et al., 1994, Proc. Natl. Acad. Sci. USA 91: 2762-2766; Thorbecke et al., 1992, Proc. Natl. Acad. Sci. USA 89:7375-7379), non-steroidal anti-inflammatory drugs (NSAIDs) (*e.g.*, aspirin, ibuprofen, celecoxib (CELEBREX™), diclofenac (VOLTAREN™), etodolac (LODINE™), fenoprofen (NALFON™), indomethacin (INDOCIN™), ketoralac (TORADOL™), oxaprozin (DAYPRO™), nabumentone (RELAFEN™), sulindac (CLINORIL™), tolmentin (TOLECTIN™), rofecoxib (VIOXX™), naproxen (ALEVE™, NAPROSYN™), ketoprofen (ACTRON™) and nabumetone (RELAFEN™)), steroidal anti-inflammatory drugs (*e.g.*, glucocorticoids, dexamethasone (DECADRON™), cortisone, hydrocortisone, prednisone (DELTASONE™), prednisolone, triamcinolone, azulfidine, and eicosanoids such as prostaglandins, thromboxanes, and leukotrienes), beta-agonists, anticholinergic agents, and methyl xanthines. In a specific embodiment, the invention provides a method of preventing, treating or ameliorating an inflammatory disorder or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention. In another embodiment, the invention provides a method of preventing, treating or ameliorating an inflammatory disorder or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention and a dose of a prophylactically or therapeutically effective amount of one or more other therapies (*e.g.*, prophylactic or therapeutic agents).

**[0298]** The invention provides methods for treating or ameliorating an inflammatory disorder or a symptom thereof in a subject refractory to conventional therapies (*e.g.*, methotrexate and a TNF- $\alpha$  antagonist (*e.g.*, REMICADE™ or ENBREL™)) for such an

inflammatory disorder, said methods comprising administering to said subject a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention. The invention also provides methods for treating or ameliorating an inflammatory disorder or a symptom thereof in a subject refractory to existing single agent therapies for such an inflammatory disorder, said methods comprising administering to said subject a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention and a dose of a prophylactically or therapeutically effective amount of one or more other therapies (e.g., prophylactic or therapeutic agents). The invention also provides methods for treating an inflammatory disorder by administering a compound identified in accordance with the methods of the invention in combination with any other therapy to patients who have proven refractory to other therapies but are no longer on these therapies. The invention also provides alternative methods for the treatment of an inflammatory disorder where another therapy has proven or may prove too toxic, *i.e.*, results in unacceptable or unbearable side effects, for the subject being treated. Further, the invention provides methods for preventing the recurrence of an inflammatory disorder in patients that have been treated and have no disease activity by administering a compound identified in accordance with the methods of the invention.

[0299] In this embodiment, target genes encoding proteins include, but are not limited to, a disintegrin and metallo proteinase domain 33; angiopoietin1; angiopoietin2; beta-catenin; chemokine (C-C) receptor; eotaxin; fibroblast growth factor 1; fibroblast growth factor 2; FMS-related tyrosine kinase 1; granulocyte - macrophage colony-stimulating factor precursor (GM-CSF) (colony-stimulating factor) (CSF) (sargramostim); GRO2 oncogene; macrophage inflammatory protein-2-alpha precursor (mip2-alpha) (growth regulated protein beta) (gro-beta); Hu antigen R; a member of the Elav (Embryonic lethal abnormal vision) family of RNA-binding proteins; insulin-like growth factor 1; interferon inducible protein; interferon 1 beta; interferon-alpha; interleukin 17F; interleukin 1-beta; interleukin 6; interleukin 10; interleukin 18; interleukin 13; interleukin 4; interleukin-9; leukemia Inhibitory factor Receptor; leukemia inhibitory factor; linker for Activation of T cells; macrophage migration inhibitory factor; monocyte chemotactic protein 1; NF-Kappa-B; nuclear factor of kappa light polypeptide gene enhancer in B-cells 1 (p105, NF-kappaB); osteopontin; p38 MAP Kinase; placental growth factor; platelet derived growth factor, beta chain; pleiotrophin; prolactin; receptor for interleukin-4; signal transducer and activator of transcription 6; TEK tyrosine kinase; and tumor necrosis factor alpha.

[0300] Inflammatory disorders that can be treated by the methods encompassed by the invention include, but are not limited to, asthma, encephalitis, inflammatory bowel disease, chronic obstructive pulmonary disease (COPD), allergic disorders, septic shock, pulmonary fibrosis, undifferentiated spondyloarthropathy, undifferentiated arthropathy, arthritis, inflammatory osteolysis, and chronic inflammation resulting from chronic viral or bacteria infections. Some autoimmune disorders are associated with an inflammatory condition, and thus, can be characterized as either or both an autoimmune disorder and/or an inflammatory disorder.

[0301] Anti-inflammatory therapies and their dosages, routes of administration and recommended usage are known in the art and have been described in such literature as the *Physician's Desk Reference* (56<sup>th</sup> ed., 2002).

### 5.9.3. Autoimmune Disorders

[0302] A compound identified in accordance with the methods of the invention may be administered to a subject in need thereof to prevent, treat or ameliorate an autoimmune disorder or one or more symptoms thereof. A compound identified in accordance with the methods of the invention may also be administered in combination with one or more other therapies (e.g., prophylactic or therapeutic agents) to a subject in need thereof to prevent, treat or ameliorate an autoimmune disorder or one or more symptoms thereof. Preferably, such therapies are useful for the prevention or treatment of an autoimmune disorder. In a specific embodiment, the invention provides a method of preventing, treating or ameliorating an autoimmune disorder or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention. In another embodiment, the invention provides a method of preventing, treating or ameliorating an autoimmune disorder or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention and a dose of a prophylactically or therapeutically effective amount of one or more other therapies (e.g., prophylactic or therapeutic agents).

[0303] The invention provides methods for treating or ameliorating an autoimmune disorder or a symptom thereof in a subject refractory to conventional therapies for such an autoimmune disorder, said methods comprising administering to said subject a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention. The invention also provides methods for treating or ameliorating an autoimmune disorder or a symptom thereof in a subject refractory to



existing single agent therapies for such an autoimmune disorder, said methods comprising administering to said subject a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention and a dose of a prophylactically or therapeutically effective amount of one or more other therapies (*e.g.*, prophylactic or therapeutic agents). The invention also provides methods for treating an autoimmune disorder by administering a compound identified in accordance with the methods of the invention in combination with any other therapy to patients who have proven refractory to other therapies but are no longer on these therapies. The invention also provides alternative methods for the treatment of an autoimmune disorder where another therapy has proven or may prove too toxic, *i.e.*, results in unacceptable or unbearable side effects, for the subject being treated. Further, the invention provides methods for preventing the recurrence of an autoimmune disorder in patients that have been treated and have no disease activity by administering a compound identified in accordance with the methods of the invention.

**[0304]** In this embodiment, target genes encoding proteins include, but are not limited to, adiponectin; alpha-glucosidase; forkhead box C2; G-CSF, colony stimulating factor 3 (granulocyte); galanin; gastric inhibitory polypeptide; ghrelin; glucagon receptor; glucagon-like peptide-1, GLP-1; glucokinase; glycogen synthase kinase-3B; glycogen synthase kinase-3A; human phosphotyrosyl-protein phosphatase (PTP-1B); I $\kappa$ B kinase; inositol polyphosphate phosphatase-like 1; insulin receptor; interleukin 10; leptin; neural cell adhesion molecule 1; neuron growth associated protein 43; peroxisome proliferator-activated receptor-gamma; phas; EIF4BP; protein kinase C, gamma; resistin; and uncoupling protein 2.

**[0305]** In autoimmune disorders, the immune system triggers an immune response when there are no foreign substances to fight and the body's normally protective immune system causes damage to its own tissues by mistakenly attacking self. There are many different autoimmune disorders which affect the body in different ways. For example, the brain is affected in individuals with multiple sclerosis, the gut is affected in individuals with Crohn's disease, and the synovium, bone and cartilage of various joints are affected in individuals with rheumatoid arthritis. As autoimmune disorders progress destruction of one or more types of body tissues, abnormal growth of an organ, or changes in organ function may result. The autoimmune disorder may affect only one organ or tissue type or may affect multiple organs and tissues. Organs and tissues commonly affected by autoimmune disorders include red blood cells, blood vessels, connective tissues, endocrine glands (*e.g.*, the thyroid or pancreas), muscles, joints, and skin. Examples of autoimmune disorders that

can be prevented, treated or ameliorated by the methods of the invention include, but are not limited to, alopecia areata, ankylosing spondylitis, antiphospholipid syndrome, autoimmune Addison's disease, autoimmune diseases of the adrenal gland, autoimmune hemolytic anemia, autoimmune hepatitis, autoimmune oophoritis and orchitis, autoimmune thrombocytopenia, Behcet's disease, bullous pemphigoid, cardiomyopathy, celiac sprue-dermatitis, chronic fatigue immune dysfunction syndrome (CFIDS), chronic inflammatory demyelinating polyneuropathy, Churg-Strauss syndrome, cicatrical pemphigoid, CREST syndrome, cold agglutinin disease, Crohn's disease, discoid lupus, essential mixed cryoglobulinemia, fibromyalgia-fibromyositis, glomerulonephritis, Graves' disease, Guillain-Barre, Hashimoto's thyroiditis, idiopathic pulmonary fibrosis, idiopathic thrombocytopenia purpura (ITP), IgA neuropathy, juvenile arthritis, lichen planus, lupus erythematosus, Ménière's disease, mixed connective tissue disease, multiple sclerosis, type 1 or immune-mediated diabetes mellitus, myasthenia gravis, pemphigus vulgaris, pernicious anemia, polyarteritis nodosa, polychondritis, polyglandular syndromes, polymyalgia rheumatica, polymyositis and dermatomyositis, primary agammaglobulinemia, primary biliary cirrhosis, psoriasis, psoriatic arthritis, Raynaud's phenomenon, Reiter's syndrome, Rheumatoid arthritis, sarcoidosis, scleroderma, Sjögren's syndrome, stiff-man syndrome, systemic lupus erythematosus, lupus erythematosus, takayasu arteritis, temporal arteritis/giant cell arteritis, ulcerative colitis, uveitis, vasculitides such as dermatitis herpetiformis vasculitis, vitiligo, and Wegener's granulomatosis.

[0306] Autoimmune therapies and their dosages, routes of administration and recommended usage are known in the art and have been described in such literature as the *Physician's Desk Reference* (56<sup>th</sup> ed., 2002).

#### 5.9.4. Genetic Disorders

[0307] A compound identified in accordance with the methods of the invention may be administered to a subject in need thereof to prevent, treat or ameliorate a genetic disorder or one or more symptoms thereof. A compound identified in accordance with the methods of may also be administered in combination with one or more other therapies (e.g., prophylactic or therapeutic agents) to a subject in need thereof to prevent, treat or ameliorate a genetic disorder or one or more symptoms thereof. Preferably, such therapies are useful for the prevention or treatment of a genetic disorder. In a specific embodiment, the invention provides a method of preventing, treating or ameliorating a genetic disorder or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention. In another embodiment, the

invention provides a method of preventing, treating or ameliorating a genetic disorder or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention and a dose of a prophylactically or therapeutically effective amount of one or more other therapies (*e.g.*, prophylactic or therapeutic agents).

[0308] In this embodiment, target genes encoding proteins include, but are not limited to, NAD(P)-dependent steroid dehydrogenase (EC 1.1.1.-h105e3 protein); peroxisome biogenesis factor 1 (peroxin-1) (peroxisome biogenesis disorder protein 1); and utrophin.

[0309] Examples of genetic disorders which can be prevented or treated in accordance with the invention include, but are not limited to, alopecia areata, alpha-1-antitrypsin deficiency, ataxia, Fragile X Syndrome, Gaucher disease, Hemophilia, Huntington disease, Niemann-Pick disease, Retinitis Pigmentosa, SCID (Severe Combined Immunodeficiency), Thalassemia, and Xeroderma Pigmentosum.

#### 5.9.5. Viral Infections

[0310] A compound identified in accordance with the methods of the invention may be administered to a subject in need thereof to prevent, treat or ameliorate a viral infection or one or more conditions or symptoms associated therewith. A compound identified in accordance with the methods of the invention may also be administered in combination with one or more other therapies (*e.g.*, prophylactic or therapeutic agents) to a subject in need thereof to prevent, treat or ameliorate a viral infection or one or more conditions or symptoms associated therewith. Preferably, such therapies are useful for the prevention or treatment of a viral infection. Examples of such therapies include, but are not limited to, amantadine, ribavirin, rimantadine, acyclovir, famciclovir, foscarnet, ganciclovir, trifluridine, vidarabine, didanosine, stavudine, zalcitabine, zidovudine, interferon, an antibiotic, amantadine, ribavirin, rimantadine, acyclovir, famciclovir, foscarnet, ganciclovir, trifluridine, vidarabine, didanosine, stavudine, zalcitabine, zidovudine, interferon, an antibiotic, PRO542 (Progenics) which is a CD4 fusion antibody useful for the treatment of HIV infection, Ostavir (Protein Design Labs, Inc., CA) which is a human antibody useful for the treatment of hepatitis B virus, and Protovir (Protein Design Labs, Inc., CA) which is a humanized IgG1 antibody useful for the treatment of cytomegalovirus (CMV). In a specific embodiment, the invention provides a method of preventing, treating or ameliorating a viral infection or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention.

In another embodiment, the invention provides a method of preventing, treating or ameliorating a viral infection or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention and a dose of a prophylactically or therapeutically effective amount of one or more other therapies (*e.g.*, prophylactic or therapeutic agents).

[0311] The invention provides methods for treating or ameliorating a viral infection or a symptom thereof in a subject refractory to conventional therapies for such a viral infection, said methods comprising administering to said subject a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention. The invention also provides methods for treating or ameliorating a viral infection or a symptom thereof in a subject refractory to existing single agent therapies for such a viral infection, said methods comprising administering to said subject a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention and a dose of a prophylactically or therapeutically effective amount of one or more other therapies (*e.g.*, prophylactic or therapeutic agents). The invention also provides methods for treating a viral infection by administering a compound identified in accordance with the methods of the invention in combination with any other therapy to patients who have proven refractory to other therapies but are no longer on these therapies. The invention also provides alternative methods for the treatment of a viral infection where another therapy has proven or may prove too toxic, *i.e.*, results in unacceptable or unbearable side effects, for the subject being treated. Further, the invention provides methods for preventing the recurrence of a viral infection in patients that have been treated and have no disease activity by administering a compound identified in accordance with the methods of the invention.

[0312] In this embodiment, target genes encoding proteins include, but are not limited to, C1q complement receptor-gC1qR; chemokine (C-X3-C) receptor 1; complement decay-accelerating factor [Precursor] Synonym CD55 antigen; cyclinT1; desmoglein 1; hepatitis A virus cellular receptor 1-havcr-1; hepatitis B virus X interacting protein-XIP; HIV Tat Specific Factor 1; human interferon gamma; human damage specific DNA binding protein-DDB1; INI1/hSNF5; interferon alpha-16 precursor (interferon alpha-wa); interferon alpha-5 precursor (interferon alpha-g) (leif g) (interferon alpha-61). human leukocyte (alpha) interferon; interferon alpha-1/13 precursor (interferon alpha-d) (leif d); interferon-beta 1; interleukin 8 precursor (il-8) (monocyte-derived neutrophilchemotactic factor) (mdnsf) (T-cell chemotactic factor) (Neutrophil-activating protein 1) (nap-1) (lymphocyte-derived

neutrophil-activating factor) (lynap) (protein 3-10c) (neutrophil-activating factor) (naf) (granulocyte chemotactic protein 1) (gsp-1)-activating factor) (naf) (granulocyte chemotactic protein 1) (gsp-1); interleukin 2; interleukin-12 beta chain precursor (il-12b) (cytotoxic lymphocyte maturation factor 40 kda subunit) (clmf p40) (nk cell stimulatory factor chain 2) (nksf2); natural resistance-associated macrophage protein 1 (nramp 1); p300/CBP associated factor (PCAF); poly(rC) binding protein 2; and virion infectivity factor.

[0313] Any type of viral infection can be prevented, treated or ameliorated in accordance with the methods of invention. Examples of viruses which cause viral infections include, but not limited to, retroviruses (*e.g.*, human T-cell lymphotropic virus (HTLV) types I and II and human immunodeficiency virus (HIV)), herpes viruses (*e.g.*, herpes simplex virus (HSV) types I and II, Epstein-Barr virus, HHV6-HHV8, and cytomegalovirus), arenaviruses (*e.g.*, lassa fever virus), paramyxoviruses (*e.g.*, morbillivirus virus, human respiratory syncytial virus, mumps, and pneumovirus), adenoviruses, bunyaviruses (*e.g.*, hantavirus), coronaviruses, filoviruses (*e.g.*, Ebola virus), flaviviruses (*e.g.*, hepatitis C virus (HCV), yellow fever virus, and Japanese encephalitis virus), hepadnaviruses (*e.g.*, hepatitis B viruses (HBV)), orthomyxoviruses (*e.g.*, influenza viruses A, B and C), papovaviruses (*e.g.*, papillomaviruses), picornaviruses (*e.g.*, rhinoviruses, enteroviruses and hepatitis A viruses), poxviruses, reoviruses (*e.g.*, rotaviruses), togaviruses (*e.g.*, rubella virus), rhabdoviruses (*e.g.*, rabies virus).

[0314] Viral infection therapies and their dosages, routes of administration and recommended usage are known in the art and have been described in such literature as the *Physician's Desk Reference* (56<sup>th</sup> ed., 2002).

#### 5.9.6. Fungal Infections

[0315] A compound identified in accordance with the methods of the invention may be administered to a subject in need thereof to prevent, treat or ameliorate a fungal infection or one or more conditions or symptoms associated therewith. A compound identified in accordance with the methods of the invention may also be administered in combination with one or more other therapies (*e.g.*, prophylactic or therapeutic agents) to a subject in need thereof to prevent, treat or ameliorate a fungal infection or one or more conditions or symptoms associated therewith. Preferably, such therapies are useful for the prevention or treatment of a fungal infection. Examples of such therapies include, but are not limited to, amphotericin B or analogs or derivatives thereof (including 14(s)-hydroxyamphotericin B methyl ester, the hydrazide of amphotericin B with 1-amino-4-methylpiperazine, and other derivatives) or other polyene macrolide antibiotics (including, *e.g.*, nystatin, candicidin,

pinaricidin and natamycin), flucytosine; flucytosine; griseofulvin; echinocandins or aureobasidins, including naturally occurring and semi-synthetic analogs; dihydrobenzo[a]naphthacenequinones; nucleoside peptide antifungals including the polyoxins and nikkomycins; allylamines such as naftifine and other squalene epoxidase inhibitors; azoles, imidazoles and triazoles such as, *e.g.*, clotrimazole, miconazole, ketoconazole, econazole, butoconazole, oxiconazole, terconazole, itraconazole or fluconazole and the like; rapamycin and rapalogs (non-immunosuppressive derivatives of rapamycin); cyclosporin A; FK506; terbinafine; and natural compounds found in goldenseal root powder, ipe roxo powder, poke root powder, lavender oil, and tea tree oil. In a specific embodiment, the invention provides a method of preventing, treating or ameliorating a fungal infection or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention. In another embodiment, the invention provides a method of preventing, treating or ameliorating a fungal infection or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention and a dose of a prophylactically or therapeutically effective amount of one or more other therapies (*e.g.*, prophylactic or therapeutic agents).

**[0316]** The invention provides methods for treating or ameliorating a fungal infection or a symptom thereof in a subject refractory to conventional therapies for such a fungal infection, said methods comprising administering to said subject a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention. The invention also provides methods for treating or ameliorating a fungal infection or a symptom thereof in a subject refractory to existing single agent therapies for such a fungal infection, said methods comprising administering to said subject a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention and a dose of a prophylactically or therapeutically effective amount of one or more therapies (*e.g.*, prophylactic or therapeutic agents). The invention also provides methods for treating a fungal infection by administering a compound identified in accordance with the methods of the invention in combination with any other therapy to patients who have proven refractory to other therapies but are no longer on these therapies. The invention also provides alternative methods for the treatment of a fungal infection where another therapy has proven or may prove too toxic, *i.e.*, results in unacceptable or unbearable side effects, for the subject being

treated. Further, the invention provides methods for preventing the recurrence of a fungal infection in patients that have been treated and have no disease activity by administering a compound identified in accordance with the methods of the invention.

[0317] In this embodiment, target genes encoding proteins include, but are not limited to, complement decay-accelerating factor [precursor] synonym CD55 antigen; desmoglein 1; human interferon gamma; interferon alpha-16 precursor (interferon alpha-wa); interferon alpha-1/13 precursor (interferon alpha-d) (leif d); interferon alpha-5 precursor (interferon alpha-g) (leif g) (interferon alpha-61). human leukocyte (alpha) interferon; interferon-beta 1; interleukin 2; interleukin-12 beta chain precursor (il-12b) (cytotoxic lymphocyte maturation factor 40 kda subunit) (clmf p40) (nk cell stimulatory factor chain 2) (nksf2); interleukin-8 precursor (il-8) (monocyte-derived neutrophilchemotactic factor) (mdnsf) (T-cell chemotactic factor) (neutrophil-activating protein 1) (nap-1) (lymphocyte-derived neutrophil-activating factor) (lynap) (protein 3-10c) (neutrophil-activating factor) (naf) (granulocyte chemotactic protein 1) (gsp-1); and natural resistance-associated macrophage protein 1 (nramp 1).

[0318] Any type of fungal infection can be prevented, treated or ameliorated in accordance with the methods of invention. Examples of fungi which cause fungal infections include, but not limited to, *Absidia* species (e.g., *Absidia corymbifera* and *Absidia ramosa*), *Aspergillus* species, (e.g., *Aspergillus flavus*, *Aspergillus fumigatus*, *Aspergillus nidulans*, *Aspergillus niger*, and *Aspergillus terreus*), *Basidiobolus ranarum*, *Blastomyces dermatitidis*, *Candida* species (e.g., *Candida albicans*, *Candida glabrata*, *Candida kerr*, *Candida krusei*, *Candida parapsilosis*, *Candida pseudotropicalis*, *Candida quilliermondii*, *Candida rugosa*, *Candida stellatoidea*, and *Candida tropicalis*), *Coccidioides immitis*, *Conidiobolus* species, *Cryptococcus neoforms*, *Cunninghamella* species, dermatophytes, *Histoplasma capsulatum*, *Microsporium gypseum*, *Mucor pusillus*, *Paracoccidioides brasiliensis*, *Pseudallescheria boydii*, *Rhinosporidium seeberi*, *Pneumocystis carinii*, *Rhizopus* species (e.g., *Rhizopus arrhizus*, *Rhizopus oryzae*, and *Rhizopus microsporus*), *Saccharomyces* species, *Sporothrix schenckii*, zygomycetes, and classes such as *Zygomycetes*, *Ascomycetes*, the *Basidiomycetes*, *Deuteromycetes*, and *Oomycetes*.

[0319] Fungal infection therapies and their dosages, routes of administration and recommended usage are known in the art and have been described in such literature as the *Physician's Desk Reference* (56<sup>th</sup> ed., 2002).

#### 5.9.7. Bacterial Infections

[0320] A compound identified in accordance with the methods of the invention may be administered to a subject in need thereof to prevent, treat or ameliorate a bacterial infection or one or more conditions or symptoms thereof. A compound identified in accordance with the methods of the invention may also be administered in combination with one or more other therapies (*e.g.*, prophylactic or therapeutic agents) to a subject in need thereof to prevent, treat or ameliorate a bacterial infection or one or more conditions or symptoms thereof. Preferably, such therapies are useful for the prevention or treatment of a bacterial infection. Examples of such therapies include, but are not limited to, amoxycillin, bacteriophages, chloramphenicol, chlorhexidine, co-trimoxazole, fluoroquinolones (*e.g.*, ciprofloxacin and ofloxacin), isoniazid, macrolides, oxazolidinones, penicillin, quinolones, rifampicin, rifamycins, streptomycin, sulfonamides, and tetracyclines. In a specific embodiment, the invention provides a method of preventing, treating or ameliorating a bacterial infection or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention. In another embodiment, the invention provides a method of preventing, treating or ameliorating a bacterial infection or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention and a dose of a prophylactically or therapeutically effective amount of one or more other therapies (*e.g.*, prophylactic or therapeutic agents).

[0321] The invention provides methods for treating or ameliorating a bacterial infection or a symptom thereof in a subject refractory to conventional therapies for such a bacterial infection, said methods comprising administering to said subject a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention. The invention also provides methods for treating or ameliorating a bacterial infection or a symptom thereof in a subject refractory to existing single agent therapies for such a bacterial infection, said methods comprising administering to said subject a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention and a dose of a prophylactically or therapeutically effective amount of one or more other therapies (*e.g.*, prophylactic or therapeutic agents). The invention also provides methods for treating a bacterial infection by administering a compound identified in accordance with the methods of the invention in combination with any other therapy to patients who have proven refractory to other therapies but are no longer on these therapies. The invention also



provides alternative methods for the treatment of a bacterial infection where another therapy has proven or may prove too toxic, *i.e.*, results in unacceptable or unbearable side effects, for the subject being treated. Further, the invention provides methods for preventing the recurrence of a bacterial infection in patients that have been treated and have no disease activity by administering a compound identified in accordance with the methods of the invention.

[0322] In this embodiment, target genes encoding proteins include, but are not limited to, ADP-ribosylation factor-4; bactericidal/permeability-increasing protein; complement decay-accelerating factor [precursor] synonym CD55 antigen; desmoglein 1; human interferon gamma; interferon alpha-16 precursor (interferon alpha-wa); interferon alpha-1/13 precursor (interferon alpha-d) (leif d); interferon alpha-5 precursor (interferon alpha-g) (leif g) (interferon alpha-61). human leukocyte (alpha) interferon; interferon-beta 1; interleukin 2; interleukin-12 beta chain precursor (il-12b) (cytotoxic lymphocyte maturation factor 40 kda subunit) (clmf p40) (nk cell stimulatory factor chain 2) (nksf2); interleukin-8 precursor (il-8) (monocyte-derived neutrophil chemotactic factor) (mdnsf) (T-cell chemotactic factor) (neutrophil-activating protein 1) (nap-1) (lymphocyte-derived neutrophil-activating factor) (lynaf) (protein 3-10c) (neutrophil-activating factor) (naf) (granulocyte chemotactic protein 1) (gsp-1); and natural resistance-associated macrophage protein 1 (nramp 1).

[0323] Any type of bacterial infection can be prevented, treated or ameliorated in accordance with the methods of invention. Examples of bacteria which cause bacterial infections include, but not limited to, the *Aquaspirillum* family, *Azospirillum* family, *Azotobacteraceae* family, *Bacteroidaceae* family, *Bartonella* species, *Bdellovibrio* family, *Campylobacter* species, *Chlamydia* species (*e.g.*, *Chlamydia pneumoniae*), clostridium, *Enterobacteriaceae* family (*e.g.*, *Citrobacter* species, *Edwardsiella*, *Enterobacter aerogenes*, *Erwinia* species, *Escherichia coli*, *Hafnia* species, *Klebsiella* species, *Morganella* species, *Proteus vulgaris*, *Providencia*, *Salmonella* species, *Serratia marcescens*, and *Shigella flexneri*), *Gardinerella* family, *Haemophilus influenzae*, *Halobacteriaceae* family, *Helicobacter* family, *Legionellaceae* family, *Listeria* species, *Methylococcaceae* family, mycobacteria (*e.g.*, *Mycobacterium tuberculosis*), *Neisseriaceae* family, *Oceanospirillum* family, *Pasteurellaceae* family, *Pneumococcus* species, *Pseudomonas* species, *Rhizobiaceae* family, *Spirillum* family, *Spirosomaceae* family, *Staphylococcus* (*e.g.*, methicillin resistant *Staphylococcus aureus* and *Staphylococcus pyrogenes*), *Streptococcus* (*e.g.*, *Streptococcus enteritidis*, *Streptococcus fasciae*, and *Streptococcus pneumoniae*) *Vampirovibr* *Helicobacter* family, and *Vampirovibrio* family.

[0324] Bacterial infection therapies and their dosages, routes of administration and recommended usage are known in the art and have been described in such literature as the *Physician's Desk Reference* (56<sup>th</sup> ed., 2002).

#### 5.9.8. Cardiovascular Diseases

[0325] A compound identified in accordance with the methods of the invention may be administered to a subject in need thereof to prevent, treat or ameliorate a cardiovascular disease or one or more symptoms thereof. A compound identified in accordance with the methods of the invention may also be administered in combination with one or more other therapies (*e.g.*, prophylactic or therapeutic agents) to a subject in need thereof to prevent, treat or ameliorate a cardiovascular disease or one or more symptoms thereof. Preferably, such therapies are useful for the prevention or treatment of a cardiovascular disease. Examples of such therapies include, but are not limited to, peripheral antiadrenergic drugs, centrally acting antihypertensive drugs (*e.g.*, methyldopa and methyldopa HCl), antihypertensive direct vasodilators (*e.g.*, diazoxide and hydralazine HCl), drugs affecting renin-angiotensin system, peripheral vasodilators, phentolamine, antianginal drugs, cardiac glycosides, inodilators (*e.g.*, amrinone, milrinone, enoximone, fenoximone, imazodan, and sulmazole), antidysrhythmic drugs, calcium entry blockers, ranitine, bosentan, and rezulin. In a specific embodiment, the invention provides a method of preventing, treating or ameliorating a cardiovascular disease or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention. In another embodiment, the invention provides a method of preventing, treating or ameliorating a cardiovascular disease or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention and a dose of a prophylactically or therapeutically effective amount of one or more other therapies (*e.g.*, prophylactic or therapeutic agents).

[0326] The invention provides methods for treating or ameliorating one or more symptoms of a cardiovascular disease in a subject refractory to conventional therapies for such a cardiovascular disease, said methods comprising administering to said subject a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention. The invention also provides methods for treating or ameliorating one or more symptoms of a cardiovascular disease in a subject refractory to existing single agent therapies for such a cardiovascular disease, said methods comprising administering to said subject a dose of a prophylactically or therapeutically

effective amount of a compound identified in accordance with the methods of the invention and a dose of a prophylactically or therapeutically effective amount of one or more other therapies (e.g., prophylactic or therapeutic agents). The invention also provides methods for treating a cardiovascular disease by administering a compound identified in accordance with the methods of the invention in combination with any other therapy to patients who have proven refractory to other therapies but are no longer on these therapies. The invention also provides alternative methods for the treatment of a cardiovascular disease where another therapy has proven or may prove too toxic, *i.e.*, results in unacceptable or unbearable side effects, for the subject being treated. Further, the invention provides methods for preventing the recurrence of a cardiovascular disease in patients that have been treated and have no disease activity by administering a compound identified in accordance with the methods of the invention.

[0327] In this embodiment, target genes encoding proteins include, but are not limited to, 3-hydroxy-3-methylglutaryl-CoA reductase; actin, alpha cardiac; acyl-coa dehydrogenase; angiotensin I-converting enzyme; bile salt export pump (atp-binding cassette, sub-family b, member 11); cardiac muscle troponin T; carnitine o-palmitoyltransferase; emotakin ATP-binding cassette, sub-family a, member 1 (atp-binding cassette transporter 1) (atp-binding cassette 1) (abc-1) (cholesterol efflux regulatory protein); erythropoietin; fibrillin; human trisosephosphate isomerase; iduronate 2-sulfatase; klotho; and thrombomodulin.

[0328] Any cardiovascular disease can be prevented, treated or ameliorated in accordance with the methods of the invention. Examples of cardiovascular diseases include, but not limited to, atherosclerosis, stroke, cerebral infarction, endothelium dysfunctions (in particular, those dysfunctions affecting blood vessel elasticity) ischemic heart disease (e.g., angina pectoris, myocardial infarction, and chronic ischemic heart disease), hypertensive heart disease, pulmonary heart disease, coronary heart disease, valvular heart disease (e.g., rheumatic fever and rheumatic heart disease, endocarditis, mitral valve prolapse, restenosis and aortic valve stenosis), congenital heart disease (e.g., valvular and vascular obstructive lesions, atrial or ventricular septal defect, and patent ductus arteriosus), and myocardial disease (e.g., myocarditis, congestive cardiomyopathy, and hypertrophic cardiomyopathy).

[0329] Cardiovascular disease therapies and their dosages, routes of administration and recommended usage are known in the art and have been described in such literature as the *Physician's Desk Reference* (56<sup>th</sup> ed., 2002).

#### 5.9.9. Central Nervous System Disorders

**[0330]** A compound identified in accordance with the methods of the invention may be administered to a subject in need thereof to prevent, treat or ameliorate a central nervous system ("CNS") disorder or one or more symptoms thereof. A compound identified in accordance with the methods of the invention may also be administered in combination with one or more other therapies (*e.g.*, prophylactic or therapeutic agents) to a subject in need thereof to prevent, treat or ameliorate a CNS disorder or one or more symptoms thereof. Preferably, such therapies are useful for the prevention or treatment of a CNS disorder. Examples of such therapies include, but are not limited to, levodopa, Parlodel (bromocriptine), Permax (pergolide), Eldepryl (selegiline hydrochloride), donepezil (Aricept®), tacrine (Cognex®), acyclovir, antibiotics, chemotherapeutics, and radiation therapy. In a specific embodiment, the invention provides a method of preventing, treating or ameliorating a CNS disorder or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention. In another embodiment, the invention provides a method of preventing, treating or ameliorating a CNS disorder or one or more symptoms thereof, said method comprising administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention and a dose of a prophylactically or therapeutically effective amount of one or more other therapies (*e.g.*, prophylactic or therapeutic agents).

**[0331]** The invention provides methods for treating or ameliorating one or more symptoms of a CNS disorder in a subject refractory to conventional therapies for such a cardiovascular disease, said methods comprising administering to said subject a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention. The invention also provides methods for treating or ameliorating one or more symptoms of a CNS disorder in a subject refractory to existing single agent therapies for such a CNS disorder, said methods comprising administering to said subject a dose of a prophylactically or therapeutically effective amount of a compound identified in accordance with the methods of the invention and a dose of a prophylactically or therapeutically effective amount of one or more other therapies (*e.g.*, prophylactic or therapeutic agents). The invention also provides methods for treating a CNS disorder by administering a compound identified in accordance with the methods of the invention in combination with any other therapy to patients who have proven refractory to other therapies but are no longer on these therapies. The invention also provides alternative methods for the treatment of a CNS disorder where another therapy has proven or may

prove too toxic, *i.e.*, results in unacceptable or unbearable side effects, for the subject being treated. Further, the invention provides methods for preventing the recurrence of a cardiovascular disease in patients that have been treated and have no disease activity by administering a compound identified in accordance with the methods of the invention.

[0332] In this embodiment, target genes encoding proteins include, but are not limited to, acetylcholinesterase; Alzheimer's disease amyloid A4 [precursor], synonyms: protease nexin-II PN-II APPI; beta-site APP-cleaving enzyme 2; catechol-O-methyltransferase; CREAM/calsenilin/KCh IP3; D-amino-acid oxidase; drebrin-1 dendritic spine protein; glutamic acid decarboxylase 2; glutamic acid decarboxylase, brain, membrane form; glutamic acid decarboxylase 3; human D-1 dopamine receptor; huntingtin; kallikrein 6; monoamine oxidase-A; monoamine oxidase-B; N-methylD-aspartate (NMDA) receptor; peroxisome assembly factor-2 (paf-2) (peroxisomal-type atpase 1) (peroxin-6); and vanilloid receptor subunit 1 (capsaicin receptor).

[0333] Any CNS disorder can be treated in accordance with the methods of the invention. Examples of CNS disorders include, but are not limited to, bacterial and viral meningitis, Alzheimers Disease, cerebral toxoplasmosis, Parkinson's disease, multiple sclerosis, brain cancers (*e.g.*, metastatic carcinoma of the brain, glioblastoma, astrocytoma, and acoustic neuroma), hydrocephalus, and encephalitis.

[0334] CNS disorder therapies and their dosages, routes of administration and recommended usage are known in the art and have been described in such literature as the *Physician's Desk Reference* (56<sup>th</sup> ed., 2002).

#### **5.10. Classification of UTRs, Compound, and Disease State**

[0335] The names of the compounds identified in accordance with the methods described herein and the names of the genes whose expression are modulated in response to said compounds can be maintained in a database. By performing an assay for modulators of untranslated region-dependent gene expression using the same assay format for a group of untranslated regions of target genes (*i.e.*, an assay format in which the only variable between each individual assay is the nucleotide sequence of untranslated regions operably linked to a reporter gene) and by storing all relevant data in a database, cluster analysis can be performed on the data and functional associations between and/or within relevant data sets (*e.g.*, (a) compounds, (b) nucleotide sequences of untranslated regions of genes, and (c) pathological conditions associated with genes) can be determined.

[0336] For example, if a common set of compounds modulates the expression of a reporter gene when the latter is operably linked to a set of untranslated regions from all untranslated regions analyzed, it can be concluded that the untranslated regions in the set

are involved in a common process or processes in post-transcriptional control of gene expression. If functional involvement in post-transcriptional control of gene expression has been reported for any of the untranslated regions in the set, the untranslated regions without known roles in gene expression regulation can be assigned a function. By performing further analysis and looking for sets of compounds that modulate sets of untranslated regions common to a particular pathological condition, the following can be identified: (a) members of biochemical reaction pathways involved in the disease, (b) targets for multiple drug intervention and/or regulation, and (c) multiple pathological conditions that can be treated with a single compound or set of compounds.

## 6. **EXAMPLE: THERAPEUTIC UNTRANSLATED REGION TARGETS**

[0337] The therapeutic targets presented herein are by way of example, and the present invention is not to be limited by the targets described herein. The therapeutic targets presented herein as DNA sequences are understood by one of skill in the art that the sequences can be converted to RNA sequences.

### 6.1. **Tumor Necrosis Factor Alpha**

[0338] See, *e.g.*, GenBank Accession # X01394.

General Target Regions:

[0339] (1) 5' Untranslated Region - nts 1 - 152 of GenBank Accession # X01394:

gcagaggacc agctaagagg gagagaagca actacagacc cccctgaaa acaaccctca gacgccacat  
ccctgacaa gctgccaggc aggttctctt cctctcatat actgaccac ggctccacc tctctccct ggaaaggaca cc  
(SEQ ID NO: 5)

[0340] (2) 3' Untranslated Region - nts 852 - 1643 of GenBank Accession # X01394:

tgaggagga cgaacatcca acctcccaa acgcctccc tgcccaatc ctttattac cccctcctc agacaccctc  
aacctctctt ggctcaaaaa gagaattggg ggcttagggc cggaacccaa gcttagaact ttaagcaaca agaccaccac  
ttcgaaacct gggattcagg aatgtgtggc ctgcacagt aattgctggc aaccactaag aattcaaact ggggcctcca  
gaactcactg gggcctacag ctttgatccc tgacatctgg aatctggaga ccaggagacc ttggttctg gccagaatgc  
tgcaggactt gagaagacct cacctagaaa ttgacacaag tggaccttag gccttcctct ctccagatgt ttccagactt  
ccttgagaca cggagcccag cctcccccatt ggagccagct cctctattt atgtttgcac ttgtgattat ttattattt  
attattttac agatgaatgt atttattgg gagaccgggg tctctgggg gacccaatgt aggagctgcc ttggtcaga  
catgtttcc gtgaaaacgg agctgaacaa taggctgttc ccatgtagcc cctggcctc tgtgccttct ttgattatg  
tttttaaaa tatttatctg attaatgtgt ctaaacaatg ctgatttgg gaccaactgt cactcattgc tgaacctctg  
ctccccaggg gagtgtgtc tgtaatgcc ctactattca gtggcgagaa ataaagttg ctt (SEQ ID NO: 6)

Initial Specific Target Motif:

[0341] (3) Group I AU-Rich Element (ARE) Cluster in 3' untranslated region:  
5' AUUUAUUUAUUUAUUUAUUUA 3' (SEQ ID NO: 7)

## 6.2. Granulocyte-macrophage Colony Stimulating Factor

[0342] See, *e.g.*, GenBank Accession # NM\_000758 or # XM\_003751.

General Target Regions:

[0343] (1) 5' Untranslated Region - nts 1 - 32 of GenBank Accession #  
NM\_000758:

g/tctggaggat gtggctgcag agcctgctgc tcttgggcac (SEQ ID NO: 8)

[0344] (2) 3' Untranslated Region - nts 468 - 789 of GenBank Accession #  
NM\_000758:

gcc ggggagctgc tctctcatga aacaagagct agaaactcag gatggctcgc ttggaggagc caaggggtgg  
gccacagcca tgggtgggagt ggccctggacc tgccctgggc cacactgacc ctgatacagg catggcagaa  
gaatgggaat atttatact gacagaaatc agtaatatat atatattat attttaaaa tatttattta tttatttatt taagttcata  
ttccatattt attcaagatg tttaccgta ataattatta ttaaaaatat gcttct (SEQ ID NO: 9)

[0345] Initial Specific Target Motif:

[0346] Group I AU-Rich Element (ARE) Cluster in 3' untranslated region:  
5' AUUUAUUUAUUUAUUUAUUUA 3' (SEQ ID NO: 10)

## 6.3. Interleukin 2

[0347] See, *e.g.*, GenBank Accession # U25676.

General Target Regions:

[0348] (1) 5' Untranslated Region - nts 1 - 47 of GenBank Accession # U25676:  
atcactctct ttaatcacta ctcacattaa cctcaactcc tgccaca (SEQ ID NO: 11)

[0349] (2) 3' Untranslated Region - nts 519- 825 of GenBank Accession # U25676:  
ta attaatgtct tccacttaa aacatatcag gccttctatt tatttattta aatatttaaa tttatatatt attgttgaat gtatgggtgc  
tacctattgt aactattatt cttaatttta aaactataaa tatggatctt ttatgattct ttttgaagc cctaggggct ctaaaatggt  
ttaccttatt tatcccaaaa atatttatta ttatgttgaa tgttaaatat agtatctatg tagattgggt agtaaaacta ttaataaat  
ttgataaata taaaaaaaaa aaacaaaaaa aaaaa (SEQ ID NO: 12)

[0350] Initial Specific Target Motifs:

Group III AU-Rich Element (ARE) Cluster in 3' untranslated region:  
5' NAUUUAUUUAUUUAN 3' (SEQ ID NO: 13)

## 6.4. Interleukin 6

[0351] See, *e.g.*, GenBank Accession # NM\_000600.

General Target Regions:

[0352] (1) 5' Untranslated Region - nts 1 - 62 of GenBank Accession #  
NM\_000600:

ttctgccctc gagcccaccg ggaacgaaag agaagctcta tctgcctcc aggagcccag ct (SEQ ID NO: 14)

[0353] (2) 3' Untranslated Region - nts 699 - 1125 of GenBank Accession #  
NM\_000600:

ta gcatgggcac ctcagattgt tgttgtaat gggcattcct tctctggtc agaaacctgt ccactgggca cagaacttat  
gttgctctct atggagaact aaaagtatga gcgtaggac actattttaa ttatttttaa ttattaata tttaaatatg tgaagctgag  
ttaatttatg taagtcatat ttatattttt aagaagtacc acttgaaaca tttatgtat tagttttgaa ataataatgg aaagtggcta  
tgcagtttga atatcctttg ttccagagcc agatcatttc ttggaaagt taggcttacc tcaataaat ggctaactta  
tacatatttt taaagaaata ttatattgt atttatataa tgtataaat gttttatac caataaatgg cattttaaaa aattc (SEQ  
ID NO: 15)

[0354] Initial Specific Target Motifs:

Group III AU-Rich Element (ARE) Cluster in 3' untranslated region:

5' NAUUUAUUUAUUUAN 3' (SEQ ID NO: 16)

### 6.5. Vascular Endothelial Growth Factor

[0355] See, *e.g.*, GenBank Accession # AF022375.

General Target Regions:

[0356] (1) 5' Untranslated Region - nts 1 - 701 of GenBank Accession # AF022375:  
aagagctcca gagagaagtc gaggaagaga gagacggggt cagagagagc gcgcgggcgt gcgagcagcg  
aaagcgacag gggcaaagt agtgacctgc tttgggggt gaccgccgga gcgcggcgtg agccctcccc  
cttgggatcc cgcagctgac cagtcgcgt gacggacaga cagacagaca ccgccccag cccagttac cacctctcc  
ccggccggcg gcggacagt gacgcggcgg cgagccgcgg gcaggggccc gagcccgccc ccggaggcgg  
gggtggagggg gtcggagctc gcggcgtcgc actgaaactt ttcgtccaac ttctgggctg ttctgcctc ggaggagccg  
tggtccgcgc gggggaagcc gagccgagcg gagccgcgag aagtgttagc tcgggccggg aggagccgca  
gccggaggag ggggaggagg aagaagagaa ggaagaggag agggggccgc agtggcgact cggcgctcgg  
aagccgggct catggacggg tgaggcggcg gtgtgcgcag acagtgtctc agcgcgcgcg ctccccagcc  
ctggccccgc ctcgggccgg gaggaagagt agctgccga ggcgccgagg agagcgggcc gccccacagc  
ccgagccgga gagggacgcg agccgcgcgc cccggtcggg cctccgaaac c (SEQ ID NO: 17)

[0357] (2) 3' Untranslated Region - nts 1275 - 3166 of GenBank Accession #  
AF022375:

tgagcc gggcaggagg aaggagcctc ctcagggtt tcgggaacca gatctctc caggaaagac tgatacagaa  
cgatcgatac agaaaccacg ctgccgccac cacaccatca ccacgacag aacagtcctt aatccagaaa cctgaaatga  
aggaagagga gactctgcgc agagcacttt gggtcgggag ggcgagactc cggcggaagc attcccgggc  
gggtgacca gcacgggtcc tcttgaatt ggattcgcca tttattttt ctgtctgta aatcaccgag cccggaagat  
tagagagttt tatttctggg attcctgtag acacaccac ccacatacat acatttatat atatatata tatatatata



taaaaataaa tatctctatt ttatatata aaaatatata tattcttttt ttaaattaac agtgctaatag ttattggtgt cttcactgga  
 tgtatttgac tgctgtggac ttgagttggg aggggaatgt tcccactcag atcctgacag ggaagaggag gagatgagag  
 actctggcat gatcttttt ttgtcccact tgggtggggcc agggctctct cccctgcccagaatgtgca aggccagggc  
 atggggggcaa atatgaccca gttttgggaa caccgacaaa cccagccctg gcgctgagcc tctctacccc aggtcagacg  
 gacagaaaga caaatcacag gttccgggat gaggacaccg gctctgacca ggagtttggg gagcttcagg acattgctgt  
 gctttgggga ttccctccac atgctgcacg cgcactcgc cccagggggc actgcctgga agattcagga gcctggggcg  
 ccttcgctta ctctcacctg cttctgagtt gccaggagg ccaactggcag atgtcccggc gaagagaaga gacacattgt  
 tggaagaagc agcccatgac agcgccccct cctgggactc gccctcatcc tcttctgct ccccttctg ggggtgcagcc  
 taaaaggacc tatgtctca caccattgaa accactagtt ctgtccccc aggaaacctg gttgtgtgtg tgtgagtggt  
 tgaccttct ccatccctg gtccctccct tccctcccg aggcacagag agacagggca ggatccacgt gccattgtg  
 gaggcagaga aaagagaaag tgtttatat acggtactta ttaatatcc cttttaatt agaaattaga acagttaatt  
 taattaaaga gtagggtttt ttctcagttat tcttggttaa tatttaattt caactattta tgagatgtat ctttgctct ctctgctct  
 cttatttga ccggttttg tatataaat tcatgttcc aatctctctc tcctgatcg gtgacagtca ctagcttate ttgaacagat  
 atttaatttt gctaacactc agctctgccc tccccgatcc cctggctccc cagcacacat tccttgaaa gagggtttca  
 atatacatct acatactata tatatatgg gcaacttgta ttgtgtgta tatatatata tatatgttta tgtatatatg tgatcctgaa  
 aaaataaaca tcgctattct gtttttata tgttcaaacc aaacaagaaa aaatagagaa ttctacatac taaatctctc  
 tcctttttta attttaatat ttgttatcat ttattattg gtgctactgt ttatccgtaa taattgtggg gaaaagatat taacatcacg  
 tctttgtctc tagtgcagtt ttccgagata ttccgtagta catatttatt tttaacaac gacaaagaaa tacagatata  
 tcttaaaaaa aaaaaa (SEQ ID NO: 18)

[0358] Initial Specific Target Motifs:

[0359] (1) Internal Ribosome Entry Site (IRES) in 5' untranslated region nts 513 - 704:

5'CCGGGCUCAUGGACGGGUGAGGCGGCGGUGUGCGCAGACAGUGCUCCAGCG  
 CGCGCGCUCCCCAGCCUUGGCCCGGCCUUGGGCCGGGAGGAAGAGUAGCUCG  
 CCGAGGCGCCGAGGAGAGCGGGCCGCCACAGCCCGAGCCGGAGAGGGACG  
 CGAGCCGCGCGCCCCGGUCGGGCCUCCGAAACCAUGAACUUUCUGCUGUCUU  
 GGGUGCAUUGGAGCCUUGCCUUGCUGCUCUACCUCCACCAUG 3' (SEQ ID NO:  
 19)

[0360] (2) Group III AU-Rich Element (ARE) Cluster in 3' untranslated region:  
 5' NAUUUAUUUAUUUAN 3' (SEQ ID NO: 20)

## 6.6. Survivin

[0361] See, *e.g.*, GenBank Accession # NM\_001168.

General Target Regions:

[0362] (1) 5' Untranslated Region - nts 1 - 49 of GenBank Accession #  
NM\_001168:

ccgccagatt tgaatcgccg gacccgttgg cagaggtggc ggcggcggc (SEQ ID NO: 21)

[0363] (2) 3' Untranslated Region - nts 479 - 1619 of GenBank Accession #  
NM\_001168:

gg cctctggccg gagctgcctg gtcccagagt ggctgcacca ctccagggt ttattccctg gtgccaccag ccttcctgtg  
ggccccttag caatgtctta ggaaaggaga tcaacatttt caaattagat gtttcaactg tgctcctgtt ttgtcttgaa  
agtggcacca gaggtgcttc tgctgtgca gcgggtgctg ctggtaacag tggctgcttc tcctctctc tcctctttt  
gggggctcat ttgtctgtt ttgattcccg ggcttaccag gtgagaagtg agggaggaag aaggcagtgt ccttttgc  
agagctgaca gctttgttcg cgtgggcaga gcctccaca gtgaatgtgt ctggacctca tgtgttgag gctgtcacg  
tcctgagtgt ggacttggca ggtgcctgtt gaatctgagc tgcaggttcc ttatctgtca cacctgtgcc tcctcagagg  
acagttttt ttgtgtgtg ttttttgtt tttttttt ggtagatgca tgacttgtgt gtgatgagag aatggagaca gagtccctgg  
ctccttact gttaacaac atggctttct tatttgtt gaattgttaa ttcacagaat agcacaaact acaattaaaa  
ctaagcaca agccattcta agtcattggg gaaacggggg gaacttcagg tggatgagga gacagaatag agtcatagga  
agcgtctggc agatactcct ttgccactg ctgtgtgatt agacaggccc agtgagccgc ggggcacatg ctggccgctc  
ctccctcaga aaaaggcagt ggcctaaatc ctttttaaat gacttggctc gatgctgtgg gggactggct gggctgctgc  
aggccgtgtg tctgtcagcc caaccttcac atctgtcacg ttctccacac gggggagaga cgagtcgcg ccaggtcccc  
gctttcttg gaggcagcag ctcccgagg gctgaagtct ggcgtaagat gatggatttg attgcctc ctccctgtca  
tagagctgca ggggtgattg ttacagcttc gctggaaacc tctggaggtc atctcggtg ttctgagaa ataaaaagcc  
tgtcatttc (SEQ ID NO: 22)

### 6.7. Epidermal Growth Factor Receptor

[0364] See, e.g., GenBank Accession # XOO588.1.

General Target Regions:

[0365] (1) 5' Untranslated Region (247 nt) (GenBank Accession No. Hu EST  
gi|6302071|gb|AW163038.1|AW163038):

ccccggcgcagcgcggccgcagcagcctccgccccgcacgggtgtgagcggcgacgcggccgagggcgccggagtccc  
gagctagccccggcggccgccgcccagaccggacgacaggccacctcgtcggcgctccgccgagtcctccgctcggc  
caacgccacaaccaccgcgcacggccccctgactccgtccagtattgatcgggagagccggagcgagctctcggggagcagc  
ag (SEQ ID NO: 23)

[0366] (2) 3' Untranslated Region (1.7 kb, 58% AT-density):

tgaccacggaggatagtagccctaaaaatccagactctttcgataccaggaccaagccacagcaggtcctcatcccaacag  
ccatgcccgcattagctcttagaccacagactggtttgcaacgtttacaccgactagccaggaagtacttccacctgggcacattt  
tggaagtgtcattcctttgtcttcaaactgtgaagcatttacagaaacgcattccagcaagaatattgtccctttgagcagaaattatctt  
tcaaagaggtatattgaaaaaaaaaaaaaagtatatgtgaggattttattgattggggatcttgaggttttcattgtcgtattgattt  
tacttcaatgggctcttccaacaaggaagaagcttgctgtagcacttgctacctgagttcatccaggcccaactgtgagcaagga

gcacaagccacaagtctccagaggatgcttgattccagtggttctgcttcaaggctccactgcaaaacactaaagatccaagaag  
gccttcatggccccagcaggccggatcggtactgtatcaagtcatggcaggtacagtaggataagccactctgtcccttctgggca  
aagaagaaacggaggggatgaattcttcttagacttactttgtaaaaatgtcccacggtacttactccccactgatggaccagtgg  
ttccagtcatgagcgttagactgacttgttgtcttccattccattgtttgaaactcagtatgccgccctgtcttgcgtcatgaaatcag  
caagagaggatgacacatcaataataactcggattccagcccacattggattcatcagcattggaccaatagcccacagctgaga  
atgtggaatacctaaggataacaccgctttgttctcgcaaaaacgtatctcctaatttgaggctcagatgaaatgcatcaggtccttg  
gggcatagatcagaagactacaaaaatgaagctgctctgaaatctccttagccatcaccccaacccccaaaattagtttggttactt  
atggaagatagttttctcttttacttcaactcaaaaagctttttactcaaagagtatatgttccctccaggtcagctgccccaaacccct  
ccttacgctttgtcacacaaaaagtgtctctgccttgagtcatttcaagcacttacagctctggccacaacaggggcattttacaggt  
gcgaatgacagtagcattatgagtagtgaattcaggtagtaaatatgaaactagggttgaaattgataatgctttcacaacatttgca  
gatgttttagaaggaaaaaagtcttctctaaataatttctctacaattggaagattggaagattcagctagttaggagccatttttctt  
aatctgtgtgtccctgttaacctgactggtaacagcagtcctttgtaaacagtgttttaaaactctcctagtcaatatccaccccatccaat  
ttatcaaggagaagaatggttcagaaaatatttcagcctacagttatgttcagtcacacacacatacaaaaatgttcttttgcttttaaagta  
attttgactccagatcagtcagagccctacagcattgtaagaaagtatttgattttgtctcaatgaaaataaaactatattcattcc  
(SEQ ID NO: 24)

#### 6.8. CCAAT/Enhancer Binding Protein

[0367] See, *e.g.*, GenBank Accession # NM\_004364.

General Target Regions:

[0368] (1) CEBP- $\alpha$ , uORF (5'UTR, 160 nt):

Tataaaagctgggcccggcgccggggccggttcgcgacccggaggtgcgcggcgccggcgagcagggctccgggtgg  
gcggcgcgacgccccgcgcaggctggaggccgcccaggctcgccatgccgggagaactctaactccccatggagtcggc  
(SEQ ID NO: 25)

[0369] (2) CEBP- $\alpha$ , uORF (3'UTR, 1306 nt):

tgaggcgcgcggtgtgggaccgccctgggcccagcctccggcggggacccaggagtggtttggggcgccggatctcaggg  
cttggccagaccgtgcgagccaggactaggagattccggctgctcctgaaagcctggcctgctccgcgtgtccctccctcctctg  
cgccggacttggtgcgtctaagatgagggggccaggcggtggcttctcctgcgaggaggggagaattcttggggctgagctgg  
gagcccggaactctagtatttaggataactgtgccttggaatgcaaaactaccgctccaatgcctactgtagggggagcaaa  
tcgtgccttgatcttttatttgagggttctgctccttcccagggtacagcagacccccatgagagaaggaggggagcaggccc  
gtggaggaggggggctcaggagctgagatcccagacagcccgccagccccagccgctcctccacgcctgtccttagaaaggg  
gtggaacatagggacttggggcttgaacctaaaggtgttccctagtctacatgaaggtggaggtctctagtccacgcctctccca  
cctccctccgcacacacccccacccagcctgctataggctggctttcccttggggctggaactcactgcgatggggtcaccaggtga  
ccagtgagagccccaccccgagtcagaccagaaagctaggctgtgggtcagctctgaggatgtataccctgggtgggagagggga  
gacctagagatctggctgtggggcgggcatgggggtgaagggccactgggacctcagccttgtttgactgtatgccttcagcat  
tgcttaggaacacgaagcacgatcagtcacccagagggaccggagttatgacaagcttccaaatatttgctttatcagccgatat  
caacacttgatctggcctctgtgccagcagtgccctgtgcaatgtaatgtaccgtctctgctaaccaccattttatttggtttgttt

gtttggttttctcgatacttgccaaaatgagactctccgctcggcagctgggggaagggtctgagactctcttcttttggttttgggatt  
acttttgatcctgggggaccaatgaggtgaggggggttctcctttgccctcagctttcccagccctccggcctgggctgcccacaag  
gcttctccccagaggccctggctcctggcgggaaggaggtgcctcccgccaacgcactactggggctgggagcagggaag  
ggaattc (SEQ ID NO: 26)

#### 6.9. Cysteine-rich, Angiogenic Inducer, 61

[0370] See, *e.g.*, Gen Bank Accession # XM\_001831.

General Target Regions:

[0371] (1) 5' Untranslated Region (GenBank Accession No.

gi|19200928|gb|BM844529.1|BM844529):

agcgagagcgccccgagcagcgcccgccctccgcgccttctccggggacctcgagcgaaagacgcccggcgccgc  
ccagccctcgctccctgccaccgggcacaccgcgccgccaccccgaccccgctgcgcacggcctgtccgctgcacaccagc  
ttgttggcgtcttcgtcgccgcgctcgccccgggctactcctgcgcgccaca (SEQ ID NO: 27)

[0372] (2) 3' Untranslated Region (687 nt) (GenBank Accession No.

12898379|emb|AL556057.1|AL556057):

taaatgctacctgggtttccagggcacacctagacaaacargggagaagagtgatcagaatcagaatcatggagaaaatgggcggg  
gggtggtgtgggtgatgggactcattgtagaaaggaagccttgctcattcttgaggagcattaaggtatttcgaaactgccaaagggtgc  
tggtgcggatggacactaatgcagccacgattggagaatactttgcttcatagtattggagcacatgttactgcttcattttggagcttgc  
ggagttgatgactttctgtttctgtttgttaaattatttgctaagcatattttcttaggctttttcttttggggttctacagtcgtaaaagaga  
taataagattagttggacagtttaaagcittttatctcctttgacaaaagtaaatgggagggcattccatcccttctgaagggggaca  
ctccatgagtgctgtgagaggcagctatctgcactctaaactgcaaacagaaatcaggtgtttaagactgaatgtttatttatcaaaa  
ttagcttttggggaggggaggggaaatgtaatactggaataattgtaaatgattttaattttatattcagtgaaaagattttatttatggaat  
taaccatttaataaagaatatttacctaaaaa (SEQ ID NO: 28)

#### 6.10. Basis Fibroblast Growth Factor

[0373] See, *e.g.*, GenBank Accession No. NM\_002006.

General Target Regions:

[0374] (1) 5' Untranslated Region:

cggccccagaaaaccgagcgagtagggggcgccgcgcaggaggaggagaactggggcgccgggaggctgggtgggtgtc  
gggggtggagatgtagaagatgtgacgcccgcccgccgggtgccagattagcggacggctgcccgcggtgcaacgggatc  
ccgggcgctgcagcttgggagggcggtctccccaggcggcgctccgcggagacacccatccgtgaacccagggtcccgggccg  
ccggctcgccgcgcaccaggggccggcgacagaagagcgccgagcggtcgaggctgggggac (SEQ ID NO:  
29)

[0375] (2) 3' Untranslated Region (5.8 kb):

ctgctaagagctgattttaatggccacatctaattcatttcacatgaaagaagaagtatttttagaaatttgtaaatgagagtaaaagaa  
aataaatgtgtatagctcagtttgataatttggtcaacaatttttatccagtagtaaaatatgtaaccattgtcccagtaaaagaaaaata

acaaaagtgtaaaaatgtatattctcccttttataattgcatctgctgttaccagtgaaagcttacctagagcaatgatcttttcacgcatttg  
ctttattcgaaaagaggcttttaaatgtgcatgtttagaacaaaaattcttcatggaaatcatatacattagaaaatcacagtcagatgtt  
taatcaatccaaaatgtccactatttctatgtcattcgttagtctacatgtttctaacaatataaatgtgaatttaaatcaattcctttcatagttt  
tataattctctggcagttccttatgatagagtttataaaacagtcctgtgtaaactgctggaagttctccacagtcagggtcaattttgtcaa  
acccttctctgtaccatacagcagcagcctagcaactctgctggatgggagtgattttcagcttcgccagggtcattgagatcca  
tccactcacatcttaagcatttctctggcaaaaatttatggatgaatgaataggctttaggcggcagatgatatacatatctgacttcca  
aaagctccaggatttgtgtgctgttgcgaatactcaggacggacctgaattctgattttataccagctcttcaaaaacttctcgaaccg  
ctgtgtctcctacgtaaaaaagagatgtacaaatcaataataattacacttttagaaactgtatcatcaaagattttcagttaaagtagca  
ttatgtaaagggtcaaaacattaccctaacaaagtaaagtttcaatacaaaattcttgccttggatatacaagaaatccaaaatatttct  
ttaccactgtaaattcaagaagcttttgaaatgctgaatatttcttggctgctacttggaggcttatctacctgtacattttggggtcagct  
cttttaacttctgctgctcttttccaaaaggtaaaaatatagattgaaaagttaaaacattttgcatggctgcagttcctttgttcttga  
gataagattccaaagaacttagattcatttctcaacaccgaaatgctggagggtgttgatcagtttcaagaaacttggaaatataaataat  
ttataattcaacaaagggtttcacattttataagggtgattttcaattaaatgcaaatgtgtggcaggattttattgccattaacatatttt  
gtggctgcttttctacacatccagatggctcctctaactgggcttctctaattttgtatgttctgtcattgtctcccaagatttttaggag  
aagcccttaaaaagctgccttctctaccatttctgtgaaagcttcacaattgtcacagacaaagattttgttccaatactcgttttgc  
ctctatttttctgttgtcaaatagtaaatgatatttgccttgcagtaattctactggtgaaaaacatgcaaagaagaggaagtcacaga  
aacatgtctcaattcccatgtgtgtgactgttagactgtcttaccatagactgtcttaccatcccctggatagctcttgttttccctcta  
atagctatggaaagatgcatagaaagagtataatgttttaaacataaggcattcatctgccattttcaattacatgctgacttcccttac  
aattgagatttgcctatagggttaaacatgggttagaacaactgaaagcataaaagaaaaatctaggccgggtgcagtggtcatgcct  
atattccctgcactttgggaggccaaagcaggaggatcgtttagcccaggagttcaagaccaacctgggtgaaacccgtctctac  
aaaaaacacaaaaatagccaggcatgggtggcgtgtacatgtgtctcagatacttgggaggctgagggtgggagggttgatcact  
tgaggctgagagggtcaaggttgcagtgagccataatcgtccactgcagtcacgcctaggcaacagagtgcagcttctcaaaa  
aaagagaaatttcttaataagaaaagtaattttactctgatgtgaatacatttgttattaaattattatttaagatggtagcactagtctt  
aaattgtataaaatatccctaacaatgtttaaatgtccattttattcattatgcttgaataaataattatggggaaatacatgtttgttataaa  
ttattattaaagatagtagcactagtcttaatttgatataacatctcctaactgtttaaattgtccattttattctttatgcttgaataaatt  
atggggatcctatttagctcttagtaccactaatcaaaagttcggcatgtagctcatgatctatgctgttctatgtcgtggaagcaccgg  
atgggggtagtgagcaaatctgccctgtcagcagtcaccatagcagctgactgaaaatcagcactgcctgagtagtttgatcagtt  
taacttgaatcactaactgactgaaaattgaatgggcaataagtctttgtctccagaglatgcgggagaccctccacctcaagat  
ggatatttctccccaaggattcaagatgaattgaaattttaatcaagatagtgtgctttattctgttgatttttattatttaataactgta  
agccaaactgaaataacatttgcgttttataggttgaagaacataggaaaaactaagagggtttgttttttttgcgtgatgaagagata  
tgtttaaatatgttgattgtttgttttagttacaggacaataatgaaatggagtttatatttgttatttctattttgttatatttaataatagaatta  
gattgaaataaaaataatgggaaataatctgcagaatgtgggttccctgggtttcctctgactctagtgcactgatgatctctgataag  
gctcagctgctttatagttctctggctaattgcagcagatacttctcctgccagtggtaatacagatttttaagaaggcagtttgcatttta  
atcttgggatacctttatactcttagggattattttatacaaaagccttgaggattgcattctattttctatatgacctcttgatatttaaaa  
aacactatggataacaatttctcatttacctagtattatgaagaatgaaggagttcaacaaatgtgttccagtttaactagggttact

gtttgagccaataataatgtttaactgtttgtgatggcagttatcctaagtacattgcatgtttcctaataacagagtttaataatttcagt  
aattcttagatgattcagcttcattcattaagaatatctttgtttatgttgagttagaatgccttcatatagacatagctttcagacctctac  
tgtcagttttcatttctagctgcttcagggtttatgaattttcaggcaaaagctttaatttataactaagcttaggaagtatggctaatagccaa  
cggcagttttttcttctaattccacatgactgagggcatatatgatctctgggtaggtgagttgtgtgacaaccacaagcacttttttttt  
ttaagaaaaaaaggtagtgaattttaatcatctggactttaagaaggattctggagtatacttaggcctgaaattatatatttgcttg  
gaaatgtgtttttctcaattacatctacaagtaagtacagctgaaattcagaggaccataagagttccatgaaaaaatcaattcatt  
gaaaaggcaagatgcaggagagaggaagccttgcaaacctgcagactgcttttcccataatagattgggtaaggctgcaaaaca  
taagcttaattagctcacatgctctgctctcacgtggcaccagtggatagtgtagagaattaggctgtagaacaaatggccttctctt  
cagcattcacaccactacaaaatcatcttttatcaacagaagaataagcataaactaagcaaaaggtaataagctgaaacca  
agattggctagagatatcttaatgcaatccattttctgatggattgtacgagttggctatataatgtatgtatggattttgattgtgtaa  
aagttttaaaaaatcaagctttaagtacatggacatttttaataaaaatatttaagacaatttagaaaatgccttaatatcattgttgctaa  
atagaataggggacatgcataattaaggaaaaggctatggagaaataatattggtatcaacaaatacattgattgtcatgatacacatt  
gaattgatccaatagtttaaggaataggtaggaaaatttggttctatttttcgatttctgtaaatcagtgacataaataattcttagcttat  
tttatatttctgtcttaataactgagctcagtaagttgtgttaggggattatttctcagttgagactttcttatatgacattttactatgtttga  
cttctgactattaaaaataaatagtagaacaaatttcataaagtgaagaattatataatcactgctttataactgactttattatattttc  
aaagttcatttaagggtactattcatcctctgtgatggaatggcaggaattgttttctcatagtttaattccaacaacaatattagtcgta  
tccaaaataacctttaatgctaaactttactgatgtatatccaaagcttctcttttcagacagattaatccagaagcagtcataaacagaa  
gaataggtggatgttctaatgatatttttactaatggaataaactgtaataatagaattatgctgctaattatcagctctgaggtat  
atttctgaaatgttcagactcagtcggaacaaattggaaaatttaatttttattcttagctataaagcaagaaagtaaacacattaattcc  
tcaacatttttaagccaattaaaaatataaaagatacacaccaatatcttcttcaggctctgacaggcctcctggaaactccacatatttt  
tcaactgcagtataaagtcagaaaataaagttaacataactttcactaacacacacatatgtagatttcacaaaatccacctataattggt  
caaagtgggtgagaatatatttttagtaattgcatgcaaaattttctagcttccatcctttctccctcgtttctcttttttgggggagctggt  
aactgatgaaatcttttccaccttttcttccaggaaatataagtgtttgtttggttaacgtgatacattctgtatgaatgaacattgga  
gggaaacatctactgaatttctgaatttaaaatattttgctgctagtttaactgaacagatagaagaatcttacagatgctgctataaat  
aagtagaaaaataaatttcactactaaaatgctatttttaaatctatttctatatattgtatttctaatcagatgtattactcttattttctatt  
gtatgtgtaatgattttatgtaaaaatgaattgcttttcatgagtagtatgaataaaattgattagttgtgttttctgtctcccgaaaaaa  
aaaaaaaaaaaaaaaaaaaaaa (SEQ ID NO: 30)

### 6.11. Cyclin D1

[0376] See, *e.g.*, GenBank Accession No. NM\_053056.

General Target Regions:

[0377] (1) 5' Untranslated Region:

cggccccagaaaacccgagcgagtagggggcggcgcgcaggaggaggagaactggggggcgcgggaggctggtgggtgtc  
gggggtggagatgtagaagatgtgacgccgcggccggcgggtgccagattagcggacggctgcccgcggttgaacgggatc  
ccgggcgctgcagcttgggaggcggctctcccaggcggcgtccgcggagacacccatccgtgaacccaggctccggggccg

**[0378] (2) 3' Untranslated Region (3.2 kb):**

- 137 -

atcctggctgcggcgtctgtctgaaccacgcggggccttgaggacgctttgtctgtcgtatggggcaagggcacaaagtcctgg  
 atgttgtgtatcgagaggccaaaggctggtggcaagtgcacggggcacagcggagtctgtcctgtgacgcgaagtctgaggg  
 tctgggcggcgggcggtggtctgtgcatttctggttcaccgcggcgcttccagcaccaacatgtaaccggcatgtttccagca  
 gaagacaaaaagacaaacatgaaagtctagaaataaaactggtaaaacccccaaaaaaaaaaaaaaaaaaaaa (SEQ ID NO: 32)

## 6.12. Murine Double Minute 2

[0379] See, e.g., GenBank Accession No. NM\_002392.

General Target Regions:

[0380] (1) 5'End/Intron 1/p53 BS for s-mdm-2: U39736:

gcaccgcggcgagcttggtgcttctggggcctgtgtggccctgtgtgtcgaaagatggagcaagaagccgagcccagggggc  
 ggccgcgacccctctgaccgagatcctgctgcttcgcagccaggagcaccgtccctccccggattagtgcgtacgagcgcgccag  
 tgccctggcccgagagtggaatgatccccgagggccaggcgctgcttccgcgcgcccgctgaaggaaactggggagtctt  
 gagggacccccgactccaagcgcgaaaaccccgatggtgaggagcaggtactggccggcagcgagcggtcacttttgggtc  
 tgggctctgacggtgtcccctctatcgtggttccagcctctgccggttcgcagcctttgtgcggttcgtgntctgggggctc  
 gcggggcgcgggggcatgggncacgtggctttgcggaggtttgttgactggggctagacagtccccgccaggaggaggcg  
 ggatttcggacggctctcgcggcggtgggggtgggggtggttcggaggtctccgcgggagttcagggtaaaggtcacggggcc  
 ggggtcgcggggcgccttcggcgcgggaggtccggatgatcgagtcctgtcgggtcactagtgtgaacgtcgcgtagtctg  
 ggcgggattgggccggttcagtgggcaggttgactcagcttttctcttgagctggtcaagttcagacacgttccgaaactgcagtaa  
 aaggagttaagtcctgactgtctccagctggggctatttaaacatgcattttccagctgtgtCAGTGGCGATTGGAG  
 GGTAGACCTGTGGGCACGGACGCACGCCACTTTTTCTCTGCTGATCCAGgtaagcac  
 cgacttgctttagcttttagtttaactgttgttatgttctttatatatgatgtattttccacagatgttcatgatttccagtttcatcgtgtctt  
 ttttctttagtgcaaatgtgcaataccaacatgtctgtacc (SEQ ID NO: 33)

[0381] (2) 3'UTR (GenBank Accession No.

gi|9150029|gb|BE275079.1|BE275079):

tagtgacctgtctataagagaattatataatttctaactatataaccctaggaattagacaacctgaaatttattcacatatataaagtga  
 gaaatgcctcaattcacatagatttcttctttagtataattgacctactttggtagtggaaatagtgaatacttactataatttgacttgaat  
 atgtagctcatcctttacaccaactcctaattttaataatttctactctgtcttaaataagaaagtacttggttttttttcttaaatatgtatatg  
 acatttaaatgtaacttattttttttgagaccgagcttctgtctgttaccaggtgagtgagtgagggtgatcttggtcactgcaag  
 ctctgccctccccgggttcgaccattctctgcctcagcctcccaattagcttggcctacagtcacttccaccacacctggctaattt  
 ttgtacttttagtagagacagggttcaccgtgttagccaggatggtctcgtatcctgacctcgtgatccgccacctcggcctcca  
 aagtgtctgggattacaggcatgagccaccgtgctctccagcctaggcaacagagtgagactctgtctccaaaaaaaaaaaaaaaaaa  
 aaggggactataacacccccagggaaggacaggtgggacattcttattcttaatttaataaattgacaggggaaagtgggcca  
 ctcttgagcttgtgggtgctcaccaggttgacccccaaaaaagaagccttcacaaaacattatttatttccctaataataccgcctot  
 gtgagttaagggataatgcatcaggactcttgcaaccagacaaaattatttaaaaacgccacttgggggggaggcggggtccctctg  
 gggattcgccttgtgggagagaaaactgcacagacttgggcaaataatgtttttgtcaccacaaaacgtattcgcgagacatttcatt



agaacgaagctttaccctaattgaactccccatttaaacagttccacacacacttagggagattttccctctgtgagttccgcagaa  
caatagttggacgggaatagaaccctgaaacactttagttcaccacgaactattataggcgagg (SEQ ID NO: 34)

### 6.13. Protein Tyrosine Phosphatase Type IVA, Member 3

[0382] See, *e.g.*, GenBank Accession No. NM\_032611.

General Target Regions:

[0383] (1) 5' Untranslated Region:

tgactatccagctctgagagacgggagtttgaggtgcccgttactttgggtgggtggggggggcgggcggtgtttgttcctttt  
ctttttaagagttgggttttctttttaattatccaaacagtgggcagcttctccccacaccaagtatgtgcacaaatgtgcgggggt  
atgggggtgggttttaaatctcgtttctcttgacaagcacaggatctcgttctcctcatttttgggggtgtgtggggacttctcaggt  
cgtgtccccagccttctctgcagtccttctgccctgccggggcccgtcgggagggcgc (SEQ ID NO: 35)

[0384] (2) 3' Untranslated Region:

tagctcaggaccttggtgggcctggctgtcatgtaggtcaggaccttggtggacctggaggccctgccagccctgctctgcc  
agcccagcaggggctccaggccttggtggccccacatgccttttctccccgacacctccgtgcactgtgtccaggagcgag  
gagccctcggggcctgggtggcctctgggcccttctcctgtctccgccactccctctggcggcgtggccgtggctctgtctct  
gaggtgggtcgggcgcctctgcccgcctccacaccagccaggtggtctcctctagcctgtttgtgtgggggtgggggtat  
atgttaaccactgggccccagccctcttttgcacccctgtcctgacctgttctggcaccttaaaattattagacccggggcag  
tcaggtgctccggacacccgaaggcaataaaacaggagccgtgaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaa  
aaaaaaaaaaaaaaaaaaaaaaaaaaaa (SEQ ID NO: 36)

### 6.14. Tissue Inhibitor of Metalloproteinase

[0385] See, *e.g.*, GenBank Accession No. XM\_003061 for TIMP-4.

General Target Regions:

[0386] (1) 5' Untranslated Region (GenBank Accession No.

gi|11293824|gb|BF346229.1|BF346229):

gctcagcaaggggtccgtccttctctgtcactgtctctttgctgttgtaattctgtctgcctctctgggactctgcctgtctcactttct  
gtctgtgcctctcctcactctgttcttctgctgaatcacagccctcagttttctgtctcatgcattgtcttttggtctttccgtcttc  
tgcccttgacaccatccctctcccagtgcttccctctgcttcagatcgcttcatgacttaggcagggaacagaggtcagggcct  
ccttcaggcttccctctgcacttactgagtatgcaggctcgaagagcctcgggtcctgcctccgcgggtggcctagagccaaagg  
aaggcggagcccgtcggggcgggattggcccttagggccacctcataaagcctggggcgaggggcacaacggccttgggaag  
gagccctgctggggcgtccagtcccccagacctcacaggctcagtcgcggatctgcagtgct (SEQ ID NO: 37)

[0387] (2) 3' Untranslated Region:

tagtagggaccagtgaccatcacatcccttaagagtcctgaagatcaagccagttctcctccctgcagagcttggccattaccac  
ctgacctctgtgccagctaataagaagtgccaaagtggacagctggccactgtcaaggcagggaaggggccatgacttttctgcc  
ctgccctcagcctgttgccctgcctccaaacccattagcttagcctgttagctgttactgcaagtgttcttctggcttagctgtttct  
aaagccaggactattcccttctccccaggaatatgtgttcttcttgaatcgatctggtaggggagaaatggcgaatgtcataca

catgagatggtatctccttgcgatgtacagaatcagaaggtggttgacagcatcataaacaggctgactggcaggaatgaaaaaa  
aaaaaaaaaaaa (SEQ ID NO: 38)

[0388] See, *e.g.*, GenBank Accession No. S48568 for TIMP-2.

General Target Regions:

[0389] (1) 5' Untranslated Region (84 % GC-rich):

ggggccgccgagagccgcagcgccgctgccccgcccccccccccagccgccccggcggaattgcgccccgcgcct  
cccctcgcgccccgagacaaagaggagagaaagtttgcgcggccgagcgggcaggtgaggagggtgagccgcgcggagg  
ggccccgctcggccccggctcagccccgcccgcgccccagcccgccgcccgcgagcagcgccccggacccccagcggcg  
ccccgccccgcccagccccggccccg (SEQ ID NO: 39)

[0390] (2) 3' Untranslated Region (GenBank Accession No.  
18505971|gb|BM456931.1|BM456931):

taagcaggcctccaacgccccgtggccaactgcaaaaaagcctccaagggttcgactggtccagctctgacatcccttctgga  
aacagcatgaataaaactcatcccatgggtccaaattaatgattctgctcccccttctccttttagacatggttggtggtctggag  
ggagacgtgggtccaaggctcatcccatcctcctctgccaggcactatgtgtctggggcttcgatccttgggtgcaggcagggc  
tgggacacgcggcttccctccagtccttgccttggcaccgtcacagatccaagcaggcagcacttagggatctccagctgggt  
tagggcagggcctggaaatgtgattttgcagaaactttgagggctgttgcaagactgtgtagcaggcctaccaggctcccttcatct  
tgagagggacatggccccctgttttctgcagcttcacgcctctgcactccctggccttgcaagtgtctccatcgccccgggtgcc  
accatgnagctccccgcacctgactccccccacatccaagggcagccctggaaccagtgggctagttccttgaaggaaagccccac  
tcattcctattaatccctcagaattcccgggggagccttccctcctgaaccttgtaaaaaatggggaacgagaaaaacccccgctt  
ggagctgtgcgttccagccccctacttgagagncttttttggggggccg (SEQ ID NO: 40)

#### 6.15. Peroxisome Proliferative Activated Receptor-g

[0391] See, *e.g.*, GenBank Accession No. NM\_138712.

General Target Regions:

[0392] (1) 5' Untranslated Region (GenBank Accession No.

12786927|emb|AL523434.1|AL523434):

cgcgccggggcccggtcggcccgaccggctccgcgcgggcagggcgcccgagcgccactcgagcccgagcccgagccg  
cagccgccgctggggcgcttgggtcggcctcaggacaccggagagggggccacgccgcccgtggccgcagatttgaaga  
agccgacactaaaccaccaatatacaacaaggccattttgtcaaagagagtcagccttaacgaaa (SEQ ID NO: 41)

[0393] (2) 3' Untranslated Region:

tagcagagagtcttagccactgccaacatttcccttctccagttgcactattctgagggaaaatctgacacctaagaaatttactgtg  
aaaaagcattttaaaaagaaaagggttttagaatatgatctattttatgcatattgtttataaagacacatttacaatttttaattaaaa  
attaccatattatgaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaa (SEQ ID NO: 42)

#### 6.16. PC-cell Derived Growth Factor/Epithelin/Granulin Precursor

[0394] See, *e.g.*, GenBank Accession No. NM\_002087.

General Target Regions:

[0395] (1) 5' Untranslated Region:

ggcacgaggggagagaggaagcagggaggagagtgtttgagtagaaaagaaacacagcattccaggctggccccacctctat  
attgataagtagccaatgggagcgggtagccctgatccctggccaatggaaactgaggtaggcgggtcatcgctgggggtctgta  
gtctgagcgctaccgggtgtgtgctgccaaggaccgcggtgagtcggacgcaggcagaccatgtggaccctgggtgagctgggtgg  
ccttaacagcaggggtgtgtggtggaacgcggtgccagatgtcagttctgcctgtggcctgctgcctggaccccgaggagc  
cagctacagct (SEQ ID NO: 43)

[0396] (2) 3' Untranslated Region:

tgagggacagtactgaagactctgcagccctcgggacccactcggagggtgccctctgctcaggcctccctagcacctcccccta  
accaaattctccctggacccattctgagctccccatcacatgggaggtggggcctcaatctaaggccttcctgtcagaagggg  
ttgtggcaaaagccacattacaagctgccatccctccccgttcagtggaccctgtggccagggtgtttccctatccacaggggtg  
ttgtgtgtgtgcgcgtgtgcgttcaataaagttgtacactttcaaaaaaaaaaaaaaaaaaaaaaaaaaaaaa (SEQ ID NO:  
44)

### 6.17. Angiogenin

[0397] See, *e.g.*, GenBank Accession No. M11567.

General Target Regions:

[0398] (1) 5' Untranslated Region:

tgtttgcattaagtcatagattataattgtaatggaatcaacaccaaagcaaattagaaagagagcccactttgtcaccagtcacg  
tcttcccatgtaaccatagaacgttgggtcctgtgtcttctagatccacagcttctgtctcagaacaggctagccacaccacaggcc  
tagtgccaggaccatggccttttttaagctcagactcccttctgtgaacagcaatatccccacaactgttacaacattggtgcttctg  
caaggggtacagaactattgatacgaatgttcattgacttacacacaagagaagcacaaaaataaaaaattaataattaattatgt  
cttgaaaaatgtaccattatttttacatttgggtcataagaattgtattacacttaagaatgcaatacaattgaagatcagattttctccc  
tttgtgagaatttctcagtatgtgtgatgactaccaagaatcatagccagtcataaattcagtgagttactcataaacaagaac  
acctacttcttggggaggttaggtctgcttccctcaactcaggatacaactgcttcaactgctttcttcacattagctgactaattagcta  
gaagcctgtcgtaaacaattttatggttgactccttccctgggctcagggttcctagaacagagaggtcccaaatcccggtctgtg  
gcctgtccgcctaagctctgcctcctgccagatcagcaggcagcattagattctcataggagctggacgcctattgtgaactgcgc  
gtgcgggatccagattgtgcactctttatgagaatctaactaatgcttgatgatctatctgaaccagaacaatttcacctgaaaccatcc  
cccaccaatccatagaaatactgtcttccacaaaaatgatccctgggtgcaaaaatgttagagaccactcccctaaaactctctcttag  
ctctcacctcctgtattactatctcatctcagtacattgaagccccatctttcccatggatgcctcatttctattaggaggagcattttt  
atTTTTgttttttttttccgagacggagtcctcgtctgtcgcaaggctggagtgcagtggcgcgatctcggctcactgcaagctcc  
gcctccgggttcacgccattctcctgcctcagcctcccaagtagctgggactacaggcggccgactacggccggctaatttttgt  
atTTTtagtagagacggggttaccgtggttagccaggatggctcctgatctcctgacctgtgatccggccgcttggcctcccaagt  
gctgggattacaggcgtgagaccgcggccggcgtcatttggatgtcttaattgtgcctcaggacctagcacagtcctggtaccca  
gtagagacctatgtaattgtctgtattcaataataatacatgaattaaagagtgagagtgattttgtaattgttacgactgatagagaaa

[0399] (2) 3' Untranslated Region:

### 6.18. Hypoxia-inducible Factor- $\alpha$

General Target Regions:

[0401] (1) 5' Untranslated Region:

tcttcagtgacagtgctgcctcgtctgaggggacaggaggatcacctcttcgtcgttcggccagtggtgcgggctgggccctga  
caagccacctgaggagaggtcggagccggggccggacccggcgattgccgcccgttctctctagctcacgaggggttccc  
gcctcgcacccccacctctggacttgcccttctctctctccgcgtgtggagggagccagcgcttaggccggagcgagcctggg  
ggccgcccgcgtgaagacatcgcggggaccgattcacc (SEQ ID NO: 47)

[0402] (2) 3' Untranslated Region (GenBank Accession No.

gi|19116743|gb|BM799920.1|BM799920):

tgagcttttcttaatttcattcctttttggacactgggtggctcactacctaagcagtcatttatattttctacatctaatttagaagcctg  
gctacaatactgcacaaacttggttagtcaattttgatccctttctacttaattacattaatgctcttttttagtatgttcttaagtctggat  
cacagacagctcattttctcagtttttggatttaaacattgcattgcagtagcatcatttaaaaaatgcacctttttattttttttgg  
ctaggaggttatcccttttcgaattttttaagaagatgccaatataattttgtaagaaggcagtaacctttcatcatgatcataggca  
gttgaaaaattttacaccttttttcacattttacataaataataatgctttgccagcagtagctggtagccacaattgcacaatatatttct  
taaaaaataccagcagttactcatggaatatattctgcgtttataaaactagttttaagaagaaattttttggcctatgaaattgttaaac  
ctggaacatgacattgttaatcatataataatgattcttaaatgctgtatggtttatttttaaatgggtaaagccattacataatatagaaa  
gatatgcataatctagaaggatgtggcatttattggataaaattctcaattcagagaaatcatctgatgtttctatagtcactttgccag  
ctcaaaagaaaacaataacctatgtagttgtggaagttatgctaattgtgtaactgatattaaacctaaatgttctgcctacctgttg  
gtataagatatatttgagcagactgtaacaagaaaaaaatcatgcattcttagcaaaattgcctagtagttaattgtctaaaata  
caatgtttgattttatgcactttgtcgtattaacatcctttttcatgtagattcaataattgagtaattttagaagcatttttaggaatata  
tagttgtcacagtaaatatctgtttttctatgtacattgtacaaattttcattccttttgcctttgtggttgatctaactaactgtattgtt  
ttgttacatcaataaacatcttctgtggaccaggaaaaaa (SEQ ID NO: 48)

### 6.19. Large Tumor Suppressor, Homolog 1

[0403] See, e.g., GenBank Accession No. XM\_015547.

General Target Regions:

[0404] (1) 5' Untranslated Region (GenBank Accession No.

gi|19008744|gb|BM695486.1|BM695486):

agacagccttaaccacgggcgcgggcgagtcgtatgggcaggggcaggcgggagcgacgtggggcgacgctcacgaacga  
tcagagctgcgggcgacgcaacgaagcccgaggccgcaggctgcgcgtccctcgagcagccgggcgggcaaaagcccc  
cagtcctcggccccgcgcaagcgacgccgggaaa (SEQ ID NO: 49)

[0405] (2) 3' Untranslated Region (GenBank Accession No.

gi|12274655|gb|BF884528.1|BF884528):

taattatttatattgtaaagaatttaacagtcctggggacttcttgaaggatcattttcacttttgctcagaagaagctctggatctatca  
aataaagaagtccttcgtgtgggctacatatagatgtttcatgaagaggagtgaagccagaaggatatagacaaatgaggcct  
aagacctttcctgccagtaactatactgtcagtagccggcaaatgttacaagaaattcgggaatcccttaggaatttatctaaacctct  
gatgctgctaaggctgagcataacatgagtaaaatgtcaaccgaagatcctcgacaagtcagaaatccacccaaatttgggacgcat  
cataaagccttgaggaaattcgaaactctctgcttccatttgcaaatgaaacaaattcttctcgagtagtacttcagaagttaattccacaa

atgcttcaagacttgcaagctgctggatttgatgaggatatggtatacaagctcttcagaaaactaacaacagaagtatagaagcag  
 caattgaattcattagtaaaatgagttaccaagatcctcgacgagagcagatggctgcagcagctgccagacctattaatgccagcat  
 gaaaccagggaatgtgcagcaatcagttaaccgcaaacagagctggaaagggttctaaagaatccttagttcctcagaggcatggcc  
 cgccactaggagaaagtgtggcctatcattctgagagtcccaactcacagacagatgtaggaagacctttgtctggatctggtatc  
 agcatttgttcaagctcaccttagcaacggacagagagtgaacccccaccaccacctcaagtaaggagtgttactcctccaccac  
 ctccaagaggccagactccccctccaagagggtacaactccacctccccctcatgggaaccaaactctcaaaacaaagcgctattctg  
 gaaacatggaatacgtaatctcccgaatctcctgtcccacctggggcatggcaagaggggtatcctccaccacctctcaacacttc  
 ccccatgaatcctcctaatacaggacagagaggcattagttctgttctgttggcagacaaccaatcatcatgcagagttctagcaaat  
 ttaactttccatcaggagacctggaatgcagaatggtactggacaaactgatttcatgatacaccaaaatgttgtccctgctggcact  
 gtgaatcggcagccaccacctccatctctgacagcagctaattggacaaagcccttctgctttacaacagggggatctgctgct  
 ccttcgtcatatacaaatggaagtattcctcagctctatgatgggtgccaacagaaatagtcataacatggaactatataacattagtgtac  
 ctggactgcaaacaattggcctcagtcattcttctgctccagccagtcacccccgagcagtgggcatgaaatccctacatggcaac  
 ctaacataccagttaggtcaaattcttttaataaccattaggaatagagcaagtcactctgctaattctcagccttctgctacaacagt  
 cactgcaattacaccagctcctattcaacagcctgtgaaaagtatgcgtgtattaaaaccagagctacagactgctttagcacctacac  
 acccttcttgataccacagccaattcaaaactgttcaaccagtccttttctgaggggaaccgcttcaaatgtgactgtgatccacctg  
 ttgctgaagctccaaactatcaaggaccaccaccacctaccctacccaaaacatctgctgcacaaaacccatctgttctccatagcagtc  
 aatcagtaagcctagcaaagaggatcagccaagcttgcccaagggaagtgcagagtgaaaagagttatgaaaatgttgatagtgagg  
 gataaagaaaagaaacagattacaacttcacctattactgttaggaaaaacaagaaagatgaagagcgaagggaatctcgtattcaa  
 agttattctcctcaagcatttaaaattctttatggagcaacatgtagaaaatgtactcaaatctcatcagcagcgtctacatcgtaaaaaac  
 aattagagaatgaaatgatcggggttgattatctcaagatgccaggatcaaatgagaaaagatgctttgccaaaaagaatctaatta  
 catccgtcttaaaagggtctaaaatggacaagtctatgtttgtgaagataaagacactaggaataggagcatttggtgaagtctgtctag  
 caagaaaagtagatactaaggctttgtatgcaacaaaaactcttcgaaagaaagatgttcttctcgaaatcaagtcgctcatgtaagg  
 ctgagagagatatcctggctgaagctgacaatgaatgggtagttcgtctatattattcattccaagataaggacaattatactttgtaat  
 ggactacattcctgggggtgatgatgagcctattaattagaatgggcatctttccagaaaagtctggcacgattctacatagcagaac  
 ttacctgtgcagttgaaagtgttcataaaatgggttttattcatagagatattaaacctgataatattttgattgatcgtgatggtcatattaa  
 attgactgactttggcctctgcactggcttcagatggacacacgattctaagtactatcagagtgggtgacctccacggcaagatagc  
 atggatttcagtaatgaatggggggatccctcaagctgtcgtatgtggagacagactgaagccattagagcggagagctgcacgcc  
 gcaccagcgtatgctagcacattcttgggtgggactcccaattatattgcacctgaagtgttgctacgaacaggatacacacagttgt  
 gtgattggtggagtgttgggtgtattcttttgaaatgttgggtgggacaacctcttcttggcacaacaccattagaaacacaaatgaa  
 ggtcacctgctgctatatacatcattggctcgagaagaaactactgaacacctgcgagagagaagcctagaaaagaaagaaagg  
 gccaaaagggtttgaactcttcatccctaatttgctacactgatcaaaaccaagtaagggtcctgaagtccatgagctatcatcaatc  
 agcacaatgctatactagtttgtaactcgggggtcagttgtgaagggaaggacagcagcttattccatattccaggaagccacagt  
 aaactgctcga (SEQ ID NO: 50)

## 6.20. P-Glycoprotein

[0406] See, *e.g.*, GenBank Accession No. M14758.

General Target Sequences:

[0407] (1) 5' Untranslated Region:

cctactctattcagatattctccagattcctaagattagagatcatttctcattctcctaggagtactcacttcaggaagcaaccagataa  
 aagagaggtgcaacgggaagccagaacattcctcctggaaattcaacctgtttcgcagtttctcgaggaatcagcattcagtcattccg

ggccgggagcagtcacatctgtggtgaggctgattggctgggcaggaacagcgccggggcgtgggctgagcacagcgcttcgctct  
ctttgccacaggaagcctgagctcattcgagtagcggctctccaagctcaaagaagcagaggccgctgttcgtttccttaggtcttt  
ccactaaagtcggagtagtcttctccaagatttcacgtcttggtggccgtccaaggagcgcgaggtcggg (SEQ ID NO:  
51)

[0408] (2) 3' Untranslated Region (GenBank Accession No.:

gi|13334786|gb|BG428280.1|BG428280):

tgaactctgactgtatgagatgttaaatacttttaataattgttagatatgacatttattcaaagttaaagcaaacttacagaattatga  
agaggtagctgtttaacatttcctcagtcaggttcagagcttcgtaattaaaggaacagagtgagagacatcatcaagt  
gagagaaatcatagttaaactgcattataaattttataacagaattaaagtagattttaaaagataaaatgtgtaatttgtttatatttccc  
atttgactgtaactgactgccttgctaaaagattatagaagtagcaaaaagtattgaaatgttgcataaagtgtctataataaaactaa  
actttcatgtgactggagtcattctgtccaaactgcctgtgaatatatcttctcaattggaatattgtagataactctgtttaaaaaagt  
ttctttaaatatacctactcattttgtgggaatgggttaagcagtttaataattcctgtgtatgtctatcacataggggtctaacagaac  
aatctggattcattatttctaggacttgatcctgctgatgctgaattgcacattaaggtgtgtaacaacacacagatcgatataa  
gaagtaaggaggtggggagaggcaattatgatgtgctatgagttagatgtatagt (SEQ ID NO: 52)

#### 6.21. CD82 Antigen

[0409] See, *e.g.*, GenBank Accession No. NM\_002231

General Target Regions:

[0410] (1) 5' Untranslated Region (GenBank Accession No.

gi|19088880|gb|BM759265.1|BM759265):

agtccgcggcgttccccggctgcagccgggagggggccgaggagtactgagccccgggctgtgcagtccgacgccgactga  
ggcacgagcgggtgacgtgggcctgcagcgcggagcagaaagcagaacccgcagagtcctccctgctgtgtgtggacgaca  
cgtgggcacaggcagaagtgggccctgtgaccagctgcactggttcgtggaaggaagctccaggactggcggg (SEQ ID  
NO: 53)

[0411] (2) 3' Untranslated Region:

tgaggcagctgctatcccc

atctccctgcctggcccccaacctcagggtctccagggtctccctggctccctcctccaggcctgcctcccacttcactgcgaaga  
ccctcttggccaccctgactgaaagtagggggctttctggggcctagcgatctctctggcctatccgctgccagccttgagccctgg  
ctgttctgtggttctctgtcaccgcccacaggggttcttataactcagagaaaaatgctccccacagcgtccctggcgcaggtg  
ggctggacttctacctgccctcaagggtgtgtatattgtataggggcaactgtatgaaaaattggggaggagggggccgggcgcgg  
tgctcacgcctgtaatccagcactttgggaggccgaggcgggtggatcacgaggtcaggagatcgagaccatctggctaact  
gggtgaaaccccgctctactaaaaatacaaaaaaatttagccgggcgcgggtggcgggcacctgtagtcccagctacttgggagggc  
tgaggcaggagaatggtgtgaacccgggagcggaggttcagtgagctgagatcgtgctactgcactccagcctggggggacaga  
aagagactccgttcaa (SEQ ID NO: 54)

#### 6.22. Bcl-2

[0412] See, *e.g.*, GenBank Accession No. M14745

General Target Regions:

[0413] (1) 5' Untranslated Region (GenBank Accession No. gi|19887364|gb|BQ061909.1|BQ061909):

tttctgtgaagcagaagtctgggaatcgatctggaaatcctcctaattttactccctctcccccgactcctgattcattgggaagttca  
aatcagctataactggagagagctgaagattgatgggacgttgcccttatgcctttgttttggtttacaaaaaggaaacttgacagagg  
atcatgctatacttaaaaaatacaacatcgagaggaagtagactcatattaaaaatacttactaataataacgtgcctcatgaagtaa  
gatccgaaaggaattggaataaaactttcctgcatctcaagccaagggggaaacaccagaatcaagtgtccgcgtgattgaagac  
acccctcgtccaagaatgcaaagcacatccaataaaagagctggattataactcctctcttctctggtgggcccgtggggtgggag  
ctggggcgagaggtgccgttgccccgttgctttcctctgggaggg (SEQ ID NO: 55)

[0414] (2) 3' Untranslated Region:

tgaagtcaacatgcctgccccaaacaaatatgcaaaagggtcactaaagcagtagaaataatatgcattgtcagtgatgtccatgaaa  
caaagctgcaggctgtttaagaaaaataacacacataataaacatcacacacacagacagacacacacacacaacaattaacag  
tcttcaggcaaaacgtcgaatcagctatttactgccaaagggaatatcttttttacattattaagaaaaaaagattttattttaag  
acagtcccatcaaaactcctgtctttggaatccgaccactaattgccaagcaccgcttcgtgtggctccacctggatgttctgtgcct  
gtaaacatagattcgctttccatgttgttgccggatcaccatctgaagagcagacggatggaaaaaggacctgatcattggggaag  
ctggctttctggtgctggaggctggggagaaggtgttcattcacttgcatcttttgccctgggggctgtgatattaacagaggagg  
gttcctgtggggggaagtccatgcctccctggcctgaagaagagactcttgcatactcacatgatgcatacctggtgggagga  
aaagagttgggaacttcagatggacctagtaccactgagatttccacgccgaaggacagcgatgggaaaaatgcccttaaatcat  
aggaaaagtatttttaagctaccaattgtgccgagaaaagcatttttagcaattatacaatatcatccagtaccttaagccctgattgtg  
atattcatatattttggatacgaccccccaactcccaactggcctctgtctgagtaagaacagaatcctctggaacttgagggaagt  
aacatttcggtgactccgcacaggaaggctagagttaccagagcatcaggccgccacaagtgcctgcttttaggagaccgaagt  
ccgcagaacctgcctgtgtccagcttgaggcctggtcctggaactgagccggggccctcactggcctcctccagggatgatca  
acagggcagtggtgtccgaatgtctggaagctgatggagctcagaattccactgtcaagaaagagcagtagagggtgtggctg  
ggcctgtcaccctggggccctccaggtagggccttttcacgtggagcatgggagccacgaccttcttaagacatgtatcactgta  
gagggaagggaacagaggccctggggccctcctatcagaaggacatggtgaaggctgggaacgtgaggagaggcaatggccac  
ggcccatittggtgtagcacatggcacgttggtgtgtggccttgcccacctgtgagtttaagcaaggctttaaatgactttggag  
agggtcacaaatcctaaaaagaagcattgaagtgggtgtcatggattaattgacctgtctatggaattacatgtaaacattatctg  
tcactgtagtttggtttatttgaaaacctgacaaaaaaaaagttccaggtgtggaatatgggggttatctgtacatcctggggcattaaa  
aaaaaatcaatggtggggaactataaagaagtaacaaaagaagtgacatcttcagcaataaactaggaaatttttttctccagttt  
agaatcagccttgaacattgatggaataactctgtggcattattgcattatataccatttatctgtattaactttggaatgtactctgtcaa  
tgtttaatgctgtggttgatattcgaaagctgctttaaaaaatacatgcatctcagcgttttttggttttaattgtattagttatggcctata  
cactatttggagcaaaggtgatcgtttctgttgagattttatctcttgattcttcaaaagcattctgagaaggtgagataagccctgag  
tctcagctacctaagaaaaacctggatgtcactggccactgaggagctttgttcaaccaagtcattgtcattccacgtcaacagaatt  
gtttattgtacagttatatctgtgtcccttgacctgtttctgaaggttctcgtccctgggcaattccgcatthaattcatggattcag



gattacatgcatgtttggftaaacccatgagattcattcagttaaaaatccagatggcaaatgaccagcagattcaaatctatgggtggtt  
gacctttagagagtgctttacgtggcctgtttcaacacagaccacccagagccctcctgccctcctccgcgggggctttctcatgg  
ctgtccttcagggcttctcctgaaatgcagtgggtgcttacgctccaccaagaaagcaggaaacctgtggtatgaagccagacctccc  
ggcgggctcaggggaacagaatgatcagacctttgaatgattctaattttaagcaaatatttttatgaaaggtttacattgtcaaagt  
gatgaatatggaatatccaatcctgtgctgctatcctgccaaaatcatttaattggagtcagttgcagtatgctccacgtggttaagatcc  
tccaagctgctttagaagtaacaatgaagaacgtggacgcttttaataataagcctgtttgtcttctgtgtgttcaaacgggattcaca  
gagtatttgaaaaatgtatatataattaagaggtcacgggggctaattgctggctggctgccttttgcctgtgggggtttgtacctggtttta  
ataacagtaaatgtgccagcctcttgccccagaactgtacagtattgtggctgcacttgcttaagagtagttgatgttgcattttct  
tattgttaaaaacatgttagaagcaatgaatgtatataaaagcctcaactagtcatttttctcctctcttttttcatatatctaatattttg  
cagttgggcaacagagaacctatccctattttgtattgaagagggttcacatctgcactttaactgctctttatgaatgaaaaaacagtc  
ctctgtatgtactcctctttacactggccagggtcagagttaaatagagtatatgcactttccaaattggggacaagggtctataaaaaa  
gccccaaaaggagaagaacatctgagaacctcctcggccctcccagtcctcgcctgcacaaatactccgcaagagaggccagaat  
gacagctgacagggtctatggccatcgggtcgtctccgaagatttggcaggggcagaaaactctggcaggcttaagatttgaata  
aagtcacagaatcaaggaagcacctcaatttagttcaacaagacgccaacattctctccacagctcacttacctctctgtgttcagat  
gtggccttccatttatatgtgatctttgtttattagtaaatgcttatcatctaaagatgtagctctggccagtgaggaaaaattaggaagt  
attataaatcgagaggagttataataatcaagattaaatgtaataatcaggggcaatcccaacacatgtctagctttcacctccaggatc  
tattgagtgaacagaattgcaaatagtctctatttgaattgaacttatcctaaaacaaatagttataaatgtgaactaaactctaattaat  
tccaactgtacttttaaggcagtggtgttttagactttctatcattatagttagtaatgtacacctactctatcagagaaaaacaggaa  
aggctcgaaatacaagccatttcaaggaaattaggagagtcagttgaaattctattctgatcttattctgtggtgtcttttcagcccagac  
aaatgtgggttacacactttttaagaaatacaattctacattgtcaagcttatgaagggtccaatcagatctttattgttattcaatttggatctt  
cagggtatttttttaattattatgggacaaaaggacatttgttgagggggtgggaggagggaacaatttttaataaaaacattccca  
agtttggatcagggagttggaagttttagaataaccagaactaagggtatgaaggacctgtattggggctgatgtgatgcctctgcg  
aagaacctgtgtgacaaatgagaaacattttgaagttgtggtacgacctttagattccagagacatcagcatggctcaaagtgcagc  
tccgtttggcagtgcaatggtataaattcaagctggatgtctaatgggtatttaacaataaatgtgcagtttaactaacaggatattt  
aatgacaaccttctggttggttagggacatctgtttctaaatgtttattatgtacaatacagaaaaaattttataaaaattaagcaatgtgaa  
actgaattggagagtataatacaagtccttagtcttaccagtgatcattctgttccatgtctttggacaacctgaccttggacaat  
catgaaatatgcactcactggatgcaaagaaaaatcagatggagcatgaatggtactgtaccgggttcatctggactgccccagaaaa  
ataacttcaagcaaacatcctatcaacaacaaggtgttctgcataccaagctgagcacagaagatgggaacactggtggaggatg  
gaaaggctcgtcaatcaagaaaattctgagactattaataaataagactgtagtgtagatactgagtaaatccatgcacctaaaccttt  
tggaatatctgccgtgggcccctccagatagctcattcattaagttttccctccaagtgagaatttgaagagtacagtggttgcatt  
tcttttggggaagctttcttttgggtgtttgtttattataaccttcttaagttttcaaccaagggttgcctttgtttgagttactgggggtattttgt  
tttaataaaaaataagtgtaataaagtgttttgtattgaaagcttttgttatcaagattttcatacttttaccttccatggctcttttaagattg  
atacttttaagagggtgctgatattctgcaacactgtacacataaaaaatacggtaaggatactttacatggttaaggtaagtaagctc  
cagttggccaccattagctataatggcactttgtttgtgtgttgaaaaagtcacattgccattaaacttcttctgtctgtctagttaatatt  
gtgaagaaaaataaagtacagtgtgagatactg (SEQ ID NO: 56)

**6.23. Insulin-like Growth Factor Binding Protein-2**

[0415] See, *e.g.*, GenBank Accession No. X16302.

General Target Regions:

[0416] (1) 5' Untranslated Region:

attcggggcgagggaggaggaagaagcggaggaggcgggtcccgctcgcagggccgtgcacctgcccggccgcccgtcgc  
tcgctcggccgcccgcgccgcgtgccgaccgccagc (SEQ ID NO: 57)

[0417] (2) 3' Untranslated Region:

tgatccagggagccccaccatccggggggaccccgagtgatcatctcttctacaatgagcagcaggaggcttgcgggggtgcacac  
ccagcggatgcagtagaccgcagccagccgggtgcctggcgccccctgcccccgccctctccaaacaccggcagaaaacgga  
gagtgcctgggtgggtgggtgctggaggatttccagttctgacacacgtatttatatttgaaagagaccagcaccgagctcggcacc  
tccccggcctctctctccagctgcagatgccacacctgctccttctgcttccccgggggaggaagggggttggtcggggag  
ctggggtacaggttggggagggggaagagaaattttattttgaaccctgtgtccctttgcataagattaaaggaaggaagaaagt  
(SEQ ID NO: 58)

**6.24. K-ras Oncogene Protein**

[0418] See, *e.g.*, GenBank Accession No. M54968.

General Target Regions:

[0419] (1) 5' Untranslated Region:

tcctaggcggcggccgcggcggcggaggcagcagcggcggcggcagtgggcggcggcgaagggtggcggcggctcggccagt  
actcccgccccccgccatttcggactgggagcagcgcggcgcagggcactgaaggcggcggcggggccagaggctcagcgg  
ctcccagggtgcgggagagaggcctgctgaaa (SEQ ID NO: 59)

[0420] (2) 3' Untranslated Region:

taaatacaattgtactttttcttaaggcatactagtagcaagtggtaattttgtacattacactaaattattagcatttgttttagcattaccta  
atTTTTcctgctccatgcagactgtagctttaccttaaatgcttattttaaaatgacagtggaggttttttctcgaagtgcagttatc  
ccagagtttgggttttgaactagcaatgcctgtgaaaaagaaactgaatacctaagatttctgtcttgggggttttgggtgcagttga  
ttacttctatttttctaccaagtgtgaatgttggtgtgaacaaattaatgaagcttttgaatcatccctattctgtgtttatctagtcacata  
aatggattaattactaatttcagttgagaccttctaattgggttttactgaaacattgaggacacaaattatgggcttcctgatgatgattc  
ttctaggcatcatgtcctatagttgtcatccctgatgaatgtaaagttacactgttcacaaagggtttgtctcctttccactgctattagtc  
ggtcactctcccaaaatattatatttttctataaaaagaaaaaatggaaaaaattacaaggcaatggaaactattataaggccattt  
ccttttcacattagataaattactataaagactcctaataagcttttctgttaaggcagaccagttatgaatgggattattatagcaacat  
tttggggctatattacatgctactaaattttataataattgaaaagatttaacaagtataaaaaattctcataggaattaaatgtagtctc  
cctgtgtcagactgctcttcatagtataactttaaatcttttctcaacttgagctttgaagatagtttaattctgcttgtgacattaaaaga  
ttatttgggccagttatagcttattagggtgtgaagagaccaaggttgcaagccaggccctgtgtgaaccttgagctttcatagagagtt  
tcacagcatggactgtgtgccccacggtcatccgagtggtgtacgatgcattggttagtcaaaaatggggaggggactagggcagtt  
tggatagctcaacaagatacaatctcactctgtggtggtcctgctgacaaatcaagagcattgctttgtttcttaagaaaacaaactcttt  
tttaaaaattacttttaataattaactcaaaagttgagatttgggggtggtggtgtgccaaagacattaatttttttaacaatgaagtga

aaagttttacaatctctagggttggttagttctttaaactggtaaattaacattgcataaacacitttcaagctgatccatatttaataat  
gctttaaaataaaaaataaaacaatcctttgataaattaaaatgttacttattttaaaataaatgaagtgagatggcatggtaggtgaa  
agtatcactggactagggtgttggtgacttaggttctagatagggtgtcttttaggactctgattttgaggacatcacttactatccatttctc  
atgttaaaagaagtcactcacaactcttagtttttttttacactatgtgatttatattccatttacataaggatacacttatttgcagctca  
gcacaatctgtaaatttttaacctatgttacaccatcttcagtgccagtcttgggcaaaattgtgcaagaggtgaagtttatattgaatatac  
cattctcgttttaggactcttctccatattagtgtcatcttgccctccctacctccacatgccccatgacttgatgcagtttaatacttgtaa  
ttcccctaaccataagatttactgctgctgtggatatctccatgaagtttcccactgagtcacatcagaaatgccctacatcttatttccct  
caggggtcaagagaatctgacagataccataaaagggtttgacctaatacctaattttcaggtgggtggctgatgctttgaacatctctt  
gctgccccaatccattagcgacagtaggattttcaaccctggatgaatagacagaaccctatccagtgggaaggagaatttaataaag  
atagtgcagaaagaattccttaggtaatctataactaggactactcctggtaacagtaatacattccattgttttagtaaccagaaatcttc  
atgcaatgaaaaatactttaattcatgaagcttacttttttttgggtgtcagagctcgtctgtcaccaggtggaatgcagtggtg  
ccatctcagctcactgcaaccttccatcttcccagggtcaagcgattctcgtgccctggcctcctgagtagctgggattacaggcggtg  
gcactacactcaactaattttgtatttttaggagagacggggttcacctgttggccaggctgggtctgaactcctgacctcaagtatt  
caccaccttggcctcataaacctgtttgcagaactcattttcagcaaatatttattgagtgacctaccagatgccagtcaccgcaca  
aggcactgggtatatggtatcccaaacaagagacataatcccggctccttaggtactgctagtgtggtctgtaatatcttactaaggcc  
tttgggtatacgacccagagataacacgatgcgtatttttagtttgcaaagaaggggttgggtctctgtgccagctctataattgttttgcta  
cgattccactgaaactcttcgatcaagctactttatgtaaactcattcattgttttaaaggaataaacttgattatattgtttttatttggcata  
actgtgattcttttaggacaattactgtacacattaagggtgtatgtcagatattcataattgacccaaatgtgtaattccagtttctctgcat  
aagtaattaaataactttaaaataatagttttatctgggtacaaataaacagtgccctgaactagttcacagacaagggaacttctat  
gtaaaaatcactatgatttctgaattgctatgtgaaactacagatctttggaacactgttttaggtagggtgttaagacttgacacagtacct  
cgtttctacacagagaaagaaatggccatacttcaggaactgcagtgcttatgaggggatatttaggcctcttgaattttgatgtagat  
gggcattttttaaggtagtggttaattacctttatgtgaactttgaatggtttaacaaaagattgttttttagagatttttaagggggaga  
attctagaaataaatgttacctaattattacagcctaaagacaaaaatcctgttgaggttttttaaaaaaagactaaattacatagactta  
ggcattaacatgtttgtggaagaatatagcagacgtatattgtatcatttgagtgaatgttcccaagtaggcattctaggctctatttaact  
gagtcacactgcataggaatttagaacctaacttttatagggtatcaaaactgtgtcaccattgcacaattttgtcctaataatatacataga  
aacittgtggggcatgttaagttacagtttcacaaagttcatctcattgtattccattgatttttttttcttaaacatttttcttcaaaaca  
gtatatataacttttttaggggatttttttagacagcaaaaaactatctgaagatttccattgtcaaaaagtaatgatttcttgataattgtg  
tagtgaatgttttttagaaccagcagttaccttgaaagctgaatttatatttagtaacttctgtgttaatactggatagcatgaattctgcat  
tgagaaactgaatagctgtcataaaatgctttcttctaaagaaagatactcacatgagttctgaagaatagtcataactagattaaga  
tctgtgttttagtttaatagtttgaagtgccgttttgggataatgataggttaatttagatgaatttaggggaaaaaaaagtatctgcagttat  
gttgagggccatctctccccccacacccccacagagctaactgggttacagtgttttatccgaaagtttccaattcc (SEQ ID  
NO: 60)

**6.25. Target of Antiproliferative Antibody**

[0421] See, *e.g.*, GenBank Accession No. M33680 and Trifillis et al., 1999, RNA 5:1071-1082.

General Target Regions:

[0422] (1) 5' Untranslated Region:

ccattgtgct ggaaaggcgc gcaacggcgg cgacggcggc gacccaccg cgcacctgc caggcctccg  
cgcccagccg cccacgcgcc cccgcgcccc gcgccccgac cctttcttcg cgcgcccgcc cctcggccccg  
ccaggccccc ttgccggcca cccgccaggc cccgcgcccgg cccgccccgc gccaggacc ggcccgcgcc  
ccgcaggccg cccgcgcccc gcgcccgc (SEQ ID NO: 61)

[0423] (2) 3' Untranslated Region:

g ccccgccagc tctggccaca gggacctctg cagtgcctcc taagtaccc ggacacttc gagggggcca  
tcaccgcctg tgtatataac gttccggta ttactctgct acacgtagcc ttttacttt tggggtttg ttttgttct gaactttct  
gttaccttt cagggtgat gtcacatgta ggtggcgtgt atgagtggag acgggcctgg gtcttgggga ctggagggca  
ggggctcttc tgccctggg gtcccagggt gctctgctg ctcagccagg cctctcttg gagccactcg cccagagact  
cagcttgcc aacttggggg gctgtgtcca cccagccgc ccgtctgtg ggctgcacag ctcacctgt tcctcctgc  
cccgttga gagccgagtc tgtgggact ctctgccttc atgcacctgt ctttctaac acgtgcctt caactgta  
cacaacatcc tgactccgtc attaataaa gaaggaacat caggcatgct aaaaaaaaa aaaaa (SEQ ID NO: 62)

**6.26. Downstream Regulatory Element-antagonist Modulator**

[0424] See, *e.g.*, GenBank Accession No. AJ131730.

General Target Regions:

[0425] (1) 5' Untranslated Region:

gaattccggc aaacatgagg cagctgccag ccggcctggg cagtcttgc tgccctggct gtgaagtgg gaggtggca  
acagtttct tcagcgccca gg (SEQ ID NO: 63)

[0426] (2) 3' Untranslated Region:

gacacgt ccaaaggagt gcatggccac agccacctcc accccaaga aacctccatc ctgccaggag cagctccaa  
gaaacttta aaaaatagat ttgcaaaaag tgaacagatt gctacacaca cacacacaca cacacacaca  
gccattcatc tgggctggca gaggggacag agttcaggga ggggctgagt ctggctaggg gccgagtcca  
gaggccccag ccagcccttc ccaggccagc gaggcgaggc tgctctggg tgagtggctg acagagcagg  
tctgcaggcc accagctgct ggatgtcacc aagaaggggc tcgagtgcc tcaggaggg tccaatctc cgtccacc  
tcgtccgtt catccattct gctttctgc cacacagtgg ccggccagg ctcccctgg ctctccccg tagccactt  
ctgcccacta cctatgttc tagaaagccc ctcacctcag gacccagag gaccagctgg ggggcagggg  
ggagagggggg taatggaggc caagcctgca gcttttga aattctccc tgggggtccc agtatccct gctactccac  
tgacctggaa gagctgggta ccaggccacc cactgtgggg caagcctgag tggtagggg ccactggcat cattctccct  
ccatggcagg aaggcggggg attcaagtt tagggattgg gtcgtgggtg agaactgag ggcactctgc cagctccaca  
ggtggatgag cctctcttg cccagtcct ggtcagtgg gaatcgagt ggtggggctg tacacacct ccagcacaga

ctgttcctc caagtcctc ttaggtccc gggaggaacg tgggtcagag actggcagcc agggagcccg gggcagagct  
 cagaggagtc tgggaagggg cgtgtccctc ctcttctgt agtgccctc ccatggccca gcagcttggc tgagccctc  
 tctgaagca gctgtgcgcc gtccctctgc cttgcacaaa aagcacaaga cattccttag cagctcagcg cagccctagt  
 gggagcccg cacactgctt ctcggaggcc aggcctcct gctggctgag cttgggcccgt gtggcccaa tatggtggcc  
 ctggggaaga ggccttgggg gtctgctctg tgcctgggat cagtggggcc ccaaagccca gcccggtga ccaacattca  
 aaagcacaaa ccctggggac tctgcttggc tgtccctcc atctggggat ggagaatgca gcccaaagct ggagccaatg  
 gtgagggtg agagggtgt ggctgggtgg tcagcagaaa cccaggagg agagagatgc tgctccgcc  
 tgattggggc ctcaccaga aggaaccgg tccagccgc atggccctc caggaacatt cccacataat acattccatc  
 acagccagcc cagctccact cagggttggc ccggggagtc ccgtgtgcc ccaagaggct agccccaggg  
 tgagcagggc ctcagagga aaggcagtat ggcggaggcc atggggggcc ctcggcattc acacacagcc  
 tggcctccc tgcggagctg catggacgcc tggctccagg ctccaggctg actggggcct ctgctccag gagggcata  
 gcttccctg gctcaggat cttccctc cctcaccg ctgccagcc ctccagctg atgtactct gccttaagc  
 caaggcctca ggagagcatc accaccacac cctgcggcct tgccttgggg ccagactggc tgcacagccc  
 aaccaggagg ggtctgcctc ccacgtggg acacagaccg gccgcatgc tgcattggcag aagcgtctc cttgccagg  
 cctgggaggg tggctctgt tctcagcatc cactaatatt cagtctgta tatttaata aaataaact gacaaaggaa  
 aaaaaaccg (SEQ ID NO: 64)

#### 6.27. 6.27. Cox2

[0427] See, *e.g.*, GenBank Accession No. M90100.

General Target Regions:

[0428] (1) 5' Untranslated Region:

gtccaggaac tctcagcag cgctcctc agtccacag ccagacgcc tcagacagca aagcctacc ccgcgccgcg  
 ccctgcccgc cgctgcg (SEQ ID NO: 65)

[0429] (2) 3' Untranslated Region:

aagtctaa tgatcatatt tattattta tatgaacct gtctattaat ttaattattt aataatattt atattaaact ccttatgtta  
 cttacatct tctgaacag aagtcagtac tctgttgcg gagaaaggag tcatactgt gaagacttt atgtcactac  
 tctaaagatt ttgctgttc tgtaagtt ggaaaacagt tttattctg tttataaac cagagagaaa tgagtttga cgtctttta  
 cttgaattc aacttatatt ataaggacga aagtaaagat gttgaatac ttaaacacta tcacaagatg ccaaaatgct  
 gaaagtttt acactgtcga tgtttccat gcatttcca tgatgcatta gaagtaacta atgtttgaaa ttttaaagta  
 ctttgggta ttttctgtc atcaacaaa acaggtatca gtgcattatt aaatgaatat taaattaga cattaccagt aattcatgt  
 ctactttta aatcagcaa tgaaacaata attgaaatt tctaaattca taggtagaa tcacctgtaa aagctgttt  
 gatttctaa agttattaaa cttgtacata taccaaaaag aagctgtctt ggatttaaat ctgtaaaac agatgaaatt  
 ttactacaat tgcctgttaa aatatttat aagtatgtt ctttttcc caagagtata aacctttta gtgtgactgt taaaactcc  
 ttttaataa aatgccaaa tttattaagg tggaggagcc actgcagtgt tatctcaaaa taagaatc ctgttgat  
 attccagaat ctgtttatat ggctggtaac atgtaaaaac ccataacc cgccaaaagg ggtcctacc ttgaacataa  
 agcaataacc aaaggagaaa agcccaaatt attggtcca aattagggt taaacttt tgaagcaaac ttttttag

ccttgtgcac tgcagacctg gtactcagat ttgctatga ggtaaatgaa gtaccaagct gtgcttgaat aacgatatgt  
 ttctcagat ttctgtgt acagttaat ttagcagtcc atacacatt gcaaaagtag caatgacctc ataaaatacc  
 tcttcaaat gcttaaatc attcacaca ttaattttat ctgagtcttg aagccaattc agtaggtgca ttggaatcaa  
 gcctggctac ctgcatgctg ttcctttct ttcttcttt tagccatttt gctaagagac acagtcttct caaacacttc gtttctcta  
 tttgtttta ctagtttta gatcagagtt cactttcttt ggactctgcc tatattttct tacctgaact ttgcaagtt tcaggtaaa  
 cctcagctca ggactgctat ttagctctc ttaagaagat taaaaaaaa aaaaaa (SEQ ID NO: 66)

## 6.28. Her-2

[0430] General Target Regions:

[0431] (1) 5' Untranslated Region:

gcgcccggccccaccctcgcagcaccgcgccccgcgccctccagccgggtccagccggagccatggggccggagcc  
 gcagtgagcaccatggag (SEQ ID NO: 67)

[0432] (2) 3' Untranslated Region:

tgaaccagaaggccaagtccgcagaagccctgatgtgtcctcagggagcaggggaaggcctgacttctgctggcatcaagaggtg  
 ggagggccctccgaccacttccaggggaacctgccatgccaggaacctgtcctaaggaaccttcttctgcttgagtccagatg  
 gctggaaggggtccagcctcgttgaagaggaacagcactggggagtctttgttgattctgaggccctgccaatgagactctagg  
 gtccagtggatgccacagcccagcttggcccttcttccagatcctgggtactgaaagccttagggaagctggcctgagagggga  
 agcggccctaagggagtgtctaagaacaaaagcgacccattcagagactgtccctgaaacctagtactgcccccatgaggaagg  
 aacagcaatggtgtcagtatccaggctttgtacagagtgttttctgttttagttttactttttgtttgttttttaaagacgaaataaagac  
 ccaggggagaatgggtgttatggggaggcaagtgtggggggccttctccacaccacttgtccatttgcaatatattttggaaa  
 ac (SEQ ID NO: 68)

## 7. EXAMPLE: VASCULAR ENDOTHELIAL GROWTH FACTOR

### 7.1. Introduction

[0433] Vascular endothelial growth factor (VEGF) plays a key role in tumor angiogenesis. Considerable evidence demonstrates that VEGF is a viable target for tumor therapy (Carmeliet & Jain, 2000, Nature 407:249-257; Sepp-Lorenzino & Pan, 2000, Angiogenesis – Research frontiers. A basic science conference of the New York Academy of Medicine. Exp. Opin. Invest. Drugs 9:1-7; and Hichlin et al., 2001, DDT 6: 517-528). There are several ongoing clinical trials (phase I-phase III) indicating that either VEGF neutralizing antibodies or VEGFR2-mediated signal transduction inhibitors are effective for tumor therapy (Carmeliet & Jain, 2000, Nature 407:249-257 and Matter, 2001, Drug Discovery Today 6:1005-1024).

[0434] VEGF protein expression is tightly regulated at both the transcriptional and post-transcriptional levels. Under hypoxic conditions, tumor cells express high levels of VEGF that can promote angiogenesis and thus support the growth of tumor cells. Increase of

VEGF protein is due to both increased transcription and enhanced mRNA stability. Hypoxia-inducible factor 1 (HIF-1) is responsible for the transcriptional activation of the VEGF gene in hypoxic cells by binding to a hypoxia response element (HRE) located 1kb upstream of the transcription initiation site. In addition, the abundance of VEGF mRNA is increased due to stabilization of the mRNA by binding of HuR to the 3' UTR (untranslated region). Under hypoxic conditions, cap-dependent translation is replaced by cap-independent translation of the VEGF mRNA which is mediated by an internal ribosome entry site (IRES) within the VEGF 5'UTR.

[0435] This Example demonstrates the generation of stable cell lines, harboring VEGF 5' and 3'UTR sequences, which can be used to identify small molecular weight compounds that inhibit VEGF IRES-dependent translation or modulate VEGF mRNA stability.

## **7.2. Materials and Methods**

### **7.2.1. Generation of VEGF 5' and 3'UTRs**

[0436] The VEGF 5'UTR was generated using PCR from human genomic DNA. The full-length 5'UTR was prepared by the ligation of two separate PCR products (FIG. 1A). The first half of the 5'UTR (designated VEGF 5'UTR2, encompassing nucleotides 1 to 498) was amplified with primer 1 (5'-AAA GTC GAC GTA ATC GCG GAG GCT TGG GGC AGC CGG-3', SEQ ID NO: 69, and primer 2 (5' TTT GCG ACT GGT CAG CTG CGG GAT CCC AAG 3', SEQ ID NO: 70). The second half of the VEGF 5'UTR (designated VEGF 5'UTR1, from nucleotide 337 to 1038, plus the first 45 bp of the VEGF open reading frame) was amplified using primer 3 (5'-AA GTC GAC GTA AGA GCT CCA GAG AGA AGT CGA G-3, SEQ ID NO: 71 and primer 4 (5'-AAA CCC GGG CAG CAA GGC AAG GCT CCA ATG CAC-3', SEQ ID NO: 72). Each PCR product was digested with BamH I, and ligated together to produce the full length 5'UTR. To facilitate downstream cloning into dicistronic plasmid p2luc-i, primers 1 and 3 were designed to include a Sal I site and a stop code (TAA), immediately after the Sal I site at the 5' end, and primer 4 for VEGF 5'UTR1 includes a Xma I site at the 5' end (FIG. 1C).

[0437] The entire VEGF 3' UTR (shown in FIG. 1B) was amplified by genomic PCR using primer 5 (5'-GCC GGG CAG GAG GAA GGA GCC TCC CTC AGG GTT TCG GGA 3', SEQ ID NO: 73) and primer 6 (5'-CTG CAC TAG AGA CAA AGA CG T GAT GTT AAT -3', SEQ ID NO: 74). The Bgl II and EcoR I restriction sites were used for subsequent cloning.

### 7.2.2. Plasmid Construction

[0438] Each PCR fragment (VEGF 5'UTR1, VEGF 5'UTR2 and VEGF 3'UTR) was cloned into pT-Adv vector for confirmation by DNA sequencing using the Clontech advantage cloning kit. A SalI-XmaI VEGF 5'UTR1 fragment was subcloned into the p2luc-i dicistronic plasmid (FIG. 2A, Grentzmann et al., 1998, RNA 4:479-486). The sequence of the polylinker site is GAA CAA ATG TCG ACG GGG GCC CCT AGC AGA TCT AGC GCT GGA TCC CCC GGG GAG CTC AUG GAA GAC (SEQ ID NO: 75, FIG. 2A). The resulting plasmid (designated p2luc/VEGF5UTR1, see FIG. 2B) contains VEGF 5'UTR1 between the two reporter genes (renilla luciferase and firefly luciferase) with a stop code (TAA) immediately after the Sal I site and a fusion translation junction between the first 15 AA of VEGF and firefly luciferase open reading frame. To construct the dicistronic plasmid containing the full length VEGF 5'UTR, VEGF 5'UTR2 was then subcloned into p2luc/VEGF5UTR1 between SalI and BamHI (designated p2luc/VEGF5UTR-fl; FIG. 2B). This plasmid also has a stop code (TAA) immediately after the Sal I site to prevent read-through from the first reporter to the second.

[0439] To map the region of the IRES essential for activity, dicistronic plasmids containing various deletions within the VEGF 5'UTR were prepared (FIG. 2B). Plasmid p2luc/veg5'utr-delta51-476 is derived from p2luc/veg5'utr-fl by removing the Nhe I fragment (nt51 to 746); plasmid p2luc/veg5utr-delta476-1038 was derived from p2luc/veg5utr-fl by removing the sequence from BamH I site to the 3'end of 5'UTR; plasmid p2luc/veg5utr-delta1-476 was derived from p2luc/veg5utr-fl by removing the sequence from BamH I to the 5'end of 5'UTR.

[0440] To generate stable cell lines for high throughput screening, a monocistronic reporter plasmid (pluc/VEGF5'+3'UTR) containing the VEGF 5' and 3'UTRs and firefly luciferase gene (FIG. 3A) was constructed. Briefly, a Sal I-Not I fragment, containing the full length VEGF 5'UTR and firefly reporter gene, from p2luc/VEGF5'UTR-fl was subcloned into pCDNA5/TO between EcoR V and Not I, and then the VEGF 3'UTR was subcloned into the intermediate plasmid at the Not I site by blunt-end ligation.

### 7.2.3. DNA Transfection and Generation of Stable Cell Lines

[0441] 293T cells were transfected with pluc/VEGF5'+3'UTR using the Fugene 6 transfection reagent (Roche) according to manufacture's instruction. 48 hours after transfection, the cells were lysed and plasmid function was monitored by measuring luciferase activity using Promega's luciferase kit according to manufacture's instruction.

[0442] To generate stable cell lines, plasmid pluc/VEGF5'+3'UTR was transfected into 293T cells as described. 48 hours after transfection, the cells were trypsinized, resuspended



in culture media plus 200 mg/ml hygromycin B, then seeded in 96 well plates at 100 to 500 cells per well for selection. The media containing hygromycin B was changed every 3 to 4 days. After 10 to 14 days of selection, hygromycin resistant clones were screened under a microscope and wells harboring a single colony were expanded under hygromycin selection for further experiments.

#### **7.2.4. Luciferase Assay**

[0443] F. luciferase and R. luciferase activities were measured using the Luciferase reporter assay system (Promega) according to manufacturer's instruction.

#### **7.2.5. Semi-quantitative PCR**

[0444] DNA and RNA were isolated from B9 cells using TRIzol reagent (GIBCO BRL) according to the manufacturer's instructions. cDNA was synthesized using Promega's reverse transcription system. Semi-quantitative PCR was performed with gene specific primers for firefly luciferase or glyceraldehyde phosphodehydrogenase (GAPDH) as an internal control. The primer pairs for firefly luciferase amplification were as follows: 5'-CGG TGT TGG GCG CGT TAT TTA TCG GAG TTG-3' (SEQ ID NO: 76) and 5'-TTG GCG AAG AAT GAA AAT AGG GTT GGT ACT-3' (SEQ ID NO: 77); the primer pairs for GAPDH were as follows: 5'-GGT GAA GGT CGG AGT CAA CGG A-3' (SEQ ID NO: 78) and 5'-GAG GGA TCT CGC TCC TGG AAG A-3' (SEQ ID NO: 79). The PCR products were separated on 1% agarose gel, stained with ethidium bromide and quantified on UVP with Labworks software.

#### **7.2.6. High Throughput Screening**

[0445] High throughput screening ("HTS") for compounds that inhibit untranslated region-dependent expression of vascular endothelial growth factor ("VEGF") is accomplished using stable cell lines described in Section 7.2.3. The 293T cell line contained stably integrated copies of the firefly luciferase gene flanked by both the 5' and 3' UTRs of VEGF. Cell lines exhibiting consistently high levels of firefly luciferase expression are further expanded and optimized for HTS.

[0446] Screening of compounds is done using one hundred 384-well plates per day. Each 384-well plate contains a standard puromycin titration curve that is used as a reference to calculate % inhibition and the statistical significance of the data points generated in the assay. This curve occurs in wells from column 3 and 4 of the 384-well plate. The concentration of puromycin is 20 mM serially diluted 2-fold to 0.078 mM plated in quadruplicate. Columns 1 and 2 contain 16 standards each of a positive control 0.5% DMSO and a negative control consisting of the puromycin at 20 mM. The difference

between the two controls is used as the window to calculate the percentage of inhibition of luciferase expression in the presence of a compound. Columns 5 through 24 contain compounds from a library of small molecules.

[0447] Two confluent T175 flasks of the VEGF stable cell line described above (B9) are split into twenty T175 flasks three days prior to screening. On each day of the HTS assay, the cells are dislodged from the flask with 3 ml of 0.25 % trypsin-EDTA (Gibco, cat no. 25200-056) and diluted to 10 ml with non-selective media. This is repeated for all twenty flasks and the cells are combined, counted and diluted to a concentration of 263.15 cells/ml. 38 to 39 ml are then added to each well containing 1 to 2 ml of compound from a small molecule library to a final compound concentration of 7.5 mM (3.75 mg/ml) in 0.5 % DMSO. The puromycin standard curve also contains 0.5 % DMSO. The stable cell line is incubated in the presence of compound overnight (approximately 16 hours) at 37 C in 5 % CO<sub>2</sub>. To monitor firefly luciferase activity, LucLite Plus (Packard cat no. 6016969) is prepared according to manufactures' instructions and 20 ml is added to each well. Following a brief incubation at room temperature (minimum 2 min.), firefly luciferase activity in each well is detected with the ViewLux 1430 ultraHTS Microplate Imager (Perkin Elmer). All data obtained is uploaded into Activity Base for % inhibition calculations and statistical analyses.

### 7.3. Results

[0448] The ability of VEGF 5'UTR sequences to modulate internal translation initiation was tested using the plasmid vector that encodes a dicistronic mRNA (FIG. 2A). The renilla luciferase is translated from the first cistron by a cap-dependent scanning mechanism, while the firefly luciferase in the second cistron is translated only if preceded by an internal ribosome entry site. In this study, five discistronic plasmids containing various deletions of the VEGF 5'UTR (FIG. 2B) were generated and transiently transfected into 293T cells to monitor IRES-dependent translation of firefly luciferase. 48 hours after transfection, extracts were prepared and assayed for renilla and firefly luciferase activities using the dual luciferase kit from Promega. As shown in FIG. 2C, deletion of either the first 336 or the first 476 nucleotides has no significant effect on firefly luciferase activity compared to full length VEGF5'UTR directed luciferase levels. However, deletion between nucleotides 51 and 746 decreased firefly luciferase activity more than 75% (33.68 $\pm$ 4.91 vs 161 $\pm$ 30.49). Deletion of nucleotides 476 to the 3' end of the VEGF 5'UTR decreased firefly luciferase activity more than 90% (12.15 $\pm$ 1.2 v.s. 161 $\pm$ 30.49). Taken together, these results confirm that the VEGF 5'UTR harbors IRES activity, and also

indicates that the region of the VEGF IRES essential for function is located within nucleotide 476 to the 3' end of VEGF 5'UTR.

[0449] To generate stable cell lines for High Throughput Screening ("HTS"), a monocistronic reporter plasmid under the transcriptional control of the CMV promoter (pluc/veg5'+3'UTR; FIG. 3A) was constructed. This plasmid contains both the VEGF 5'- and 3'-UTRs separated by the firefly luciferase gene. After confirmation of luciferase production by transient transfection (data not shown), transfected 293-T cells were seeded in 96 well plates at a concentration of 100-500 cells per well, and then cultured under hygromycin B selection. After two weeks of selection, 19 clones were screened for luciferase activity, three of which demonstrated high levels of luciferase activities (clones B9, D3, H6; FIG. 3B). To determine which cell line demonstrated the highest level of expression, the luciferase activities of clones B9, D3, H6 were compared and normalized against the protein concentrations extracted from each cell line. The results shown in FIG. 4 demonstrate that the luciferase activity from B9 cells was two fold greater than H6 cells, and more than three fold higher than D3 cells.

[0450] To determine if the B9 cells are stable, these cells were maintained under hygromycin selection for more than three months, with intermittent monitoring of luciferase activity. The results indicate that this cell line is stable and sustains a high level of luciferase expression when continuously cultured in vitro for more than three months (FIG. 5). Sustained expression of luciferase by B9 cells indicated that the monocistronic plasmid integrated into the genomic DNA. Semi-quantitative PCR was performed to determine the number of copies of the reporter plasmid integrated per B9 cell. As FIG. 6A shows, series diluted plasmid pluc5'+3'vegf-UTR were included as positive control to make sure the reaction for sample (genomic DNA from B9 cells) was within the linear range, i.e., not saturated. The PCR standard curve was plotted with the PCR product intensity against the amount of positive plasmid control loaded for PCR (FIG. 6B). Sigma plot regression indicated that PCR product intensity for B9 genomic DNA (50 ng) is about the same level of 6.4 pg plasmid control. As 1 mg of 8 kb plasmid roughly contains  $10^{11}$  copies and  $10^6$  cells have 10 mg genomic DNA, the results here indicated that approximately 100 copies of the plasmid were integrated per cell.

[0451] High throughput screening ("HTS") for compounds that inhibit untranslated region-dependent expression of vascular endothelial growth factor ("VEGF") is accomplished with the generated stable cell lines.

## 8. EXAMPLE: SURVIVIN

### 8.1. Introduction

[0452] Survivin, a member of IAP (inhibitor of apoptosis proteins) gene family, is critically required for suppression of apoptosis and ensuring cell division through the G2/M phase of the cell cycle. It is absent in normal adult tissues, but highly expressed in all of the most common cancers in a cell cycle-regulated manner. Disruption of survivin expression/function by antisense or dominant-negative mutation resulted in deregulation of mitotic progression and spontaneous apoptosis. It has been demonstrated that survivin targeting in vivo increased apoptosis and reduced proliferation of tumor cells, but did not affect cell viability of proliferating normal cells. It has also been showed that survivin targeting induces apoptosis and sensitizes tumor cells to chemical agents. Therefore, inhibition of survivin may be of great benefit for refractory cancer therapy when combined with standard chemotherapy. Another benefit for this project will be low toxicity because survivin expression is absent in normal adult tissues. Taken together, these indicated survivin is valid target for cancer therapy.

[0453] Translation of survivin might be cap-independent/IRES dependent since it is maximally expressed in metaphase. In G2/M phase, the eIF4E-binding proteins (4E-BPs) become hypophosphorylated. 4E-BPs compete with eIF4G for eIF4E binding, thus preventing eIF4F formation and cap-dependent translation initiation. In addition, survivin mRNA has a long 3'UTR featuring a poly(U) sequence and multiple CU repeats.

[0454] This Example demonstrates the generation of stable cell lines, harboring survivin 5' and 3'UTR sequences, to identify small molecular weight compounds that inhibit survivin IRES-dependent translation or modulate survivin mRNA stability.

### 8.2. Materials and Methods

#### 8.2.1. Generation of Survivin 5' and 3' UTRs

[0455] The 5' UTR of survivin was generated by filling-in partially overlapping oligonucleotides with Taq polymerase. A 5' UTR forward oligonucleotide (5' AAAGTCGACGTAACCGCCAGATTTGAATCGCGGGACCCGTTGGCAGAGGTGGC GG 3', SEQ ID NO: 80) encompassing nucleotides 1 to 42 of the 5' UTR of survivin and a 5' UTR reverse oligonucleotide (5' AAAGGATCCGGGCAACGTCGGGGCACCCATGCCGCGCCGCCACCTCTGCCAA C 3', SEQ ID NO: 81) encompassing nucleotides 26 to 49 of the 5' UTR of survivin as well as the first 21 nucleotides of the open reading frame of survivin were annealed at 45 C and

extended at 72 C with Taq polymerase. The Sal I and BamH I restriction sites (underlined) were used for subsequent cloning.

[0456] The 3' UTR of survivin was amplified from human genomic DNA using the 3' UTR forward oligonucleotide (5'

AAAGCGGCCGCGGCCTCTGCCGGAGCTGCCTGGTCCCAGA 3', SEQ ID NO: 82)

and the 3' UTR reverse oligonucleotide (5'

AAATCTAGACTCAGGAACAGCCGAGATGACCTCCAGA 3', SEQ ID NO: 83). The

Not I and Xba I restriction sites were used for subsequent cloning.

### 8.2.2. Plasmid Construction

[0457] The survivin 3' UTR PCR product generated in Section 8.2.1 was cloned into the pT-Adv for sequence verification. A positive clone was subsequently digested with Not I and Xba I and the resulting 1.1 kb survivin 3' UTR PCR fragment was subcloned into pcDNA3.1/Hygro (Invitrogen cat. no. V87020) to generate the intermediate plasmid Surv3'UTR/pcDNA3.1/Hygro.

[0458] The survivin 5' UTR DNA fragment generated in Section 8.2.1 was digested with Sal I and BamH I and was subcloned into p2luci (see, *e.g.*, Grentzmann et al., 1998, RNA 4:479-486) to generate the intermediate plasmid, Surv5'UTR/p2luci, which contains the 5' UTR of survivin between the open reading frames of the renilla and firefly luciferase reporter genes. Surv5'UTR/p2luci was then digested with either Sal I and Not I or BamH I and Not I to isolate and gel purify the 1.75 kb survivin 5' UTR-firefly luciferase or the 1.7 kb firefly luciferase DNA fragments. The Sal I 5' overhang of the 1.75 kb survivin 5' UTR-firefly luciferase fragment was filled-in with T4 DNA polymerase and was subcloned into both Surv3'UTR/pcDNA3.1/Hygro and pcDNA3.1/Hygro digested with EcoR V and Not I to generate the plasmids, Surv5'UTR-Fluc-Surv3'UTR/pcDNA3.1/Hygro and Surv5'UTR-Fluc/pcDNA3.1/Hygro. The former plasmid contains the firefly luciferase reporter gene surrounded by both the 5' and 3' untranslated regions of survivin while the latter plasmid contains the firefly luciferase reporter gene preceded only by the 5' UTR of survivin and will be used as a "5' UTR-only" control plasmid in future experiments. The 1.7 kb BamH I-Not I firefly luciferase fragment was subcloned into both Surv3'UTR/pcDNA3.1/Hygro and pcDNA3.1/Hygro to generate the plasmids, Fluc-Surv3'UTR/pcDNA3.1/Hygro and Fluc/pcDNA3.1/Hygro. The former plasmid contains the firefly luciferase reporter gene followed only by the 3' UTR of survivin and will be used as a "3' UTR-only" control plasmid in future experiments. The latter plasmid contains only the firefly luciferase reporter gene lacking any surrounding untranslated regions of survivin and will be used in subsequent studies as a "no UTR" control plasmid.

[0459] Since the 5' UTR of survivin is small (49 nucleotides), it is likely that cap-dependent and cap-independent firefly luciferase expression in the survivin expression plasmids described above cannot be distinguished. One method of separating cap-dependent from cap-independent translation is through the introduction of a stable hairpin or secondary structure upstream or near the 5' end of the 5' UTR of the expression vector (see, *e.g.*, Muhlrads et al., 1995 Mol. Cell. Biol. 15:2145-2156). Therefore, two complementary oligonucleotides, SL top (5' CTAGAAGCTTAGGGCCGCGGATCCGCGCGCGGTTCGCCGCGCGCGGATCCGCG GTAGCAAGTTAGTC 3', SEQ ID NO: 84) and SL bottom (5' GACTAAGCTTGCTACCGCGGATCCGCGCGCGGCGAACC GCGCGCGGATCCGCG GCCCTAAGCTTCTAG 3', SEQ ID NO: 85) were synthesized, digested with Hind III, annealed and subcloned into the survivin expression vectors described above. A stable stem-loop structure with an 18 base-pair stem and a UUCG loop sequence will form and effectively block cap-dependent translation (see, *e.g.*, Beelman & Parker, 1994 J. Biol. Chem. 269:9687-9692 and Muhlrads et al., 1995 Mol. Cell. Biol. 15:2145-2156).

### 8.2.3. DNA Transfection and Stable Cell Line Generation

[0460] 293T cells were transiently transfected with equal amounts of each of the survivin expression vectors described in Section 8.2.2 (both with and without each of the 5' and 3' UTRs of survivin and in the presence and absence of the stem-loop secondary structure) using the Fugene 6 transfection reagent (Roche) according to manufacture's instruction. Untranslated region-dependent firefly luciferase activity was monitored forty-eight hours post-transient transfection according to manufacture's instruction (see, *e.g.*, Section 8.2.4.).

[0461] To generate stable cell lines, 293T cells were transiently transfected as above. Instead of lysing the transiently transfected 293T cells to monitor firefly luciferase activity, the cells were trypsinized, counted and seeded (10 ml) in 10 cm petri dishes at a concentration of 5000 cells/ml. The following day, hygromycin B was added in culture media to a final concentration of 200 mg/ml to select for cells in which the transiently transfected plasmid has stably integrated into the genome. Following ten to fourteen days of hygromycin B selection, individual hygromycin-resistant clones were expanded by transferring the cells from the petri dish to a single well in a six or twenty-four well plate using trypsin-soaked filter discs according to manufacture's instructions. Individual cell lines are then selected for further studies based on firefly luciferase expression levels.

#### **8.2.4. Luciferase Assays**

[0462] Firefly luciferase activity was measured with the luciferase reporter assay system (Promega) according to manufacture's instructions.

#### **8.2.5. High Throughput Screening**

[0463] High throughput screening ("HTS") for compounds that inhibit untranslated region-dependent expression of survivin is accomplished with stable cell lines described in Section 8.2.3. The 293T cell line contained stably integrated copies of the firefly luciferase gene flanked by both the 5' and 3' UTRs of survivin. Cell lines exhibiting consistently high levels of firefly luciferase expression are further expanded and optimized for HTS.

[0464] Screening of compounds is done using one hundred 384-well plates per day. Each 384-well plate contains a standard puromycin titration curve that is used as a reference to calculate % inhibition and the statistical significance of the data points generated in the assay. This curve occurs in wells from column 3 and 4 of the 384-well plate. The concentration of puromycin is 20 mM serially diluted 2-fold to 0.078 mM plated in quadruplicate. Columns 1 and 2 contain 16 standards each of a positive control 0.5% DMSO and a negative control consisting of the puromycin at 20 mM. The difference between the two controls is used as the window to calculate the percentage of inhibition of luciferase expression in the presence of a compound. Columns 5 through 24 contain compounds from a library of small molecules.

[0465] Two confluent T175 flasks of the survivin stable cell line described above are split into twenty T175 flasks three days prior to screening. On each day of the HTS assay, the cells are dislodged from the flask with 3 ml of 0.25 % trypsin-EDTA (Gibco, cat no. 25200-056) and diluted to 10 ml with non-selective media. This is repeated for all twenty flasks and the cells are combined, counted and diluted to a concentration of 263.15 cells/ml. 38 ml are then added to each well containing 2 ml of compound from a small molecule library to a final compound concentration of 7.5 mM (3.75 mg/ml) in 0.5 % DMSO. The puromycin standard curve also contains 0.5 % DMSO. The stable cell line is incubated in the presence of compound overnight (approximately 16 hours) at 37 C in 5 % CO<sub>2</sub>. To monitor firefly luciferase activity, LucLite Plus (Packard cat no. 6016969) is prepared according to manufactures' instructions and 20 ml is added to each well. Following a brief incubation at room temperature (minimum 2 min.), firefly luciferase activity in each well is detected with the ViewLux 1430 ultraHTS Microplate Imager (Perkin Elmer). All data obtained is uploaded into Activity Base for % inhibition calculations and statistical analyses.

### 8.5. Results

[0466] To determine the effect of the survivin untranslated regions on post-transcriptional control of gene expression, transient transfections of the survivin expression vectors described in Section 8.1.2., containing both, one or none of the 5' and 3' UTRs of survivin both in the absence or presence of the stem-loop secondary structure were performed. In the absence of the stem-loop secondary structure, cap-dependent and cap-independent translation are equally favored and no significant difference in firefly luciferase expression could be detected when either or both of the 5' and 3' UTRs are present or absent (FIG. 7A). This results confirms the earlier notion that, in the survivin expression vectors without the stem-loop secondary structure, the 5' UTR of survivin is unable to block cap-dependent translation. In the presence of the stem-loop secondary structure, a 3-fold increase in firefly luciferase expression can be detected only in the survivin expression vectors that contain the 5' UTR of survivin (FIG. 7B). This result strongly suggests that the 5' UTR of survivin can function as an internal ribosome entry site and promote cap-independent translation and helps explain the increase in the endogenous levels of survivin in the G2/M phase of the cell cycle when overall translation is dramatically reduced.

[0467] High throughput screening ("HTS") for compounds that inhibit untranslated region-dependent expression of survivin is accomplished with the generated stable cell lines.

## 9. EXAMPLE: HER-2

[0468] This Example demonstrates the generation of stable cell lines, harboring Her-2 5' and 3'UTR sequences, to identify small molecular weight compounds that inhibit Her-2 5' UTR-dependent translation or modulate Her-2 mRNA stability.

### 9.1. Her-2 Constructs

#### 9.1.1. Generation of Her-2 in vitro Expression Constructs

[0469] The 99 nucleotide 5' UTR of Her-2 was PCR-amplified from a human genomic DNA (Promega) using the following primers: Sense/HindIII: CAAGAAGCTTgcgccggccccccaccctcg (SEQ ID NO: 86) and Antisense/NcoI: AGCCCATGGtgcctcactgcggctccggccc (SEQ ID NO: 87). The Advantage-GC2-PCR kit was used according to the manufacturer's instructions (Clontech) with the following conditions: PCR cycle conditions were 94 C, 3 minutes, followed by 35 cycles of 94 C, 30 seconds, and 68 C, 30 seconds. The PCR-amplified product was cloned using the pT Adv kit (Clontech) according to the manufacturer's instructions. All clones were confirmed by



sequencing. The resulting clone was digested with HindIII/NcoI and the fragment was cloned into pT7Luc, upstream of the luciferase gene, to generate pT7Luc/5'UTR.

[0470] The 615 nucleotide 3'UTR was PCR-amplified from human genomic DNA (Promega) using the following primers: sense/BglII: agactctgaaccagaaggccaa (SEQ ID NO: 88) and antisense/KpnI: ctcggtaccagttttccaaaatatatttgcaaatgg (SEQ ID NO: 89). The Titanium Taq kit (Clontech) was used according to the manufacturer's instructions with the following amplification conditions: 94 C, 1 minute, followed by 35 cycles at 94 C, 30 seconds to denature, 60 C 30 seconds to anneal, 72 C 1 minute to extend. The product was gel purified and cloned using pT Adv (Clontech) according to the manufacturer's instructions. All clones were sequenced. The resulting clone was digested with BglII/KpnI and cloned into a BglII/KpnI digested pT7Luc and pT7Luc/5'UTR to generate pT7Luc/3'UTR and pT7Luc/5'and3'UTR, respectively.

### **9.1.2. Generation of Her-2 in vivo Expression Constructs**

[0471] Constructs for cell-based expression were generated by isolation of Her-2 containing fragments of pT7Luc/5'UTR, pT7Luc/3'UTR and pT7Luc/5' and 3'UTR digested with HindIII and KpnI and cloned into pcDNA (+) (Invitrogen).

### **9.1.3. Generation of Her-2 uORF Mutants**

[0472] The uORF contained within the Her-2 5' UTR was removed by extending the overlapping long primers. The overlapping sequence is underlined. The sense minus uORF HindIII primer is: cccaagcttcgcgcccggccccccaccctcgcagcaccgcgccccgcgcctccc (SEQ ID NO: 90) and the antisense minus uORF NcoI primer is: ggcccatggctccggctggaccggctgggaccggctgggagggcgcgaggcgcg (SEQ ID NO: 91). The primers (10 micrograms) were denatured at 95 C for 2 minutes, annealed at 60 C for 5 minutes and extended at 72 C for 10 minutes using Taq polymerase (Clontech). After buffer-exchange, the product was digested with NcoI and HindIII and cloned in the HindIII/NcoI sites of the in vitro expression vector pT7Luc and pT7Luc/3'UTR, yielding pT7Luc/5'UTR minus uORF and pT7Luc/5'UTR minus uORF and 3' UTR. Both plasmids were digested with HindIII and KpnI and the Her-2 containing fragment was subcloned into the HindIII/KpnI site of pcDNA (+) (Invitrogen) for cell-based studies.

## **9.2. Stable Cell Line Production**

[0473] Stable cell lines were generated in HeLa, 293T, and MCF-7. First, transient transfection was carried out using the Fugene 6 transfection reagent (Roche) according to manufacturer's instructions. Untranslated region-dependent firefly luciferase activity was

monitored forty-eight hours post-transient transfection with the luciferase reporter assay system (Promega) according to manufacture's instructions.

[0474] Next, stable cell lines were generated by first transiently transfecting the above cell lines. Instead of lysing the transiently transfected 293T cells (or other), the cells were trypsinized, counted and seeded (10 ml) in 10 cm petri dishes at a concentration of 5000 cells/ml. The following day, hygromycin B was added in culture media to a final concentration of 100 mg/ml for 293T cells and 200 mg/ml for MCF-7 and Hela, to select for cells in which the transiently- transfected plasmid has stably integrated into the genome. Following ten to fourteen days of hygromycin B selection, individual hygromycin-resistant clones were expanded by transferring the cells from the petri dish to a single well in a six or twenty-four well plate using trypsin-soaked filter discs according to manufacture's instructions. Individual cell lines are then selected for further studies based on firefly luciferase expression levels.

### 9.3. *In vitro* High Throughput Screen

[0475] Construct pT7Luc/5' and 3'UTR is utilized as a template for large-scale T7 polymerase transcription according to the manufacturer's protocol (Ambion). The mRNA template containing the Her-2 5' and 3' UTR and the Luciferase ORF is uncapped and used at 100 nanograms/reaction for a typical in vitro HTS. The number of samples run determines the amount of transcription yield that must be obtained. For example, for 100,000 reactions, using 100 nanograms of RNA/reaction, 10 milligrams of RNA must be produced. Typical yields from the Ambion T7 Transcription Kit for this template are 5 mg/ml of transcription.

[0476] Screening of compounds is done using one hundred 384-well plates per day. Each 384-well plate contains a standard puromycin titration curve that is used as a reference to calculate % inhibition and the statistical significance of the data points generated in the assay. This curve occurs in wells from column 3 and 4 of the 384-well plate. The concentration of puromycin is 20 mM serially diluted 2-fold to 0.078 mM plated in quadruplicate. Columns 1 and 2 contain 16 standards each of a positive control of 4% DMSO and a negative control consisting of the puromycin at 20 mM. The difference between the two controls is used as the window to calculate the percentage of inhibition of luciferase expression in the presence of a compound. Columns 5 through 24 contain compounds from a library of small molecules.

[0477] The *in vitro* translation reaction of the Her-2 driven Luciferase ORF consists of four microliters of rabbit reticulocyte lysate (Green Hectares) supplemented with 0.013 mgs/ml hemin (Sigma), 0.05 mgs/ml creatine kinase (Roche), and 0.125 mgs/ml tRNA

(Sigma Type XII, rabbit liver), 100 nanograms uncapped mRNA and buffer containing 100 mM KOAc, 0.5 mM Mg(OAc)<sub>2</sub>, 10 mM creatine phosphate, 0.03 mM amino acid mix, in a reaction volume of 20 ml. The reaction is then incubated at 30 C for 45 minutes. At the end of incubation, 20 mL of LucLite (Packard) is added to the reaction and the light output resulting from luciferase catalyzed conversion of luciferin, is monitored on a ViewLux uHTS Plate reader (Perkin Elmer).

#### 10. **EXAMPLE: CELL EXPRESSION VECTORS**

[0478] Stable cell line expression vectors (pCMR1 and pCMR2) are shown in FIGS. 8A and 8C. pCMR1, a high-level stable and transient mammalian expression vector designed to randomly integrate into the genome and pCMR2 is an episomal mammalian expression vector. pMCP1 (FIG. 8B) is a high level stable and transient mammalian expression vector designed to site-specifically integrate into the genome of cells genetically engineered to contain the FRT site-specific recombination site via the Flp recombinase (see, *e.g.*, Craig, 1988, Ann. Rev. Genet. 22: 77-105; and Sauer, 1994, Curr. Opin. Biotechnol. 5: 521-527). The nucleotide sequences are presented below.

##### [0479] pCMR1 (SEQ ID NO: 92)

gacggatcgggagatctcccgatccctatggtgactctcagtacaatctgctctgatgccgcatagttaagccagtatctgctccct  
gcttgtgtgttgaggctcgtgagtagtgcgcgagcaaaatttaagctacaacaaggcaaggcttgaccgacaattgcatgaagaat  
ctgcttaggggttaggcgttttgcgctgcttcgcgatgtacgggcccagatatacgcgttgacattgattattgactagttattaatagtaatc  
aattacgggggtcattagttcatagcccatatatggagttccgcgttacataacttacggtaaatggcccgcctggctgaccgcccac  
gacccccgcccattgacgtcaataatgacgtatgttcccatagtaacgccaatagggactttccattgacgtcaatgggtggagtattt  
acggtaaaactgcccacttggcagtagcatcaagtgtatcatatgccaaagtacgccccctattgacgtcaatgacggtaaatggccgc  
ctggcattatgccagtagcatgacctatgggactttcctacttggcagtagcatctacgtattagtcacgcctattaccatgggtgatgcgg  
tttggcagtagcatcaatgggcgtggatagcgggttgactcacggggatttccaaagtctccaccctattgacgtcaatgggagtttgtt  
tggcaccaaaatcaacgggactttccaaaatgtcgtacaactccgccccattgacgcaaattggcggttaggcgtgtacgggtggga  
ggctctatataagcagagctctctggctaactaagctttcggcgcgccgaggtaccatgggatccgaagacgccaaaaacataaga  
aaggccccggcgccattctatcctctagaggatggaaccgctggagagcaactgcataaggctatgaagagatacgcctgtgtcct  
ggaacaattgcttttacagatgcacatcgcaggtgaacatcacgtacgcggaatacttcgaaatgtccgttcggttggcagaagctat  
gaaacgatatgggctgaatacaaatcacagaatcgtcgtatgcagtgaactctcttcaattctttatgccggtgttggcgcgcttatt  
atcggagttgcagttgcgcccgcgaacgacattataatgaacgtgaattgctcaacagtatgaacatttgcagcctaccgtagtgtt  
tgtttccaaaagggggttgcaaaaatttgaacgtgcaaaaaaattaccaataatccagaaaattattatcatggattctaaaacgga  
ttaccagggtttcagtcgatgtacacgttcgtcacatctcatctacctcccggtttaatgaatacgattttgtaccagagtcctttgatcg  
tgacaaaacaattgcactgataatgaattcctctggatctactgggttacctaagggtgtggcccttcgcgcatagaactgcctgcgtca  
gattctcgcagatgccagatcctatttttgcaatcaaatcattccggatactgcgatttaagtgttgtccattccatcacggtttggaa  
tgtttactacactcggatattgatattgtggatttcgagtcgtcttaattgtatagattgaagaagagctgttttacgatcccttcaggatta

caaaattcaaagtgcgttgctagtagaccaacctattttcattcttcgccaaaagcactctgattgacaaatacatttatctaatttacacg  
aaattgcttctgggggacacacctttcgaaagaagtcggggaagcgggttgcaaacgcttccatcttcaggagatacgacaaggat  
atgggctcactgagactacatcagctattctgattacacccgaggggagatgataaacgggcgcggtcggtaaagttgtccattttt  
gaagcgaagggttggtatctggataccgggaaaacgctgggaggttaatacagagaggcgaattatgtgtcagaggacctatgattat  
gtccgggttatgtaacaatccggaagcgaccaacgccttgattgacaaggatggatggctacattctggagacatagcttactggga  
cgaagacgaacacttctcatagttgaccgctgaagtccttaataatacaaaaggatatcaggtggcccccgctgaattggaatcga  
tattgttacaacaccccaacatcttcgacgcgggctggcaggtcttcccgacgatgacgccggtgaacttcccgccgccgttggtg  
tttgagcacggaaagacgatgacggaaaagagatcgtggattacgtcgccagtcagtaacaaccgcgaaaagttgcgcgg  
aggagttgtgttggtgacgaagtaccgaaaggcttaccggaaaactcgacgcaagaaaaatcagagagatcctcataaaaggcca  
agaaggcggaagtcgcaattgcgcggccgctaactcgagaataaaatgaggaaattgcatcgcttctgagtaggtgtcattc  
tattctggggggtgggggtgggacagcaagggggaggattgggaagacaatagcaggcatgctggggatgcggtggggt  
ctatggcttctgaggcggaagaaccagctggggctctagggggtatccccacgcgccctgtagcggcgcatgaagcgcgggcg  
gtgtggtggttacgcgcagcgtgaccgctacacttgccagcgccctagcgccgctccttctgcttcttcccttcttctcgcacgt  
tcgccggcttccccgtcaagctctaaatcgggggctcccttaggggtccgatttagtcttacggcacctcgacccccaaaaactt  
gattagggtgatggttcacgtagtgggccatcgccctgatagacggtttctgccccttgacgttgagtcacgttcttaatagtgga  
ctctgttccaaactggaacaacactcaaccctatctcggcttattctttgattataagggttttgcgatttggcctattggttaaaaa  
atgagctgatttaacaaaaatttaacgcgaattaattctgtggaatgtgtgtcagttagggtgtggaaagtcggcaggtcccgagcag  
gcagaagtatgcaaagcatgcatctcaattagtcagcaaccagggtgtgaaagtcggcaggtcccgagcagcagaagtatgca  
aagcatgcatctcaattagtcagcaaccatagtcggcccttaactccgcccataactccgcccaggtccgcccattct  
ccgccccatggctgactaattttttatttatgcagaggccgaggccgcctctgcctctgagctattccagaagtagtgaggaggctt  
tttgaggccctaggcttttgcaaaaagctccgggagcttgatatccatttctggatctgatcagcacgtgatgaaaaagcctgaact  
caccgcgacgtctgtcgagaagttctgatcgaaaagttcgacagcgctccgacctgatgcagctctcgaggggcgaagaatctc  
gtgcttctcagcttgatgtaggagggcggtgatgtctcgggtaaatagctgcgccgatggttctacaaagatcgttatgtttatc  
ggcactttgcatcgccgcgctcccgaattccggaagtgttgacattggggaattcagcgagagcctgacctattgcatctccgcc  
gtgcacagggtgtcacgttgcaagacctgcctgaaaccgaactgccgctgttctgcagccggtcgcgaggccatggatgcgat  
cgctgcggccgatcttagccagacgagcgggttcggccattcggaccgcaaggaatcgttcaatacactacatggcgtgattca  
tatgcgcgattgctgatccccatgtgtatcactggcaaaactgtgatggacgacaccgtcagtcgctccgctcgcgaggctctcgtatg  
agctgatgctttggccgaggactgccccgaagtccggcacctcgtgcacgcggatttgggtccaacaatgtcctgacggacaat  
ggccgcataacagcgggtcattgactggagcagggcgatgttcggggattcccaatacagaggtcgccaacatcttcttctggaggcc  
gtggttggttgatgtagcagcagacgcgctacttcgagcggaggcatccggagcttgaggatcgccgcgggtccgggctat  
atgctccgattggtcttgaccaactctatcagagcttggtgacggcaatttcgatgatgcagcttgggcgagggctgatgcgacg  
caatcgtccgatccggagccgggactgtcggcgctacacaaatcgccgcagaagcgcggccgtctggaccgatggctgtgtag  
aagtactcgccgatagtggaaaccgacggccagcactgtccgaggggcaaaggaatagcacgtgctacgagatttcgattccac  
cgccgccttctatgaaagggtgggcttcggaatcgtttccgggacgccggctggatgatcctccagcgcggggatctcatgctgga  
gttcttcgccaccccaactgtttattgcagcttataatggttacaaataaagcaatagcatcacaatttcacaaataaagcattttttc

actgcattctagttgtggttgcctcaactcatcaatgtatcttatcatgtctgtataccgtcgacctctagctagagcttggcgtaatcatg  
gtcatagctgttctgtgtgaaattgtatccgctcacaattccacacaacatacagaccggaagcataaagtgtaaagcctgggggtg  
cctaagttagtgagctaaactcacattaattgcgttgcgtcactgcccgtttccagtcgggaaacctgtcgtgccagctgcattaatga  
atcggccaacgcgcggggagaggcgggttgcgtattgggcgctcttccgcttctcgtcactgactcgtcgcctcggcgttcgg  
ctgcggcgagcgggtatcagctcactcaaaggcggtaatacggttatccacagaatcaggggataacgcaggaaagaacatgtgag  
caaaaggccagcaaaaggccagggaaccgtaaaaaggccgcgttgcgtggcgttttccataggctccgccccctgacgagcatca  
caaaaatcgacgtcaagtcagaggtggcgaaacccgacaggactataaagataaccaggcgtttccccctggaagctccctcgtg  
cgctctcctgttccgacctgccgcttaccggataacctgtccgcctttctcccttcgggaagcgtggcgctttctcatagctcacgctgt  
aggatctcagttcgggtgtaggtcgttcgctccaagctgggctgtgtgcacgaacccccgttcagcccagaccgctgcgccttatcc  
ggtaactatcgtcttagtccaacccggtaagacacgacttatcgccactggcagcagccactggtaacaggattagcagagcgag  
gtatgtaggcgggtgctacagagttctgaagtgggtggcctaactacggctacactagaagaacagtatttggtagtctgcgtctgctga  
agccagttaccttcggaaaaagagttggtagctcttgatccggcaaaacaaaccaccgctggtagcgggtttttgtttgcaagcagca  
gattacgcgcagaaaaaaggatctcaagaagatcctttgatctttctacgggggtctgacgctcagtggaaacgaaaactcacgttaa  
gggattttggtcatgagattatcaaaaaggatcttcacctagatccttttaataaaaaatgaagttttaaatcaatctaaagtatatatgag  
taaacttggctgacagttaccaatgcttaatcagtgaggcacctatctcagcgatctgtctatttcgttcacatagttgcctgactccc  
cgctcgttagataactacgatacgggagggccttaccatctggccccagtgctgcaatgataccgcgagaccacgctcaccggctc  
cagatttatcagcaataaaccagccagccggaaggccgagcgcagaagtggctcctgcaactttatccgcctccatccagcttatta  
attgttgccgggaagctagagtaagtagttcgccagttaatagtttgcgaacgttgttgccattgctacaggcatcgtgggtgcacgct  
cgctgtttggtatggcttcattcagctccggttccaacgatcaaggcgagttacatgatccccatgttgtgcaaaaaagcggttagc  
tccttcggctcctccgatcgttgcagaagtaagttggccgcagtggtatcactcatggttatggcagcactgcataattctcttactgtcat  
gccatccgtaagatgcttttctgtgactgggtgagtactcaaccaagtcattctgagaatagtgtagtgcggcgaccgagttgctcttggcc  
ggcgtcaatacgggataataccgcgccacatagcagaactttaaaagtgtcatcattgaaaacgttcttcggggcgaaaactctc  
aaggatcttaccgctgttgagatccagttcgtatgaaccactcgtgcaccaactgatcttcagcatcttttactttaccagcgtttctg  
ggtagcaaaaaacggaaggcaaaatgccgaaaaaagggaataaaggcgacacggaaatgtgaatactcatactcttctctttt  
caatattattgaagcatttatcagggtattgtctcatgagcggatacatattgaatgtatttagaaaaataaacaatagggggtccgcg  
cacatttccccgaaaagtgccacctgacgtc

[0480] pCMR2 (SEQ ID NO: 93)

gttgacattgattattgactagttattaatagtaatcaattacggggtcattagttcatagcccatatatggagttccgcgttacataactta  
cggtaaatggccgcctggctgaccgccaacgacccccgccattgacgtcaataatgacgtatgttccatagtaacgccaata  
gggactttcattgacgtcaatgggtggagtatttacggtaactgccacttggcagtagcatcaagtgtatcatatgccaagtccgcc  
ccctattgacgtcaatgacggtaaatggccgcctggcattatgccagtagcatgaccttacgggactttcctacttggcagtagcatct  
acgtattagtcacgctattaccatgggtgatgcgggtttggcagtagacaccaatgggcgtggatagcgggttgactcacggggatttoca  
agtctccaccccaattgacgtcaatgggagttgttttggcaccaaaatcaacgggactttccaaaatgtcgttaataaccccgccccgtt  
gacgcaaatgggcggtaggcgtgtacgggtgggaggtctatataagcagagctcgtttagtgaaccgtaaagctttcggcgccacg  
gtaccatgggatccgaagacgccaaaaacataaagaaaggccggcgccattctatcctctagaggatggaaccgctggagagc

aactgcataaggctatgaagagatacgccttggtcctggaacaattgctttacagatgcacatatcagggtgaacatcacgtacgc  
ggaatacttcgaaatgtccgttcggttggcagaagctatgaaacgatatgggctgaatacaaatcacagaatcgtcgtatgcagtga  
aactctcttcaattcttattgccggtgttggcgcggttattatcggagttgcagttgcgccgcgaacgacatttataatgaacgtgaatt  
gctcaacagtatgaacatttcgcagcctaccgtagtggttgttccaaaaagggttgcaaaaaatttgaaactgcaaaaaaattacc  
aataatccagaaaattattatcatggattctaaaacggattaccagggttgcagtcgatgtacacgttcgtcacatctcatctacctccc  
gggtttaatgaatacatttgtaccagagtcctttgatcgtgacaaaacaattgcactgataatgaattcctctggatctactgggttacc  
taagggtgtggcccttcgcatagaactgcctgcgtcagattctcgcagtcgagatcctatttttgcaatcaaatcattccggatac  
tgcgattttaagtgtgttccattccatcacgggttggaaatgttactacactcggatatttgatatgtggatttcgagtcgtcttaattgata  
gatttgaagaagagctgttttacgatcccttcaggattacaaaattcaaagtgcgttgctagtaccaacctattttcattcttcgcaaa  
agcactctgattgacaaatacatttatctaatttacacgaaattgcttctgggggcgcacctcttcgaaagaagtgggggaagcgg  
tgcaaacgcttccatctccaggatacgaacaggatagggtcactgagactacatcagctattctgattacacccgagggggat  
gataaacgggcgcggtcggttaaagttgttccatttttgaagcgaaggttggatctggataccgggaaaacgctgggcgttaac  
agagaggcgaattatgtgtcagaggacctatgattatgtccggttatgtaacaatccggaagcgaccaacgccttgattgacaagg  
atggatggctacattctggagacatagcttactgggacgaagacgaacactcttcatagttgaccgcttgaagctttaattaataca  
aaggatatcaggtggcccccgtgaattggaatcgaattgttacaacaccccaacatcttcgacgcgggcgtggcaggtcttccc  
acgatgacgccgggtgaactcccgcgcggttgttggagcacggaaagacgatgacggaaaaagagatcgtggattacgtc  
gccagtcaagtaacaaccgcgaaaaagttgcgcggaggagttgtgttggacgaagtaccgaaagcttaccggaaaactcg  
acgcaagaaaaatcagagagatcctcataaaggccaagaaggcggaaagtccaaattgcgcggccgctaactcgagaataaac  
aagtaacaacaacaattgcattcattttatgttccaggttcagggggaggtgtgggaggtttttaagcaagtaaacctctacaaatg  
tggtatggctgattatgatccggctgcctcgcgcgttccggtgatgacggtgaaaacctctgacacatgcagctcccggagacggtc  
acagcttgtctgtaagcggatgccgggagcagacaagcccgtaggcgtcagcgggtgttggcgggtgtcggggcgcagccatg  
aggctgactctagaggatcgtgccccgccccggacgaactaaacctgactacgacatctctcccccttctcgcggggcagtgca  
tgtaatcccttcagttggttggtacaacttgccaactgggccctgttccacatgtgacacggggggggaccaaacacaaaagggttc  
tctgactgtagtgcacatccttataaatggatgtgcacatttgccaacactgagtggctttcatcctggagcagacttgcagctgtgga  
ctgcaacacaacattgcctttatgtgtaactcttggtgaagctcttacaccaatgctgggggacatgtacctcccagggggccagga  
agactacgggaggctacaccaacgtcaatcagaggggcctgtgtagctaccgataagcggacctcaagagggcattagcaata  
gtgtttataaggcccccttgtaaccctaaacgggtagcatatgcttccgggtagtagtatatactatccagactaacctaatcaata  
gcatatgttaccacacgggaagcatatgctatcgaattagggttagtaaaagggtcctaaggaacagcgatatctcccaccccatga  
gctgtcacgggtttttacatggggtcaggattccacgagggtagtgaaccatttttagtcacaagggcagtggtgaagatcaagga  
gcgggcagtgaaactctcctgaatcttcgctgcttcttcattctccttcgtttagctaataagaataactgctgagttgtgaacagtaaggt  
gtatgtgaggtgtcgaacaaggtttcaggtgacgccccagaataaaatttgacgggggggtcagtggtggcattgtgctatg  
acaccaatataaccctcacaaacccttgggcaataaatactagtgtaggatgaaacattctgaatatcttaacaatagaatccatg  
gggtgggggacaagccgtaaaagactggatgtccatctcacacgaatttatggctatgggcaacacataatcctagtgcataatgatact  
gggggttattaagatgtgtccaggcagggaccaagacaggtgaaccatgttgttactctatttgaacaaggggaaagagagtgg  
acgccgacagcagcggactccactggtgtcttaacacccccgaaaattaaacgggggtccacgccaatggggcccataaaca

[illegible]



cctcgtccatggttatccccctcttctttaggtccactgccgccggagcctcttggtccagatgtgtctccctctctcctagggccat  
ttccaggtcctgtacctggccccctgctcagacatgattcacactaaaagagatcaatagacatcttattagacgacgctcagtgaata  
cagggagtgcagactcctgccccctcaacagccccccaccctcatccccctcatggctgctgacagacatccaggtctgaaa  
attccccatcctccgaaccatcctcgtcctcatcaccattactgcagcccgaaaactcccgtgaacatcctcaagattgcgtcc  
tgagcctcaagccaggcctcaaatcctcgtcccccttttctggtgacggtagggatggggattctcgggacccctcctcttctcttc  
aaggtcaccagacagagatgctactggggcaacggaagaaaagctgggtgctggcctgtgaggatcagcttatcgatgataagctg  
tcaaacatgagaattcttgaagacgaaagggcctcgtgatacgcctatttttataggtaatgtcatgataataatggttcttagacgtca  
gggtggcacttttggggaaatgtgcgcggaaccctatttgtttattttctaaatacattcaaatatgtatccgctcatgagacaataacc  
ctgataaatgcttcaataattgaaaaaggaagagtatgagtattcaacattccgtgctgcccttattccctttttgctggcattttgcctt  
cctgttttctcaccagaaacgctggtgaaagtaaaagatgctgaagatcagttgggtgcacgagtgggttacatcgaactggatc  
tcaacagcggtaagatccttgagagtttgcgccgaagaacgtttccaatgatgagcacttttaaagtctgctatgtggcgcggtat  
tatcccggtgttgacgccgggcaagagcaactcggctgccgcatacactattctcagaatgacttggttagtactaccagtcacaga  
aaagcatcttacggatggcatgacagtaagagaattatgcagtgtgccataaccatgagtataactgcggccaacttacttctg  
acaacgatcggaggaccgaaggagctaaccgctttttgcacaacatgggggatcatgtaactgccttgatcgttgggaaccgga  
gctgaatgaagccataccaaacgacgagcgtgacaccacgatgcctgcagcaatggcaacaacgttgcgcaaacatttaactggc  
gaactacttacttagcttcccggaacaattaatagactggatggagggcgataaagtgcaggaccacttctgcgctcggcccttc  
cggctggctggtttattgctgataaatctggagccggtagcgtgggtctcgcggtatcattgcagcactggggccagatggtaagc  
cctcccgatcgtagtattctacacgacggggagtcaggcaactatggatgaacgaaatagacagatcgtgagataggtgcctcac  
tgattaagcattggttaactgtcagaccaagttactcatatatacttttagattgatttaaaacttcatttttaatttaaaaggatctaggtgaa  
gatcctttttgataatctcatgacaaaaatcccttaacgtgagtttctgttccactgagcgtcagaccccgtagaaaagatcaaaggatc  
ttcttgagatccttttttctgcgcgtaactgtgctgttgcacaacaaaaaccaccgctaccagcgggtggtttgttccggatcaagag  
ctaccaactctttttccgaaggtaactggcttcagcagagcgcagataccaaatactgtccttctagttagccgtagttaggccacca  
cttcaagaactctgtagcaccgcctacatacctcgtctgctaactcgttaccagtggtgctgctgccagtggcgataagtcgtgtcttac  
cgggttgactcaagacgatagttaccggataaggcgcagcggctcgggtgaacgggggggttcgtgcacacagcccagcttgga  
gcgaacgacctacaccgaactgagatacctacagcgtgagctatgagaaagcggcacgcttccgaaggagaaaggcgggaca  
ggatatccggtaagcggcagggtcggaaacaggagagcgcacgaggagcttccagggggaaacgcctggatatctttatagtcctgt  
cgggtttcggcacctctgacttgagcgtcgattttgtgatgctcgtcagggggggcgagcctatggaaaaacgccgcaacgcgg  
cctttttacggttctgaccttttctggccttgaagctgtccctgatggctgtcatctacctgcctggacagcatggcctgcaacgcgg  
gcatcccgatgccgccggaagcgagaagaatcataatgggggaaggccatccagcctcgcgtcgcgaacgccagcaagacgtag  
cccagcgcgtcggccccgagatgcgccgcgtcgggtgctggagatggcggacgcgatggatatgttctgccaagggttggttg  
cgattcacagttctccgcaagaattgattggtccaatcttggagtggatgaatccgttagcgaggtgccgccctgcttcatccccgt  
ggcccgttgctcgcgttctggcgggtgtccccggaagaaatataatttgcattgtttagttctatgatgacaaaaccccgccagcg  
tcttgtcattggcgaattcgaacacgcagatgcagtcggggcgggcggtccgaggtccacttcgatattaaggtgacgcgtgtg  
gcctcgaacaccgagcgacctgcagcgacctgctaacagcgtcaacagcgtccgcagatccgggggggcaatgagatatg  
aaaaagcctgaactcaccgcgacgtctgtcgagaagtttctgatcgaaaagttcgacagcgtctccgacctgatgcagctctcggag



ggcgaagaatctcgtgctttcagcttcgatgtaggagggcggtggatatgtcctgcgggtaaataagctgcgccgatggtttctacaaag  
atcgttatgtttatcggcactttgcatcggccgcgctcccattccggaagtgttgacattggggaattcagcgagagcctgacctat  
tgcattctcccgccgtgcacaggggtgcacgttgcaagacctgcctgaaaccgaactccccgctgttctgcagccggtcgcggagg  
ccatggatgcgacgctgcggccgatcttagccagacgagcgggttcggccattcggaccgcaagggaatcggtaataactac  
atggcgtgattcatatgcgcgattgctgatccccatgtgtatcactggcaaactgtgatggacgacaccgtcagtgctccgtcgcg  
caggctctcgtatgagctgatgctttgggcccaggactgccccgaagtccggcacctcgtgcacgcggatttcggctccaacaatgt  
cctgacggacaatggccgcataacagcgggtcattgactggagcgaggcgatgttcggggattcccaatacagggtcgccaacatc  
tttcttggaggccgtgggttgcttgatggagcagcagacgcgctacttcgagcggaggcatccggagcttcgaggatcgccgcg  
gctccggggtatatgtccgcattggctttagcaactctatcagagcttgggtgacggcaatttcgatgatgcagcttgggcgag  
ggtcgatgcgacgcaatcgtccgatccggagccgggactgtcggggtacacaaatcgccgcagaagcgcggccgtctggac  
cgatggctgtgtagaagtactcgccgatagtggaaaccgacgccccagcactcgtccggatcgggagatgggggaggctaactg  
aaacacggaaggagacaataaccggaaggaaaccgcgctatgacggcaataaaaagacagaataaaacgcacgggtgttgggtc  
gtttgttcataaacgcgggggttcgggtcccagggtggcactctgtcgataccccaccgagacccattggggccaatacggccgcg  
tttcttcttttccccacccccacccccaaagtccgggtgaaggcccagggtcgcagccaacgtcggggcgagggccctgccata  
gccactggccccgtgggttagggacggggtccccatggggaatggtttatggttcgtgggggttatttttggcggtgcgtggg  
gtcaggtccacgactggactgagcagacagacctgggttttggatggcctgggcatggaccgcatgtactggcgcgacacgaa  
caccggcgctctgtgggtgccaacacccccgacccccaaaaaccaccgcgcggatttctggcgtgccaagctagtcgaccaatt  
ctcatgtttgacagcttatcatcgcagatccgggcaacgttgttgcattgctgcaggcgagaaactggtaggtatggaagatctatac  
attgaatcaatattggcaattagccatattagtcattggttatatagcataaatcaatattggctattggccattgcatacgttgcattat  
cataatgtacatttatattggctcatgtccaatatgaccgccaat

[0481] pMCP1 (SEQ ID NO: 94)

gacggatcgggagatctcccgatcccctatggtgcaactctcagtacaatctgctctgatccgcatagttaagccagtatctgctccct  
gcttgtgtgttgagggtcgtgagtagtgcgcgagcaaaatttaagctacaacaaggcaaggcttgaccgacaattgcatgaagaat  
ctgcttaggggttaggcgttttgcgtgcttcgcgatgtacgggcccagatatacgcgttgacattgattattgactagttattaatagtaac  
aattacgggggtcattagttcatagcccataatggagttccgcgttacataacttacggtaaatggcccgcctggctgaccgccaac  
gacccccgccattgacgtcaataatgacgtatgttccatagtaacgccaatagggactttcattgacgtcaatgggtggagtattt  
acggtaaaactgcccacttggcagtacatcaagtgtatcatatgccaaagtacgccccctattgacgtcaatgacggtaaatggccgc  
ctggcattatgccagttacatgacctatgggactttcctacttggcagtacatctacgtattagtcacgctattaccatgggtgatcgg  
tttggcagttacatcaatggggtggatagcgggttgactcacggggatttccaagtctccacccattgacgtcaatgggagtttgtt  
tggcaccaaaatcaacgggactttccaaaatgtcgtacaactccgccccattgacgcaaatggcggttaggcgtgtacgggtggga  
ggcttatataagcagagctctctggctaactaagctttcggcgcgccgaggtaccatgggatccgaagacgccccaaacataaaga  
aaggccccggcgccattctatcctctagaggatggaaccgctggagagcaactgcataaggctatgaagagatacggcctgggtcct  
ggaacaattgcttttacagatgcacatatcgagggtgaacatcacgtacgcggaatacttcgaaatgtccgttcggttggcagaagctat  
gaaacgatatgggctgaatacaaatcacagaatcgtcgtatgcagtgaaaactcttcaattctttatgccggtgttggcgcggttattt  
atcggagttgcagttgcgcccgcgaacgacattataatgaacgtgaattgtcaacagtatgaacatttcgagcctaccgtagtgtt

tgtttccaaaaaggggttgcaaaaaatttgaacgtgcaaaaaaattaccaataatccagaaaattattatcatggattctaaaacgga  
ttaccagggtttcagtcgatgtacacgttcgtcacatctcatctacctcccggtttaatgaatacgaattttgtaccagagtcctttgatcg  
tgacaaaacaattgcactgataatgaattcctctggatctactgggttacctaaggggtgtggccctccgcatagaactgcctgcgtca  
gattctcgcagatgccagagatcctatTTTTGGCAATCAATCATTCCGGATACTGCGATTTAAGTGTGTCCATTCCATCAGGTTTTGGA  
TGTTTACTACACTCGGATATTTGATATGTGGATTTCGAGTCGTCTTAATGTATAGATTGAAGAAGAGCTGTTTTACGATCCCTCAGGATTA  
CAAAATCAAAGTGCCTTGCTAGTACCAACCCTATTTTCATTCTTCGCCAAAAGCACTCTGATTGACAAATACGATTATCTAATTACACG  
AAATTGCTTCTGGGGGCGCACCTCTTTCGAAAGAAGTCGGGGAAGCGGTTGCAAAACGCTTCCATCTTCAGGGATACGACAAGGAT  
ATGGGCTCACTGAGACTACATCAGCTATTCTGATTACACCCGAGGGGGATGATAAACCGGGCGCGGTGCGTAAAGTTGTCCATTTTT  
GAAGCGAAGGTTGTGGATCTGGATACCGGGAAAACGCTGGGCGTAAATCAGAGAGGCGAATTATGTGCAGAGGACCTATGATTAT  
GTCCGGTTATGTAACAATCCGGAAGCGACCAACGCCTTGATTGACAAGGATGGATGGCTACATTCTGGAGACATAGCTTACTGGGA  
CGAAGACGAACACTTCTCATAGTTGACCGCTTGAAGTCTTAATAATAACAAGGATATCAGGTGGCCCCCGCTGAATTGGAATCGA  
TATTGTTACAACACCCCAACATCTTCGACGCGGGCGTGGCAGGTCTTCCGACGATGACGCCGGTGAACCTCCCGCCCGCTTGTGT  
TTGGAGCACGGAAAGACGATGACGGAAAAAGAGATCGTGGATTACGTCGCCAGTCAAGTAACAACCGCAAAAAGTTGCGCGG  
AGGAGTTGTGTTGTGGACGAAGTACC GAAAGTCTTACC GAAAAC TGACGCAAGAAAAATCAGAGAGATCCTCATAAAGGCCA  
AGAAGGGCGGAAAGTCCAAATTGCGCGGCCGCTAACTCGAGAATAAAATGAGGAAATTGCATCGCATTGTCTGAGTAGGTGCATTC  
TATTCTGGGGGGTGGGGTGGGGCAGGACGCAAGGGGGAGGATTGGGAAGACAATAGCAGGCGATGCTGGGGATGCGGTGGGCT  
CTATGGCTTCTGAGGCGGAAAGAACCAGCTGGGGCTCTAGGGGGTATCCCCACGCGCCCTGTAGCGGGCGCATTAAAGCGCGGGG  
GTGTGGTGGTTACGCGCAGCGTGACCGCTACACTGCCAGCGCCCTAGCGCCCGCTCTTTCGCTTCTTCCCTTCTTCTCGCCACGT  
TCGCCGGCTTCCCCGTCAAGCTCTAAATCGGGGTCCCTTAGGGTTCGATTAGTGCTTTACGGCACCTCGACCCAAAAAACTTG  
ATTAGGTGATGGTTCACGTACCTAGAAGTTCCTATTCCGAAGTTCCTATTCTCTAGAAAGTATAGGAACCTCCTTGCCAAAAAGCCTG  
AATCACCGCGACGTCTGTCGAGAAGTTCTGATCGAAAAGTTCGACAGCGTCTCCGACCTGATGCAGCTCTCGGAGGGCGAAGAA  
TCTCTGTCTTTCAGCTTCGATGTAGGAGGGCGTGGATATGTCCTCGGGTAAATAGCTGCGCCGATGGTTTCTACAAAGATCGTTATGTT  
ATCGGCACCTTGCATCGGCCGCGCTCCCGATTCCGGAAGTGCTTGACATTGGGGAATTCAGCGAGAGCCTGACCTATTGCATCTCCCG  
CCGTGCACAGGGTGTACGTTGCAAGACCTGCTGAAACCGAACTGCCGCTGTTCTGCAAGCGGTGCGGAGGCCATGGATGCG  
ATCGCTGCGGCCGATCTTAGCCAGACGAGCGGGTTCGGCCATTCCGACCGCAAGGAATCGGTCAATACACTACATGGCGTGATT  
CATATGCGCGATTGCTGATCCCCATGTGTACTGGCAAACGTGTATGGACGACACCGTCAGTGCCTCGTCGCGCAGGCTCTCGAT  
GAGCTGATGCTTTGGCCGAGGACTGCCCCGAAGTCCGGCACCTCGTGCAGCAAAACAACCCGCTGGTAGCGGTTTTTTGTTG  
CAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAGATCCTTGATCTTTCTACGGGGTCTGACGCTCAGTGAACGAAAA  
CTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGGATCTTACCTAGATCCTTTAAATAAAAATGAAGTTTAAATCAATCTAAG  
TATATAGAGTAAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTCTGTTACCATAGTTGC  
CTGACTCCCCGTCGTGTAGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGACCCACGCT  
CACC GGCTCCAGATTATCAGCAATAAACCGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGTCTGCAACTTATCCGCCTCCATC  
CAGTCTATTAATTGTGCCGGGAAGCTAGAGTAAGTAGTTCGCCAGTTAATAGTTTGCACAACGTTGTTGCCATTGCTACAGGCATCGT  
GTGTCACGCTCGCTGTTGGTATGGCTTCACTCAGCTCCGGTCCCAACGATCAAGGCGAGTTACATGATCCCCATGTTGTGAAAAA  
AGCGGTTAGCTCCTTGGTCTCCGATCGTTGTCAGAAGTAAGTTGGCCGAGTGTTATCACTCATGGTTATGGCAGCACTGCATAATC

tcttactgtcatgccatccgtaagatgcttttctgtgactggtgagtactcaaccaagtcattctgagaatagtgtatgcggcgaccgag  
ttgctcttgcccggtcaatacgggataataccgcgccacatagcagaactttaaagtgtcatcattggaaaacgttcttcgggg  
cgaaaactctcaaggatcttaccgctgttgagatccagttcgtatgaaccactcgtgcaccaactgatcttcagcatctttactttca  
ccagcgtttctgggtgagcaaaaacaggaaggcaaaatgccgcaaaaagggaataaggcgacacggaaatgttgaatactcat  
actcttccttttcaatattattgaagcattatcagggttattgtctcatgagcggatacatattgaatgtatttagaaaaataaacaata  
gggggtccgcgcacatttccccgaaaagtgccacctgacgtc

[0482] Equivalents:

[0483] The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described will become apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications are intended to fall within the scope of the appended claims.

[0484] Various publications are cited herein, the disclosures of which are incorporated by reference in their entireties.

**WHAT IS CLAIMED IS:**

1. A method for identifying a compound that modulates untranslated region-dependent expression of a vascular endothelial growth factor (VEGF) gene, said method comprising:
  - (a) contacting a member of a library of compounds with a cell containing a nucleic acid comprising a reporter gene operably linked to an UTR of the VEGF gene; and
  - (b) detecting a reporter protein translated from said reporter gene, wherein a compound that modulates untranslated region-dependent expression of a VEGF gene is identified if the expression of said reporter gene in the presence of a compound is altered as compared to the expression of said reporter gene in the absence of said compound or the presence of a control.
2. A method for identifying a compound that modulates untranslated region-dependent expression of a VEGF gene, said method comprising:
  - (a) contacting a member of a library of compounds with a cell-free translation mixture and a nucleic acid comprising a reporter gene operably linked to an UTR of the VEGF gene; and
  - (b) detecting the expression of said reporter gene, wherein a compound that modulates untranslated region-dependent expression of a VEGF gene is identified if the expression of said reporter gene in the presence of a compound is altered as compared to the expression of said reporter gene in the absence of said compound or the presence of a control.
3. The method of claim 1 or 2, wherein the UTR of the VEGF gene is the 5' untranslated region (5' UTR) of a VEGF gene.
4. The method of claim 3 wherein the 5' UTR of the VEGF gene is operably linked upstream of the reporter gene.
5. The method of claim 1 or 2, wherein the UTR of VEGF gene is the 3' untranslated region (3' UTR) of a VEGF gene.

6. The method of claim 5, wherein the 3' UTR of the VEGF gene is operably linked downstream of the reporter gene.
7. The method of claim 3, wherein the nucleic acid further comprises the 3' UTR of a VEGF gene.
8. The method of claim 7, wherein the 3' UTR of the VEGF gene is operably linked downstream of the reporter gene.
9. The method of claim 1 or 2, wherein the reporter gene further comprises an intron.
10. The method of claim 1 or 2, wherein the UTR of the VEGF gene comprises an iron response element ("IRE"), internal ribosome entry site ("IRES"), upstream open reading frame ("uORF"), or AU-rich element ("ARE").
11. The method of claim 1 or 2, wherein the nucleic acid is further polyadenylated at the 3' end.
12. The method of claim 1 or 2, wherein the nucleic acid is not capped at the 5' end.
13. The method of claim 1 or 2, wherein the reporter gene encodes firefly luciferase, renilla luciferase, click beetle luciferase, green fluorescent protein, yellow fluorescent protein, red fluorescent protein, cyan fluorescent protein, blue fluorescent protein, beta-galactosidase, beta-glucoronidase, beta-lactamase, chloramphenicol acetyltransferase, or alkaline phosphatase.
14. The method of claim 1, wherein said cell is stably transfected with said nucleic acid.
15. The method of claim 1, wherein said cell is transiently transfected with said nucleic acid.
16. The method of claim 1, wherein said cell is transfected with an episomal expression vector comprising said nucleic acid.

17. The method of claim 1 or 2 further comprising measuring the effect of said compound on the expression of the VEGF gene.

18. The method of claim 1, wherein the cell is a human cell, a yeast cell, a mouse cell, a rat cell, a Chinese hamster ovary ("CHO") cell, a MCF-7 cell, a primary cell, or an undifferentiated cancer cell.

19. The method of claim 18 wherein the human cell is a HeLa cell or a 293 cell.

20. The method of claim 3, wherein the cell-free translation mixture is a cell extract.

21. The method of claim 20, wherein the cell extract is derived from is a human cell, a yeast cell, a mouse cell, a rat cell, a Chinese hamster ovary ("CHO") cell, a *Xenopus* oocyte, a MCF-7 cell, a primary cell, an undifferentiated cancer cell, a reticulocyte, or a rye embryo.

22. The method of claim 1 or 2, wherein the compound is selected from a combinatorial library of compounds comprising peptoids, random biooligomers, diversomers, vinylogous polypeptides, nonpeptidal peptidomimetics, oligocarbamates, peptidyl phosphonates, peptide nucleic acid libraries, antibody libraries, carbohydrate libraries, and small organic molecule libraries.

23. The method of claim 22, wherein the small organic molecule libraries are libraries of benzodiazepines, isoprenoids, thiazolidinones, metathiazanones, pyrrolidines, morpholino compounds, or diazepindiones.

24. The method of claim 1, wherein the step of contacting a library of compounds with a cell is in an aqueous solution comprising a buffer and a combination of salts.

25. The method of claim 24, wherein the aqueous solution approximates or mimics physiologic conditions.

26. The method of claim 24, wherein the aqueous solution further comprises a detergent or a surfactant.

27. The method of claim 1 or 2 further comprising (c) determining the structure of the compound that modulates untranslated region-dependent expression of the VEGF gene.
28. The method of claim 27, wherein the structure of the compound is determined by mass spectroscopy, NMR, vibrational spectroscopy, or X-ray crystallography.
29. The method of claim 1 or 2, wherein the compound directly binds to an RNA transcribed from the VEGF gene.
30. The method of claim 1 or 2, wherein the compound binds to one or more proteins that modulate untranslated region-dependent expression of the VEGF gene.
31. The method of claim 7, wherein the compound disrupts an interaction between the 5' UTR and the 3' UTR of the VEGF gene.
32. A method of modulating the expression of a VEGF gene comprising contacting a cell expressing the VEGF gene with a compound, or a pharmaceutically acceptable salt thereof, identified according the method of claim 1 or 2.
33. A method of treating, preventing or ameliorating cancer or one or more symptoms thereof, said method comprising administering to a subject in need thereof an effective amount of a compound, or a pharmaceutically acceptable salt thereof, identified according to the method of claim 1 or 2, wherein said effective amount decreases the expression of the VEGF gene.
34. A method of inhibiting or reducing angiogenesis, said method comprising administering to a subject in need thereof a prophylactically or therapeutically effective amount of a compound, or a pharmaceutically acceptable salt thereof, identified according to the method of claim 1 or 2, wherein said effective amount decreases the expression of the VEGF gene.
35. A method of identifying a compound that inhibits or reduces angiogenesis, said method comprising:

- (a) contacting a member of a library of compounds with a cell containing a nucleic acid comprising a reporter gene operably linked to an UTR of a VEGF gene; and
- (b) detecting the expression of said reporter gene, wherein if a compound that reduces the expression of said reporter gene relative to the expression of said reporter gene in the absence of said compound or the presence of a control is detected in (b), then
- (c) contacting the compound with a tumor cell and detecting the proliferation of said tumor cell, so that if the compound reduces or inhibits the proliferation of the tumor cell, the compound is identified as a compound that inhibits or reduces angiogenesis.

36. The method of claim 38 further comprising (d) testing said compound in an animal model for angiogenesis, wherein said testing comprises administering said compound to said animal model and verifying that angiogenesis is inhibited by said compound in said animal model.

37. A method of treating, preventing or ameliorating cancer or one or more symptoms thereof, said method comprising administering to a subject in need thereof an effective amount of a compound, or a pharmaceutically acceptable salt thereof, identified according to the method of claim 35.

38. A method of inhibiting or reducing angiogenesis, said method comprising administering to a subject in need thereof an effective amount of a compound, or a pharmaceutically acceptable salt thereof, identified according to the method of claim 35.

39. A method of identifying a compound that inhibits or reduces angiogenesis, said method comprising:

- (a) contacting a member of a library of compounds with a cell-free translation mixture and a nucleic acid comprising a reporter gene operably linked to an UTR of a VEGF gene; and
- (b) detecting the expression of said reporter gene, wherein if a compound that reduces the expression of said reporter gene relative to the expression of said reporter gene in the absence of said compound or the presence of a control is detected in (b), then




- (c) contacting the compound with a tumor cell and detecting the proliferation of said tumor cell, so that if the compound reduces or inhibits the proliferation of the tumor cell, the compound is identified as a compound that inhibits or reduces angiogenesis.

40. The method of claim 39 further comprising (d) testing said compound in an animal model for angiogenesis, wherein said testing comprises administering said compound to said animal model and verifying that angiogenesis is inhibited by said compound in said animal model.


41. A method of inhibiting or reducing angiogenesis, said method comprising administering to a subject in need thereof an effective amount of a compound, or a pharmaceutically acceptable salt thereof, identified according to the method of claim 39.

42. A method of treating, preventing or ameliorating cancer or one or more symptoms thereof, said method comprising administering to a subject in need thereof an effective amount of a compound, or a pharmaceutically acceptable salt thereof, identified according to the method of claim 39.

43. The method of claim 1 or 2 further comprising determining the specificity of the compound for the VEGF untranslated region.





FIGURES 1A TO 1C



## Polylinker sites

p2luc1	Sall	ApaI	Eco471II	BamHI	SmaI	SacI
GAA CAA ATG TCG ACG GCG GCG CTT AGG AGA TCT AGC GCT GGA TCC CCC GGC GAG CTC AUG GAA GAC						
RLUC	M	S				N E FLUC

FIGURE 2A




FIGURES 2B TO 2C

A.

pLuc/VEGF5'+3'UTR



B.



FIGURES 3A TO 3B




FIGURE 4

## Stability of stable cell line B9







FIGURE 5




FIGURES 6A TO 6B



A.



B.



FIGURES 7A TO 7B

# Stable Cell Line Expression Vector

Hind III Asc I Kpn I BamHI | Reporter | Not I Xho I




FIGURE 8A

## Stable Cell Line Expression Vector




FIGURE 8B

# Episomal Mammalian Expression Vector

Hind III Asc I Kpn I BamH I Not I Xho I

Reporter




FIGURE 8C

## SEQUENCE LISTING

<110> PTC Therapeutics, Inc.

<120> METHODS FOR IDENTIFYING COMPOUNDS THAT MODULATE UNTRANSLATED  
REGION-DEPENDENT GENE EXPRESSION AND METHODS OF USING SAME

<130> 10589-012-228

<140>  
<141>

<150> 60/441,637  
<151> 2003-01-21

<160> 94

<170> PatentIn version 3.2

<210> 1  
<211> 14  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: Motif

<220>  
<221> misc\_feature  
<222> 3, 7, 8, 11  
<223> n = a, t, c, or g

<220>  
<221> misc\_feature  
<222> (7)..(8)  
<223> This represents one form of the sequence as described, other forms  
described may have up to five nucleotides in this variable region

<400> 1  
ggntggnggg ntgg 14

<210> 2  
<211> 14  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: Motif

<220>  
<221> misc\_feature  
<222> 3, 4, 7, 8, 11, 12  
<223> n = a, t, g or c

<220>  
<221> misc\_feature  
<222> (2)..(12)

<223> This represents one form of the sequence as described, other forms described have longer variable regions, typical is 2 - 10 nucleotides

<400> 2  
ggnnnggnngg nngg 14

<210> 3  
<211> 14  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: Motif

<220>  
<221> misc\_feature  
<222> 3, 4, 7, 8, 11, 12  
<223> n = a, t, g, or c

<220>  
<221> misc\_feature  
<222> (2)..(12)  
<223> This represents one form of the sequence as described, other forms described have longer variable regions, typical is 2 - 10 nucleotides

<400> 3  
ggnnnggnngg nngg 14

<210> 4  
<211> 19  
<212> RNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: Motif

<400> 4  
ccccrcccuc uuccccaag 19

<210> 5  
<211> 152  
<212> DNA  
<213> Homo sapiens

<400> 5  
gcagaggacc agctaagagg gagagaagca actacagacc cccctgaaa acaaccctca 60  
gacgccacat cccctgacaa gctgccaggc aggttctctt cctctcacat actgaccac 120  
ggctccaccc tctctcccct ggaaaggaca cc 152

<210> 6  
<211> 792  
<212> DNA

<213> Homo sapiens

<400> 6

```

tgaggaggac gaacatccaa ccttcccaaa cgcctcccct gcccgaatcc ctttattacc      60
ccctccttca gacaccctca acctcttctg gctcaaaaag agaattgggg gcttaggggc      120
ggaacccaag cttagaactt taagcaacaa gaccaccact tcgaaacctg ggattcagga      180
atgtgtggcc tgcacagtga attgctggca accactaaga attcaaactg gggcctccag      240
aactcactgg ggcctacagc tttgatccct gacatctgga atctggagac cagggagcct      300
ttggttctgg ccagaatgct gcaggacttg agaagacctc acctagaaat tgacacaagt      360
ggaccttagg ccttcctctc tccagatgtt tccagacttc cttgagacac ggagcccagc      420
cctcccatg gagccagctc cctctattta tgtttgcact tgtgattatt tattatttat      480
ttattattta tttatttaca gatgaatgta tttatttggg agaccggggg atcctggggg      540
accaatgta ggagctgcct tggctcagac atgttttccg tgaaaacgga gctgaacaat      600
aggctgttcc catgtagccc cctggcctct gtgccttctt ttgattatgt tttttaaaat      660
atztatctga ttaagttgtc taaacaatgc tgatttggtg accaactgtc actcattgct      720
gagcctctgc tcccagggg agttgtgtct gtaatcgccc tactattcag tggcgagaaa      780
taaagtttgc tt                                                    792

```

<210> 7

<211> 21

<212> RNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: Motif

<400> 7

```

auuuauuuau uuauuuauuu a                                                    21

```

<210> 8

<211> 40

<212> DNA

<213> Homo sapiens

<400> 8

```

kctggaggat gtggctgcag agcctgctgc tcttgggcac                                40

```

<210> 9

<211> 289

<212> DNA

<213> Homo sapiens

<400> 9

```

gccggggagc tgctctctca tgaaacaaga gctagaaact caggatgggc atcttggagg      60

```

gaccaagggg tgggccacag ccatggtggg agtggcctgg acctgccctg ggccacactg 120  
 accctgatac aggcattggca gaagaatggg aatattttat actgacagaa atcagtaata 180  
 tttatatatt tatattttta aaatattttat ttattttatt atttaagttc atattccata 240  
 tttattcaag atgttttacc gtaataatta ttattaaaaa tatgcttct 289

<210> 10  
 <211> 21  
 <212> RNA  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: Motif

<400> 10  
 auuuauuuau uuauuuauuu a 21

<210> 11  
 <211> 47  
 <212> DNA  
 <213> Homo sapiens

<400> 11  
 atcactctct ttaatcacta ctcacattaa cctcaactcc tgccaca 47

<210> 12  
 <211> 307  
 <212> DNA  
 <213> Homo sapiens

<400> 12  
 taattaaagt cttcccactt aaaacatatc aggccttcta tttattttatt taaatattta 60  
 aattttatat ttattgttga atgtatggtt gctacctatt gtaactatta ttcttaactct 120  
 taaaactata aatatggatc ttttatgatt ctttttgtaa gccctagggg ctctaaaatg 180  
 gtttacctta tttatcccaa aaatattttat tattatgttg aatgttaa atagtatcta 240  
 tgtagattgg ttagtaaaac tatttaataa atttgataaa tataaaaaaa aaaaacaaaa 300  
 aaaaaaa 307

<210> 13  
 <211> 15  
 <212> RNA  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: Motif

<220>

<221> misc\_feature  
 <222> (1)..(15)



<223> n = a, t, g or c

<400> 13  
nauuuauuuu uuan

15

<210> 14  
<211> 62  
<212> DNA  
<213> Homo sapiens

<400> 14  
ttctgcccctc gagcccaccg ggaacgaaag agaagctcta tctcgccctcc aggagcccag 60  
ct 62

<210> 15  
<211> 427  
<212> DNA  
<213> Homo sapiens

<400> 15  
tagcatggggc acctcagatt gttgttggtta atgggcattc cttctttctgg tcagaaacct 60  
gtccactggg cacagaactt atgttggttct ctatggagaa ctaaaagtat gagcgtagg 120  
acactatttt aattattttt aatttattaa tatttaaata tgtgaagctg agttaattta 180  
tgtaagtcatt atttatattt ttaagaagta ccacttgaaa cattttatgt attagttttg 240  
aaataataat ggaaagtggc tatgcagttt gaatatacctt tgtttcagag ccagatcatt 300  
tcttggaag ttaggctta cctcaaataa atggctaact tatacatatt tttaaagaaa 360  
tatttatatt gtatttatat aatgtataaa tggtttttat accaataaat ggcattttta 420  
aaaattc 427

<210> 16  
<211> 15  
<212> RNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: Motif

<220>  
<221> misc\_feature  
<222> (1)..(15)  
<223> n = a, t, g or c

<400> 16  
nauuuauuuu uuan

15

<210> 17  
<211> 701  
<212> DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 17

```

aagagctcca gagagaagtc gaggaagaga gagacggggt cagagagagc gcgcgggctg      60
gcgagcagcg aaagcgacag gggcaaagtg agtgacctgc ttttgggggt gaccgccgga      120
gcgcgggctg agccctcccc cttgggatcc cgcagctgac cagtcgcgct gacggacaga      180
cagacagaca cggcccccag cccagttac cacctcctcc cgggccggcg gcggacagtg      240
gacgcggcgg cgagccgcgg gcagggggccg gagcccgcgc ccggaggcgg ggtggagggg      300
gtcggagctc gcggcgctgc actgaaactt ttctgtccaa ttctgggctg ttctcgcttc      360
ggaggagccg tgggtccgcgc gggggaagcc gagccgagcg gagccgcgag aagtgtctagc      420
tcgggccggg agggagccgca gccggaggag ggggaggagg aagaagagaa ggaagaggag      480
agggggccgc agtggcgact cggcgctcgg aagccgggct catggacggg tgaggcggcg      540
gtgtgcgcag acagtgtctc agcgcgcgcg ctccccagcc ctggcccggc ctcgggccgg      600
gaggaagagt agctcgccga ggcgccgagg agagcgggccc gcccacagc ccgagccgga      660
gagggacgcg agccgcgcgc cccgggtcggg cctccgaaac c              701

```

&lt;210&gt; 18

&lt;211&gt; 1892

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 18

```

tgagccgggc aggaggaagg agcctccctc agggtttcgg gaaccagatc tctctccagg      60
aaagactgat acagaacgat cgatacagaa accacgctgc cgccaccaca ccatcaccat      120
cgacagaaca gtccttaatc cagaaacctg aaatgaagga agaggagact ctgcgcagag      180
cactttgggt ccggagggcg agactccggc ggaagcattc ccgggcgggt gaccagcac      240
ggtccctctt ggaattggat tcgccatttt atttttcttg ctgctaaatc accgagcccg      300
gaagattaga gagttttatt tctgggattc ctgtagacac accacccac atacatacat      360
ttatatatat atatattata tatatataaa aataaatatc tctattttat atatataaaa      420
tatatatatt ctttttttaa attaacagtg ctaatgttat tgggtgtcttc actggatgta      480
tttgactgct gtggacttga gttgggaggg gaatgttccc actcagatcc tgacagggaa      540
gaggaggaga tgagagactc tggcatgata ttttttttgt ccacttggt ggggccaggg      600
tcctctcccc tgcccaagaa tgtgcaaggc cagggcatgg gggcaaatat gaccagttt      660
tgggaaacacc gacaaaccca gccctggcgc tgagcctctc taccacaggt cagacggaca      720
gaaagacaaa tcacaggttc cgggatgagg acaccggctc tgaccaggag tttggggagc      780
ttcaggacat tgctgtgctt tggggattcc ctccacatgc tgcacgcgca tctcgcccc      840

```

```

aggggcactg cctggaagat tcaggagcct gggcggcctt cgcttactct cacctgcttc      900
tgagttgccc aggaggccac tggcagatgt cccggcggaag agaagagaca cattgttgga      960
agaagcagcc catgacagcg ccccttcctg ggactcgccc tcatcctctt cctgctcccc     1020
ttcctgggggt gcagcctaaa aggacctatg tcctcacacc attgaaacca ctagttctgt     1080
ccccccagga aacctgggtg tgtgtgtgtg agtgggtgac cttcctccat cccctgggtcc     1140
ttcccttccc ttcccgaggc acagagagac agggcaggat ccacgtgccc attgtggagg     1200
cagagaaaag agaaagtgtt ttatatacgg tactttattta atatcccttt ttaattagaa     1260
attagaacag ttaatttaat taaagagtag ggtttttttt cagtattctt gggttaatatt     1320
taatttcaac tatttatgag atgtatcttt tgctctctct tgctctctta tttgtaccgg     1380
tttttgtata taaaattcat gtttccaatc tctctctccc tgatcgggtga cagtcactag     1440
cttatcttga acagatattt aattttgcta acactcagct ctgccctccc cgatcccttg     1500
gctccccagc acacattcct ttgaaagagg gtttcaatat acatctacat actatatata     1560
tattgggcaa cttgtatttg tgtgtatata tatatatata tgtttatgta tatatgtgat     1620
cctgaaaaaa taaacatcgc tattctgttt tttatatgtt caaaccaaac aagaaaaaat     1680
agagaattct acatactaaa tctctctcct tttttaattt taatatttgt tatcatttat     1740
ttattggtgc tactgtttat ccgtaataat tgtggggaaa agatattaac atcacgtctt     1800
tgtctctagt gcagtttttc gagatattcc gtagtacata tttattttta aacaacgaca     1860
aagaaataca gatatatctt aaaaaaaaaa aa                                     1892

```

<210> 19  
 <211> 249  
 <212> RNA  
 <213> Homo sapiens

```

<400> 19
ccgggcucau ggacggguga ggcggcggug ugcgcagaca gugcuccagc gcgcgcgcuc      60
cccagcccug gcccggccuc gggccgggag gaagaguagc ucgccgaggc gccgaggaga     120
gcggggccgc ccacagcccg agccggagag ggacgcgagc cgcgcgcccc ggucgggccu     180
ccgaaaccau gaacuucug cugucuuggg ugcauuggag ccuugccuug cugcucuacc     240
uccaccaug                                     249

```

<210> 20  
 <211> 15  
 <212> RNA  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: Motif

<220>

<221> misc\_feature

<222> (1)..(15)

<223> n = a, t, g or c

<400> 20

nauuuauuuu uuuuan

15

<210> 21

<211> 49

<212> DNA

<213> Homo sapiens

<400> 21

ccgccagatt tgaatcgcg gacccgttgg cagaggtggc ggcggcggc

49

<210> 22

<211> 1141

<212> DNA

<213> Homo sapiens

<400> 22

ggcctctggc cggagctgcc tgggtcccaga gtggctgcac cacttccagg gtttattccc 60

tgggtgccacc agccttccctg tgggcccctt agcaatgtct taggaaagga gatcaacatt 120

ttcaaattag atgtttcaac tgtgctcctg ttttgtcttg aaagtggcac cagaggtgct 180

tctgcctgtg cagcgggtgc tgctggtaac agtggctgct tctctctctc tctctctttt 240

ttgggggctc atttttgctg ttttgattcc cgggcttacc aggtgagaag tgaggaggga 300

agaaggcagt gtcccttttg ctagagctga cagctttgtt cgcgtgggca gagccttcca 360

cagtgaatgt gtctggacct catgttggtg aggtgtcac agtcctgagt gtggacttgg 420

caggtgcctg ttgaatctga gctgcagggt ccttatctgt cacacctgtg cctcctcaga 480

ggacagtttt tttgttggtg tgtttttttt tttttttttt ttggtagatg catgacttgt 540

gtgtgatgag agaatggaga cagagtccct ggctcctcta ctgtttaaca acatggcttt 600

cttattttgt ttgaattgtt aattcacaga atagcacaaa ctacaattaa aactaagcac 660

aaagccattc taagtcattg gggaaacggg gtgaacttca ggtggatgag gagacagaat 720

agagtgatag gaagcgtctg gcagatactc cttttgccac tgctgtgtga ttagacaggc 780

ccagtgagcc gcggggcaca tgctggccgc tcctccctca gaaaaaggca gtggcctaaa 840

tcctttttta atgacttggc tcgatgctgt gggggactgg ctgggctgct gcaggccgtg 900

tgtctgtcag cccaaccttc acatctgtca cgttctccac acgggggaga gacgcagtcc 960

gcccaggtcc ccgctttctt tggaggcagc agctcccgca gggctgaagt ctggcgtaag 1020

atgatggatt tgattcgccc tcctccctgt catagagctg caggggtggat tgttacagct 1080  
 tcgctggaaa cctctggagg tcatctcggc tggtcctgag aaataaaaag cctgtcattt 1140  
 c 1141

<210> 23  
 <211> 247  
 <212> DNA  
 <213> Homo sapiens

<400> 23  
 ccccggcgca gcgcggccgc agcagcctcc gcccccgca cgggtgtgagc gcccgcgcgc 60  
 gccgaggcgc ccggagtccc gagctagccc cggcggccgc cgccgcccag accggacgac 120  
 aggccacctc gtcggcgctcc gcccgagtcc ccgcctcgcc gccaacgcca caaccaccgc 180  
 gcacggcccc ctgactccgt ccagtattga tcgggagagc cggagcgagc tcttcggggga 240  
 gcagcag 247

<210> 24  
 <211> 1716  
 <212> DNA  
 <213> Homo sapiens

<400> 24  
 tgaccacgga ggatagtatg agccctaaaa atccagactc tttcgatacc caggaccaag 60  
 ccacagcagg tcctccatcc caacagccat gccgcatta gctcttagac ccacagactg 120  
 gttttgcaac gtttacaccg actagccagg aagtacttcc acctcgggca cattttggga 180  
 agttgcattc ctttgtcttc aaactgtgaa gcatttacag aaacgcatcc agcaagaata 240  
 ttgtcccttt gagcagaaat ttatctttca aagagggtata tttgaaaaaa aaaaaaaaag 300  
 tataatgtgag gattttttatt gattggggat cttggagttt ttcattgtcg ctattgattt 360  
 ttacttcaat gggctcttcc aacaaggaag aagcttgctg gtagcacttg ctaccctgag 420  
 ttcattccagg cccaactgtg agcaaggagc acaagccaca agtcttcagc aggatgcttg 480  
 attccagtgg ttctgcttca aggcttccac tgcaaaacac taaagatcca agaaggcctt 540  
 catggcccca gcaggccgga tcggtactgt atcaagtcac ggcagggtaca gtaggataag 600  
 ccactctgtc ccttcctggg caaagaagaa acggagggga tgaattcttc cttagactta 660  
 cttttgtaaa aatgtcccca cggctacttac tccccactga tggaccagtg gtttccagtc 720  
 atgagcggtta gactgacttg tttgtcttcc attccattgt tttgaaactc agtatgccgc 780  
 ccctgtcttg ctgtcatgaa atcagcaaga gaggatgaca catcaaataa taactcggat 840  
 tccagcccac attggattca tcagcatttg gaccaatagc ccacagctga gaatgtggaa 900  
 tacctaagga taacaccgct tttgttctcg caaaaacgta tctcctaatt tgaggctcag 960

atgaaatgca tcagggtcctt tggggcatag atcagaagac tacaaaaatg aagctgctct 1020  
 gaaatctcct ttagccatca cccaaccccc ccaaaattag tttgtgttac ttatggaaga 1080  
 tagttttctc cttttacttc acttcaaaaag ctttttactc aaagagtata tgttccctcc 1140  
 aggtcagctg cccccaacc cctccttac gctttgtcac acaaaaagtg tctctgcctt 1200  
 gagtcattcta ttcaagcact tacagctctg gccacaacag ggcattttac aggtgcgaat 1260  
 gacagtagca ttatgagtag tgtgaattca ggtagtaaat atgaaactag ggtttgaaat 1320  
 tgataatgct ttcacaacat ttgcagatgt tttagaagga aaaaagttcc ttcctaaaat 1380  
 aatttctcta caattggaag attggaagat tcagctagtt aggagcccat tttttcctaa 1440  
 tctgtgtgtg ccctgtaacc tgactgggta acagcagtc tttgtaaaca gtgtttttaa 1500  
 ctctcctagt caatatccac cccatccaat ttatcaagga agaaatgggt cagaaaaatat 1560  
 tttcagccta cagttatggt cagtcacaca cacatacaaa atgttccttt tgctttttaa 1620  
 gtaatttttg actcccagat cagtcagagc ccctacagca ttgttaagaa agtatttgat 1680  
 ttttgtctca atgaaaataa aactatatcc atttcc 1716

<210> 25  
 <211> 160  
 <212> DNA  
 <213> Homo sapiens

<400> 25  
 tataaaagct gggccggcgc gggccgggccc attcgcgacc cggaggtgcg cgggcgcggg 60  
 cgagcagggg ctccgggtgg gcggcgcgac gccccgcgca ggctggaggc cgccgaggct 120  
 cgccatgccg ggagaactct aactccccca tggagtcggc 160

<210> 26  
 <211> 1306  
 <212> DNA  
 <213> Homo sapiens

<400> 26  
 tgaggcgcgc ggctgtggga ccgccctggg ccagcctccg gcggggaccc agggagtggg 60  
 ttggggctgc cggtatctga ggcttgccca gaccgtgcga gccaggacta ggagattccg 120  
 gtgcctcctg aaagcctggc ctgctccgcg tgtccctcc cttcctctgc gccggacttg 180  
 gtgcgtctaa gatgaggggg ccaggcgggt gcttctccct gcgaggaggg gagaattctt 240  
 ggggctgagc tgggagcccc gcaactctag tatttaggat aacttgtgcc ttggaaatgc 300  
 aaactcaccg ctccaatgcc tactgagtag ggggagcaaa tcgtgccttg tcattttatt 360  
 tggaggtttc ctgcctcctt cccgaggcta cagcagaccc ccatgagaga aggaggggag 420

caggccccgtg gaggaggggg gctcagggag ctgagatccc gacaagcccc ccagccccag 480  
 ccgctcctcc acgcctgtcc ttagaaaagg gtggaaacat agggacttgg ggcttggaac 540  
 ctaaggttgt tccctagtcc tacatgaagg tggaggtctc tagttccacg cctctccac 600  
 ctccctccgc acacacccca ccagcctgc tataggctgg ctttcccttg gggctggaac 660  
 tcaactgcgat ggggtcacca ggtgaccagt ggagccccca ccccgagtca gaccagaaag 720  
 ctaggtcgtg ggtcagctct gaggatgtat acccctgggt ggagagggag acctagagat 780  
 ctggctgtgg ggcgggcatg ggggggtgaag ggccactggg accctcagcc ttgtttgtac 840  
 tgtatgcctt cagcattgcc taggaacacg aagcacgata agtccatcca gagggaccgg 900  
 agttatgaca agcttcccaa atattttgct ttatcagccg atatcaacac ttgtatctgg 960  
 cctctgtgcc cagcagtgcc ttgtgcaatg tgaatgtacc gtctctgcta aaccaccatt 1020  
 ttattttggtt ttgttttgtt tggttttctc ggatacttgc caaaatgaga ctctccgtcg 1080  
 gcagctgggg gaaggggtctg agactctctt tcccttttgg tttgggatta cttttgatcc 1140  
 tgggggacca atgaggtgag ggggggtctc ctttgccctc agctttccca gccctccggc 1200  
 ctgggctgcc cacaaggctt ctccccaga ggccctggct cctggtcggg aaggaggtg 1260  
 cctcccgcca acgcatcact ggggctggga gcagggaagg gaattc 1306

<210> 27  
 <211> 216  
 <212> DNA  
 <213> Homo sapiens

<400> 27  
 agcgagagcg cccccgagca gcgcccgcgc cctccgcgcc ttctccgccg ggacctcgag 60  
 cgaaaagacgc ccgcccgcgc ccagccctc gcctccctgc ccaccgggca caccgcgccg 120  
 ccaccccgac ccgctgcgc acggcctgtc cgctgcacac cagcttggtg gcgtcttcgt 180  
 cgccgcgctc gcccggggt actcctgcgc gccaca 216

<210> 28  
 <211> 687  
 <212> DNA  
 <213> Homo sapiens

<400> 28  
 taaatgctac ctgggtttcc agggcacacc tagacaaaca rgggagaaga gtgtcagaat 60  
 cagaatcatg gagaaaatgg gcgggggtgg tgtgggtgat gggactcatt gtagaaagga 120  
 agccttgctc attcttgagg agcatthaagg tatttcgaaa ctgccaaagg tgctgggtgcg 180  
 gatggacact aatgcagcca cgattggaga atactttgct tcatagtatt ggagcacatg 240  
 ttactgcttc attttgagc ttgtggagtt gatgactttc tgttttctgt ttgtaaatta 300

ttgctaagc atattttctc taggcttttt tccttttggg gttctacagt cgtaaaagag 360  
 ataataagat tagttggaca gtttaaagct tttattcgtc ctttgacaaa agtaaatggg 420  
 agggcattcc atcccttctt gaagggggac actccatgag tgtctgtgag aggcagctat 480  
 ctgcactcta aactgcaaac agaaatcagg tgttttaaga ctgaatgttt tatttatcaa 540  
 aatgtagctt ttggggaggg aggggaaatg taatactgga ataatttgta aatgatttta 600  
 attttatatt cagtgaagag attttattta tggaattaac catttaataa agaaatatat 660  
 acctaaaaaa aaaaaaaaaa aaaaaaa 687

<210> 29  
 <211> 310  
 <212> DNA  
 <213> Homo sapiens

<400> 29  
 cggccccaga aaacccgagc gagtaggggg cggcgcgcag gagggaggag aactggggggc 60  
 gggggaggct ggtgggtgtc ggggggtggag atgtagaaga tgtgacgccg cggcccggcg 120  
 ggtgccagat tagcggacgg ctgccgcggg ttgcaacggg atcccggggc ctgcagcttg 180  
 ggaggcggct ctccccaggc ggcgtccgcg gagacacca tccgtgaacc ccagggtccc 240  
 ggccgcccgc tcgccgcgca ccagggggccg gcggacagaa gagcggccga gcggctcgag 300  
 gctggggggac 310

<210> 30  
 <211> 5882  
 <212> DNA  
 <213> Homo sapiens

<400> 30  
 ctgctaagag ctgattttta tggccacatc taatctcatt tcacatgaaa gaagaagtat 60  
 attttagaaa ttgtttaatg agagtaaaag aaaataaatg tgtatagctc agtttggata 120  
 attggtcaaa caatttttta tccagtagta aaatatgtaa ccattgtccc agtaaagaaa 180  
 aataacaaaa gttgtaaaat gtatatctc ccttttatat tgcattctgt gttaccagct 240  
 gaagcttacc tagagcaatg atctttttca cgcatttgct ttattcgaaa agaggctttt 300  
 aaaatgtgca tgtttagaaa caaaatttct tcatggaaat catatacatt agaaaatcac 360  
 agtcagatgt ttaatcaatc caaaatgtcc actatttctt atgtcattcg ttagtctaca 420  
 tgtttctaaa catataaatg tgaatttaat caattccttt catagtttta taattctctg 480  
 gcagttcctt atgatagagt ttataaaaca gtcctgtgta aactgctgga agttcttcca 540



cagtcaggtc	aatthttgtca	aacccttctc	tgtacccata	cagcagcagc	ctagcaactc	600
tgctggatgat	gggagttgta	ttttcagttc	tcgccaggtc	attgagatcc	atccactcac	660
atcttaagca	ttcttcctgg	caaaaattta	tggtgaatga	atatggcttt	aggcggcaga	720
tgatatacat	atctgacttc	ccaaaagctc	caggatttgt	gtgctggtgc	cgaataactca	780
ggacggacct	gaattctgat	tttataccag	tctcttcaaa	aactttctcg	accgctgtgt	840
ctcctacgta	aaaaaagaga	tgtacaaatc	aataataatt	acacttttag	aaactgtatc	900
atcaaagatt	ttcagttaaa	gtagcattat	gtaaaggctc	aaaacattac	cctaacaaag	960
taaagttttc	aatacaaatt	ctttgccttg	tgatatacaa	gaaatcccaa	aatatthttct	1020
taccactgta	aattcaagaa	gcttttgaaa	tgctgaatat	ttctttggct	gctacttgga	1080
ggcttatcta	cctgtacatt	tttgggggtca	gctctthttta	acttcttgct	gctctthtttc	1140
ccaaaaggta	aaaatataga	ttgaaaagtt	aaaacatttt	gcatggctgc	agttcctttg	1200
tttcttgaga	taagattcca	aagaacttag	attcatttct	tcaacaccga	aatgctggag	1260
gtgtttgatc	agttttcaag	aaacttgga	tataaataat	tttataattc	aacaaagggt	1320
ttcacatttt	ataaggttga	tttttcaatt	aatgcaaatt	ttgtgtggca	ggattthttat	1380
tgccattaac	atathttttgt	ggctgctttt	tctacacatc	cagatgggtcc	ctctaactgg	1440
gctttctcta	atthttgtgat	gttctgtcat	tgtctcccaa	agtatttagg	agaagccctt	1500
taaaaagctg	ccttcctcta	ccactttgct	ggaaagcttc	acaattgtca	cagacaaaaga	1560
tttttgttcc	aataactcgtt	ttgcctctat	ttttcttggt	tgtcaaatag	taaatgatat	1620
ttgcccttgc	agtaattcta	ctggtgaaaa	acatgcaaag	aagagggaagt	cacagaaaca	1680
tgtctcaatt	cccatgtgct	gtgactgtag	actgtcttac	catagactgt	cttaccatc	1740
ccctggatat	gctcttggtt	tttccctcta	atagctatgg	aaagatgcat	agaaagagta	1800
taatgtthta	aaacataagg	cattcatctg	ccattthttca	attacatgct	gacttccctt	1860
acaattgaga	tttgcccata	ggttaaacad	ggttagaaac	aactgaaagc	ataaaagaaa	1920
aatctaggcc	gggtgcagtg	gctcatgcct	atattccctg	cactttggga	ggccaaagca	1980
ggaggatcgc	ttgagcccag	gagttcaaga	ccaacctggg	gaaaccccg	ctctacaaaa	2040
aaacacaaaa	aatagccagg	catggtggcg	tgtacatgtg	gtctcagata	cttggggaggc	2100
tgagggtggga	gggttgatca	cttgaggctg	agagggtcaag	gttgacagtga	gccataatcg	2160
tgccactgca	gtccagccta	ggcaacagag	tgagactttg	tctcaaaaaa	agagaaattt	2220
tccttaataa	gaaaagtaat	ttttactctg	atgtgcaata	catttggttat	taaatttatt	2280
atttaagatg	gtagcactag	tcttaaatgt	tataaaatat	cccctaacad	gtttaaatgt	2340
ccattthttat	tcattatgct	ttgaaaaata	attatggggg	aatacatggt	tgthtattaaa	2400

tttattatta	aagatagtag	cactagtctt	aaatttgata	taacatctcc	taacttgttt	2460
aaatgtccat	ttttattctt	tatgcttgaa	aataaattat	ggggatccta	tttagctctt	2520
agtaccacta	atcaaaagtt	cggcatgtag	ctcatgatct	atgctgtttc	tatgtcgtgg	2580
aagcaccgga	tgggggtagt	gagcaaactc	gccctgctca	gcagtcacca	tagcagctga	2640
ctgaaaatca	gcactgcctg	agtagttttg	atcagtttaa	cttgaatcac	taactgactg	2700
aaaattgaat	gggcaaataa	gtgcttttgt	ctccagagta	tgcgggagac	ccttccacct	2760
caagatggat	atttcttccc	caaggatttc	aagatgaatt	gaaattttta	atcaagatag	2820
tgtgctttat	tctgttgtat	tttttattat	tttaataatac	tgtaagccaa	actgaaataa	2880
catttgctgt	tttatagggt	tgaagaacat	aggaaaaact	aagaggtttt	gtttttattt	2940
ttgctgatga	agagatatgt	ttaaatatgt	tgtattgttt	tgtttagtta	caggacaata	3000
atgaaatgga	gtttatattt	gttatttcta	ttttgttata	tttaataata	gaattagatt	3060
gaaataaaat	ataatgggaa	ataatctgca	gaatgtgggt	ttcctggtgt	ttcctctgac	3120
tctagtgcac	tgatgatctc	tgataaggct	cagctgcttt	atagttctct	ggctaattgca	3180
gcagatactc	ttcctgccag	tggttaatacg	attttttaag	aaggcagttt	gtcaatttta	3240
atcttggtga	tacctttata	ctcttaggggt	attattttat	acaaaagcct	tgaggattgc	3300
attctatttt	ctatatgacc	ctcttgatat	ttaaaaaaca	ctatggataa	caattcttca	3360
tttacctagt	attatgaaag	aatgaaggag	ttcaaacaaa	tgtgtttccc	agttaactag	3420
ggtttactgt	ttgagccaat	ataaatgttt	aactgtttgt	gatggcagta	ttcctaaagt	3480
acattgcatg	ttttcctaaa	tacagagttt	aaataatttc	agtaattctt	agatgattca	3540
gcttcatcat	taagaatatc	ttttgtttta	tgttgagtta	gaaatgcctt	catatagaca	3600
tagtctttca	gacctctact	gtcagttttc	atttctagct	gctttcaggg	ttttatgaat	3660
tttcaggcaa	agctttaatt	tatactaagc	ttaggaagta	tggctaattgc	caacggcagt	3720
ttttttcttc	ttaattccac	atgactgagg	catatatgat	ctctgggtag	gtgagttggt	3780
gtgacaacca	caagcacttt	tttttttttt	aaagaaaaaa	aggtagtgaa	tttttaatca	3840
tctggacttt	aagaaggatt	ctggagtata	cttaggcctg	aaattatata	tatttggtct	3900
ggaaatgtgt	ttttcttcaa	ttacatctac	aagtaagtac	agctgaaatt	cagaggaccc	3960
ataagagttc	acatgaaaaa	aatcaattca	tttgaaaagg	caagatgcag	gagagaggaa	4020
gccttgcaaa	cctgcagact	gcttttttgc	caatatagat	tgggtaaggc	tgcaaaacat	4080
aagcttaatt	agctcacatg	ctctgctctc	acgtggcacc	agtggatagt	gtgagagaat	4140
taggctgtag	aacaaatggc	cttctctttc	agcattcaca	ccactacaaa	atcatctttt	4200

```

atatcaacag aagaataagc ataaactaag caaaagggtca ataagtacct gaaaccaaga 4260
ttggctagag atatatctta atgcaatcca ttttctgatg gattggttacg agttggctat 4320
ataatgtatg tatgggtatgt tgatttgtgt aaaagtttta aaaatcaagc ttaagtaca 4380
tggaacatgtt taaataaaat atttaaagac aatttagaaa attgccttaa tatcattggt 4440
ggctaaatag aataggggac atgcatatta aggaaaaggc catggagaaa taatattggt 4500
atcaaacaaa tacattgatt tgtcatgata cacattgaat ttgatccaat agtttaagga 4560
ataggtagga aaatttggtt tctatttttc gatttctgtt aaatcagtga cataaataat 4620
tcttagctta ttttatatgt ccttgtctta aatactgagc tcagtaagtt gtgttagggg 4680
attattttctc agttgagact ttcttatatg acattttact atgttttgac ttcctgacta 4740
ttaaaaataa atagtagaaa caattttcat aaagtgaaga attatataat cactgcttta 4800
taactgactt tatttatatgt atttcaaagt tcattttaaag gctactatct atcctctgtg 4860
atggaatggc caggaatttg ttttctcata gttaaattcc aacaacaata ttagtcgtat 4920
ccaaaataac ctttaatgct aaactttact gatgtatatc caaagcttct ccttttcaga 4980
cagattaatc cagaagcagt cataaacaga agaatagggtg gtatgttcct aatgatatta 5040
tttctactaa tggaataaac tgtaatatta gaaattatgc tgctaattat atcagctctg 5100
aggtaatttc tgaaatgttc agactcagtc ggaacaaatt ggaaaattta aatttttatt 5160
cttagctata aagcaagaaa gtaaacacat taatttcctc aacattttta agccaattaa 5220
aaatataaaa gatacacacc aatatcttct tcaggctctg acaggcctcc tggaaacttc 5280
cacatatgtt tcaactgcag tataaagtca gaaaataaag ttaacataac tttcactaac 5340
acacacatat gtagatttca caaaatccac ctataattgg tcaaagtggc tgagaatata 5400
tttttttaga attgcatgca aaatttttct agcttccatc ctttctccct cgtttcttct 5460
ttttttgggg gagctggtaa ctgatgaaat cttttccac cttttctctt caggaaatat 5520
aagtggtttt gtttggttta cgtgatacat tctgtatgaa tgaaacattg gagggaaaca 5580
tctactgaat ttctgtaatt taaaatattt tgctgctagt taactatgaa cagatagaag 5640
aatcttacag atgctgctat aaataagtag aaaatataaa tttcatcact aaaatagct 5700
attttaaaat ctatttctta tattgtatgt ctaatcagat gtattactct tattatttct 5760
attgtatgtg ttaatgattt tatgtaaaaa tgtaattgct tttcatgagt agtatgaata 5820
aaattgatta gtttgtgttt tcttgtctcc cgaaaaaaaa aaaaaaaaaa aaaaaaaaaa 5880
aa 5882

```

<210> 31  
<211> 310

<212> DNA  
 <213> Homo sapiens

<400> 31  
 cggccccaga aaacccgagc gagtaggggg cggcgcgcag gagggaggag aactggggggc 60  
 gcgggagggt ggtgggtgtc ggggggtggag atgtagaaga tgtgacgccg cggcccggcg 120  
 ggtgccagat tagcggacgg ctgcccgcgg ttgcaacggg atcccggggc ctgcagcttg 180  
 ggaggcgggt ctccccaggc ggcgtccgcg gagacaccca tccgtgaacc ccagggtcccg 240  
 ggccgcggcg tcgcccgcga ccagggggcg gcggacagaa gagcggccga gctggctcgag 300  
 gctggggggac 310

<210> 32  
 <211> 3212  
 <212> DNA  
 <213> Homo sapiens

<400> 32  
 tgaggggcgcc aggcaggcgg gcgccaccgc caccgcgcag gagggcggag ccggccccag 60  
 gtgctccccct gacagtccct cctctccgga gcattttgat accagaaggg aaagcttcat 120  
 tctccttggt gttggttggt ttttcctttg ctctttcccc ctccatctc tgacttaagc 180  
 aaaagaaaaa gattacccaa aaactgtctt taaaagagag agagagaaaa aaaaaatagt 240  
 atttgcataa ccctgagcgg tgggggagga gggttgtgct acagatgata gaggatttta 300  
 taccccaata atcaactcgt ttttatatta atgtacttgt ttctctgttg taagaatagg 360  
 cattaacaca aaggaggcgt ctggggagag gattagggtc catcctttac gtgtttaaaa 420  
 aaaagcataa aaacatttta aaacataga aaaattcagc aaaccatttt taaagtagaa 480  
 gagggtttta ggtagaaaaa catattcttg tgcttttcct gataaagcac agctgtagtg 540  
 gggttctagg catctctgta ctttgcttgc tcatatgcat gtagtcactt tataagtcac 600  
 tgtatgttat tatattccgt aggtagatgt gtaacctctt caccttattc atggctgaag 660  
 tcacctcttg gttacagtag cgtagcgtgg ccgtgtgcat gtcctttgcg cctgtgacca 720  
 ccacccaac aaaccatcca gtgacaaaacc atccagtga ggtttgtcgg gcaccagcca 780  
 gcgtagcagg gtccggaaag gccacctgtc ccactcctac gatacgctac tataaagaga 840  
 agacgaaata gtgacataat atattctatt ttataactct tcctatTTTT gtagtgacct 900  
 gtttatgaga tgctggTTTT ctacccaacg gccctgcagc cagctcacgt ccaggttcaa 960  
 cccacagcta cttgggtttgt gttcttcttc atattctaaa accattccat ttccaagcac 1020  
 tttcagtcca ataggtgtag gaaatagcgc tgtttttggt gtgtgtgcag ggagggcagt 1080  
 tttctaattg aatgggtttg gaatatccat gtacttgttt gcaagcagga ctttgaggca 1140

```

agtgtgggcc actgtggtgg cagtggaggt ggggtgtttg ggaggctgcg tgccagtcaa 1200
gaagaaaaaag gtttgcattc tcacattgcc aggatgataa gttcctttcc ttttctttaa 1260
agaagttgaa gtttaggaat cctttggtgc caactggtgt ttgaaagtag ggacctcaga 1320
ggtttaccta gagaacaggt ggtttttaag ggttatctta gatgtttcac accggaaggt 1380
ttttaaacac taaaatatat aatttatagt taaggctaaa aagtatatatt attgcagagg 1440
atgttcataa ggccagtatg atttataaat gcaatctccc cttgatttaa acacacagat 1500
acacacacac acacacacac acacacaaac cttctgcctt tgatgttaca gatttaatac 1560
agtttatatt taaagataga tccttttata ggtgagaaaa aaacaatctg gaagaaaaaa 1620
accacacaaa gacattgatt cagcctgttt ggcgtttccc agagtcactc gattggacag 1680
gcatgggtgc aaggaaaatt agggactca acctaagttc ggttccgatg aattcttctc 1740
ccctgcccct tcctttaaaa aacttagtga caaaatagac aatttgacac tcttggtctat 1800
gtaattcttg taatttttat ttaggaagtg ttgaaggag gtggcaagag tgtggaggct 1860
gacgtgtgag ggaggacagg cgggaggagg tgtgaggagg aggctcccga ggggaagggg 1920
cggtgccac accggggaca ggccgcagct ccattttctt attgcgctgc taccgttgac 1980
ttccaggcac ggtttggaat tattcacatc gcttctgtgt atctctttca cattgtttgc 2040
tgctattgga ggatcagttt tttgttttac aatgtcatat actgccatgt actagtttta 2100
gttttctctt agaacattgt attacagatg ccttttttgt agtttttttt ttttttatgt 2160
gatcaatttt gacttaatgt gattactgct ctattccaaa aaggttgctg tttcacaata 2220
cctcatgctt cacttagcca tgggtggacc agcggggcagg ttctgcctgc tttggcgggc 2280
agacacgcgg gcgcgatccc acacaggctg gcgggggccc gcccgcaggc cgcgtgctg 2340
agaaccgcgc cgggtgtccc agagaccagg ctgtgtccct cttctcttcc ctgcgcctgt 2400
gatgctgggc acttcatctg atcgggggcg tagcatcata gtagttttta cagctgtgtt 2460
attctttgcg ttagctatg gaagttgcat aattattatt attattatta taacaagtgt 2520
gtcttacgtg ccaccacggc gttgtacctg taggactctc attcgggatg attggaatag 2580
cttctggaat ttgttcaagt tttgggtatg ttaaatctgt tatgtactag tgttctgttt 2640
gttattgttt tgtaattac accataatgc taatttaaag agactccaaa tctcaatgaa 2700
gccagctcac agtgctgtgt gcccgggtca cctagcaagc tgccgaacca aaagaatttg 2760
caccocgctg cgggcccacg tggttggggc cctgccctgg cagggtcatc ctgtgctcgg 2820
aggccatctc gggcacaggc ccaccccgcc ccaccctcc agaacacggc tcacgcttac 2880
ctcaaccatc ctggctgctg cgtctgtctg aaccacgcgg gggccttgag ggacgctttg 2940

```

```

tctgtcgtga tggggcaagg gcacaagtcc tggatgttgt gtgtatcgag aggccaaagg 3000
ctggtggcaa gtgcacgggg cacagcggag tctgtcctgt gacgcgcaag tctgagggtc 3060
tgggcggcgg gcggtctgggt ctgtgcattt ctggttgcac cgcggcgctt cccagcacca 3120
acatgtaacc ggcattgttt cagcagaaga caaaaagaca aacatgaaag tctagaaata 3180
aaactggtaa aacccccaaaa aaaaaaaaaa aa 3212

```

```

<210> 33
<211> 1043
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (409)..(444)
<223> n = a, t, g or c

```

```

<400> 33
gcaccgcggc gagcttggct gcttctgggg cctgtgtggc cctgtgtgtc ggaaagatgg 60
agcaagaagc cgagcccgag gggcggccgc gaccctctg accgagatcc tgctgctttc 120
gcagccagga gcaccgtccc tccccggatt agtgcgtacg agcgcgccagt gccctggccc 180
ggagagtgga atgatccccg aggccagggg cgtcgtgctt ccgcgcgccc cgtgaaggaa 240
actggggagt cttgaggggac ccccgactcc aagcgcgaaa accccggatg gtgaggagca 300
ggtaactggc cggcagcgag cggtcacttt tgggtctggg ctctgacggg gtcccctcta 360
tcgctgggtc ccagcctctg cccgttcgca gcctttgtgc ggttcgtgnc tgggggctcg 420
gggcgcgggg cgcggggcat gggncacgtg gctttgcgga ggttttgttg gactggggct 480
agacagtccc cgccaggagg gagggcgagg tttcggacgg ctctcgcggc ggtgggggtg 540
ggggtgggtc ggaggtctcc gcgggagttc agggtaaagg tcacggggcc ggggctgcgg 600
gccgcttcgg cgcgggaggt ccggatgatc gcagtgcctg tcgggtcact agtgtgaacg 660
ctgcgcgtag tctgggcggg attgggccgg ttcagtgggc aggttgactc agcttttcct 720
cttgagctgg tcaagttcag acacgttcgg aaactgcagt aaaaggagtt aagtcctgac 780
ttgtctccag ctggggctat ttaaaccatg cattttccca gctgtgttca gtggcgattg 840
gagggtagac ctgtgggcac ggacgcacgc cactttttct ctgctgatcc aggttaagcac 900
cgacttgctt gtagcttttag ttttaactgt tgtttatgtt ctttatatat gatgtathtt 960
ccacagatgt ttcattgatt ccagttttca tcgtgtcttt tttttccttg taggcaaagt 1020
tgcaatacca acatgtctgt acc 1043

```

<210> 34  
 <211> 1153  
 <212> DNA  
 <213> Homo sapiens

<400> 34  
 tagttgacct gtctataaga gaattatata tttctaacta tataacccta ggaatttaga 60  
 caacctgaaa tttattcaca tatatcaaag tgagaaaatg cctcaattca catagatttc 120  
 ttctcttttag tataattgac ctactttggt agtggaatag tgaatactta ctataatttg 180  
 acttgaatat gtagctcatc ctttacacca actcctaatt ttaaataatt tctactctgt 240  
 cttaaagtag aagtacttgg tttttttttt cttaaatatg tatatgacat ttaaagttaa 300  
 cttattatatt tttttgagac cgagtcttgc tctgttacc aggctggagt gcagtgggtg 360  
 atcttggtc actgcaagct ctgccctccc cgggttcgca ccattctcct gcctcagcct 420  
 cccaattagc ttggcctaca gtcactctgcc accacacctg gctaattttt tgtactttta 480  
 gtagagacag ggtttcaccg tgtagccag gatggtctcg atctcctgac ctctgatcc 540  
 gccacctcg gcctccaaa gtgctgggat tacaggcatg agccaccgtg ctctccagcc 600  
 taggcaacag agtgagactc tgtctccaaa aaaaaaaaaa aaaaaggagg actataacac 660  
 cccagggaa agggacaggt gggacattct tattcttaat ttaaataaat tgacagggga 720  
 aagttgggcc actcttgagc ttgtgggtgc tcaccagggt gaccccaaaa aaagaagcct 780  
 tccacaaaac attaatttat ttcctaata taccgcctc tgtgagttaa gggataatgc 840  
 atcaggactc ttgcaaccag acaaaattat ttaaaaacgc cacttggggg ggaggcgggt 900  
 ccctcctggg gattcgctt tgtgggagag aaaactgcac agacttgggc aaataatggt 960  
 ttttgtcacc ccaaaacgta ttcgogagac atttcattag aacgaagctt taccctaata 1020  
 ttgaactccc catttaaaca gtttccacac acacttaggg agatttttcc ctctgtgagt 1080  
 tccgcagaac aatagttgga cgggaataga accctgaaac actttagttc accacgaact 1140  
 attatagggc ggg 1153

<210> 35  
 <211> 334  
 <212> DNA  
 <213> Homo sapiens

<400> 35  
 tgactatcca gctctgagag acgggagttt ggagttgcc gctttacttt ggttgggttg 60  
 gggggggcgg cgggctgttt tgttcctttt cttttttaag agttgggttt tcttttttaa 120  
 ttatccaaac agtgggcagc ttctccccc acaccaagt atttgcacaa tatttgtgcg 180  
 gggatatggg gtgggttttt aaatctcgtt tctcttgag aagcacaggg atctcgttct 240

cctcattttt tgggggtgtg tggggacttc tcaggtcgtg tccccagcct tctctgcagt 300  
cccttctgcc ctgccgggcc cgtcgggagg cgcc 334

<210> 36  
<211> 543  
<212> DNA  
<213> Homo sapiens

<400> 36  
tagctcagga ccttggctgg gcctggctgt catgtaggtc aggaccttgg ctggacctgg 60  
aggccctgcc cagccctgct ctgcccagcc cagcaggggc tccaggcctt ggctggcccc 120  
acatcgcctt ttctccccg acacctcgt gcacttgtgt ccgaggagcg aggagcccct 180  
cgggcccctgg gtggcctctg ggcctttct cctgtctccg ccactccctc tggcggcgct 240  
ggccgtggct ctgtctctct gaggtgggtc gggcgccctc tgcccgcctc ctcccacacc 300  
agccaggctg gtctcctcta gcctgtttgt tgtgggggtg gggatatatt tgtaaccact 360  
gggccccag cccctctttt ggcacccctt gtcctgacct gttctcggca ccttaaatta 420  
ttagaccccg gggcagtcag gtgtccgga cccccgaagg caataaaaca ggagccgtga 480  
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 540  
aaa 543

<210> 37  
<211> 511  
<212> DNA  
<213> Homo sapiens

<400> 37  
gctcagcaag gggtcogtcc ttctctgtca ctgtctcttt tgccgtgtgt aattctgtct 60  
gcctctctgg gactctgcct gtctcactct ttctgtctgt gcctctcctc actcttgctc 120  
tttctgcctg aatcacagcc ctcagttttt ctgtcctcat gcatttgtct ttgtggctct 180  
ttccgtcttt ctgcccttga caccatcccc tctcccagtg cttccctct gcttccagat 240  
cgcttcatga cttaggcagg gaaacagagg tcagggcctc cttccaggct tccctctgca 300  
tcttactgag tatgcaggtc ggaagagcct cgggtcctgc ctccgcgggt ggcctagagc 360  
caaaggaagg cggagcccg cggggcgagg ttggccctta gggccacctc ataaagcctg 420  
gggcgagggg cacaacggcc ttgggaagga gccctgctgg ggccgtccag tccccagac 480  
ctcacaggct cagtcgcgga tctgcagtgt c 511

<210> 38  
<211> 458  
<212> DNA  
<213> Homo sapiens



```

<400> 38
tagtaggggac cagtgaccat cacatccctt caagagtcct gaagatcaag ccagttctcc      60
ttccctgcag agctttggcc attaccacct gacctcttgc tgccagctaa taagaagtgc      120
caagtggaca gtctggccac tgtcaaggca ggggaaggggc catgactttt ctgccctgcc      180
ctcagcctgt tgccctgcct cccaaacccc attagtctag ccttgtagct gttactgcaa      240
gtgtttcttc tggcttagtc tgttttctaa agccaggact attccctttc ctccccagga      300
atatgtgttt tcctttgtct taatcgatct ggtaggggag aaatggcgaa tgtcatacac      360
atgagatggg atatccttgc gatgtacaga atcagaagggt ggtttgacag catcataaac      420
aggctgactg gcaggaatga aaaaaaaaaa aaaaaaaaaa      458

```

```

<210> 39
<211> 270
<212> DNA
<213> Homo sapiens

```

```

<400> 39
ggggccgccc agagccgcag cgccgctcgc ccgccgcccc ccaccccgcc gccccgcccg      60
gcgaattgcg ccccgcgccc tcccctcgcg ccccgagac aaagaggaga gaaagtttgc      120
gcggccgagc gggcagggtga ggaggggtgag ccgcgcggag gggcccgccct cggccccggc      180
tcagcccccg cccgcgcccc cagcccgccg ccgcgagcag cgcccggaacc cccagcggc      240
ggccccgccc gccagcccc ccggcccgcc      270

```

```

<210> 40
<211> 751
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (535)..(739)
<223> n = a, t, g or c

```

```

<400> 40
taagcaggcc tccaacgccc ctgtggccaa ctgcaaaaaa agcctccaag ggtttcgact      60
ggtccagctc tgacatccct tcctggaaac agcatgaata aaacactcat cccatgggtc      120
caaattaata tgattctgct ccccccttct ccttttagac atggttgtgg gtctggaggg      180
agacgtgggt ccaaggctct catcccatcc tccctctgcc aggcactatg tgtctggggc      240
ttcgatcctt ggggtgcaggc agggctggga cacgcggctt ccctcccagt ccctgccttg      300
gcaccgtcac agatgccaaag caggcagcac ttagggatct ccagctggg ttagggcagg      360
gcctggaaat gtgcattttg cagaaacttt tgagggtcgt tgcaagactg tgtagcaggc      420

```

ctaccaggtc cctttcatct tgagagggac atggccctt gttttctgca gcttccacgc 480  
ctctgactc cctgcccctg gcaagtgtc ccatcgcccc cggtgccac catgnagctc 540  
cccgcacctg actccccca catccaaggg cagccctgga accagtgggc tagttccttg 600  
aaggaagccc cactcattcc tattaatccc tcagaattcc cggggggagc cttccctcct 660  
gaaccttggg aaaaaatggg gaacgagaaa aacccccgct tggagctgtg cgtttccagc 720  
ccctacttga gagncttttt tttggggggc g 751

<210> 41  
<211> 229  
<212> DNA  
<213> Homo sapiens

<400> 41  
cgcgccgggc ccggctcggc ccgacccggc tccgcgcggg caggcggggc ccagcgcact 60  
cggagcccga gcccgagccg cagccgcccg ctggggcgct tgggtcggcc tcgaggacac 120  
cggagagggg cgccacgccc ccgtggccgc agatttgaaa gaagccgaca ctaaaccacc 180  
aatatacaac aaggccatth tgtcaaacga gagtcagcct ttaacgaaa 229

<210> 42  
<211> 233  
<212> DNA  
<213> Homo sapiens

<400> 42  
tagcagagag tcctgagcca ctgccaacat ttcccttctt ccagttgcac tattctgagg 60  
gaaaatctga cacctaagaa atttactgtg aaaaagcatt ttaaaaagaa aaggtttttag 120  
aatatgatct attttatgca tattgtttat aaagacacat ttacaattta cttttaatat 180  
taaaaattac catattatga aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaa 233

<210> 43  
<211> 349  
<212> DNA  
<213> Homo sapiens

<400> 43  
ggcacgaggg gcgagaggaa gcagggagga gagtgatttg agtagaaaag aaacacagca 60  
ttccaggctg gcccacctc tatattgata agtagccaat gggagcgggt agccctgac 120  
cctggccaat ggaaactgag gtaggcgggt catcgcgctg gggctctgtag tctgagcgct 180  
acccggttgc tgctgcccga ggaccgcgga gtcggacgca ggcagaccat gtggaccctg 240  
gtgagctggg tggccttaac agcagggctg gtggctggaa cgcggtgcc agatgggtcag 300  
ttctgccctg tggcctgctg cctggacccc ggaggagcca gctacagct 349

<210> 44  
 <211> 337  
 <212> DNA  
 <213> Homo sapiens

<400> 44  
 tgagggacag tactgaagac tctgcagccc tcgggacccc actcggaggg tgcctctctgc 60  
 tcaggcctcc ctagcacctc cccctaacca aattctccct ggacccatt ctgagctccc 120  
 catcaccatg ggaggtgggg cctcaatcta aggccttccc tgtcagaagg gggttgtggc 180  
 aaaagccaca ttacaagctg ccatcccctc cccgtttcag tggaccctgt ggccaggtgc 240  
 ttttccctat ccacaggggt gtttgtgtgt gtgcgcgtgt gcgtttcaat aaagtttgta 300  
 cactttcaaa aaaaaaaaaa aaaaaaaaaa aaaaaaa 337

<210> 45  
 <211> 1700  
 <212> DNA  
 <213> Homo sapiens

<400> 45  
 tgtttgcatt aagttcatag attataatth gtaatggaat caacacccaa tgcaaattag 60  
 aaagagagcc cactttgctc acccagtcac gtcttcccat gtaaccatag aacgttgggg 120  
 tctgtgtctt ttctagatcc acagtcttgc tctcagaaca ggctagccac accacaggcc 180  
 tagtgccagg acccatggcc tttttttaag ctcagactcc cttctgtgaa cagcaatatc 240  
 cccacaactt gtacaacatt ggtgcttcct gcaagggcta cagaactatt tgatacgaaa 300  
 atgttcattg acttacacac aagagaagca caaaataaaa aattaataat taatttaattg 360  
 tctttgaaaa tgtaccatth atttttacat ttgggggtcat aagaattgta ttacacttaa 420  
 gaatgcaata caatttgaag atcagattht tctccctttg tgagaatttc tcagtatgtg 480  
 tgatgactac caagaaatca tagccagtca taaattcagt gagttactca taaacgaaca 540  
 agaaccacct acttcttggg gaggtaggtc tgcttccctt caactcagga tacaactgct 600  
 ttcaactgct ttcttcacat tagctgacta attagctaga agcctgtcgt aaacaattth 660  
 atggttgact ctttccctgg gctcagggtt ccctagaaca gagagggtccc caaatcccgg 720  
 tctgtggcct gtccgcctaa gctctgcctc ctgccagatc agcaggcagc attagattct 780  
 cataggagct ggacgcctat tgtgaactgc gcatgtgcgg gatccagatt gtgcactctt 840  
 tatgagaatc taactaatgc ttgatgatct atctgaacca gaacaatttc atcctgaaac 900  
 catccccac caatccatag aaatactgtc ttccacaaaa atgatccctg gtgccaaaaa 960  
 tgtagagac cactccccta aaactctctt cttagctctc acctcctgta ttactatctc 1020

atctcagtac attgaagccc ccatcttttc cccatggatg cctcatttcc tattagggag 1080  
 gcattttttt attttttggt tttatttttt tccgagacgg agtctcgctc tgtcgccaag 1140  
 gctggagtgct agtggcgcca tctcggctca ctgcaagctc cgcctcccgg gttcacgcca 1200  
 ttctcctgcc tcagcctccc aagtagctgg gactacaggc gcccgacta cgcccggcta 1260  
 attttttgta tttttagtag agacgggggt tcaccgtggt agccaggatg gtctcgatct 1320  
 cctgacctcg tgatccgccc gccttggcct cccaaagtgc tgggattaca ggcgtgagac 1380  
 cgcgcccggc cgtcatttgg tatgtcttaa tgtgcctcag gacctagcac agtccctggt 1440  
 acccagtaga gacctatgta atgttcgtta ttcaataata aatacatgaa ttaaagagtg 1500  
 agagtggatt ttgtaatggt acgactgata gagaaatact cagtgattct aagggatggg 1560  
 gaagaacggg tggagctaga gggtgtgctc aggaaactat taaatagacg ttccgcagga 1620  
 agggattgac gaagtgtgag gttaatgagg aagggaat aaatataaa atttggtggt 1680  
 ggaaaagatc tgattcatga 1700

<210> 46  
 <211> 2419  
 <212> DNA  
 <213> Homo sapiens

<400> 46  
 taaccagcgg gccctggctc aagtgtggc tctgtgtcc ttgccttcca tttcccctct 60  
 gcaccagaa cagtgggtggc aacattcatt gccaagggcc caaagaaaga gctacctgga 120  
 ccttttgttt tctgtttgac aacatgttta ataaataaaa atgtcttgat atcagtaaga 180  
 atcagagtct tctcactgat tctgggcata ttgatcttcc cccatttttc tctacttggc 240  
 tgctccctga gaggactgca taggatagaa atgccttttt cttttctttt cgtttttttt 300  
 tttttttttt tttgagatgg agtctcactc tgtcgcccag gcttaagtgc aatggcacia 360  
 tctcggctca ctgcaacctc tctctcctgg gttcaagtga ttctcctgcc tcagcctccc 420  
 aaatagctga gattacaggc atgcaccacc acacctggct aatttttgtg tttttagtag 480  
 agacagggtt tcaccgtttt ggccagggtg gtcttgaact cctgacctcg ggagatccgc 540  
 ccaccttggc ctctctttgt gctgggatta caggcatgag ccaactgagcc gggccacttt 600  
 ttccttatca gtcagttttt acaagtcatt agggaggtag actttacctc tctgtgaagg 660  
 aaagtatggt atgttgatct acagagagag atggaaaaat tccagggtc gtagctacta 720  
 agcagaattt ccaagatagg caaattgttt tttctgtcaa ataataagct aatattactt 780  
 ctacaaatat gagaccttgg agagaagttt ccaaggacca agtaccaaca taccaacaga 840

```

ttattatagt ttctctcact cttacacaca cacacacaca tatacacata tgtaatccag      900
catgaataacc aaaattcatt cagggtagcc accttttgtc ttaatcgaga gataattttg      960
atgtttgaat ggaatgctcc caggatattc tcttgtcatg gttattttat ataaaattca     1020
aaaaccaatt acattatttc ctctgtaatc ttttacttta tcaactaatg tctggcaagt     1080
gtgatgtttt ggggaagtta tagaagattc cggccaggcg cttatctcac gcttgtaatc     1140
cagcactttg ggaagctgag gcggacagat cagcagggtca agagatcaag accatcctgg     1200
acaacatggg gaaaccttgt ctctactaaa aatgtgaaaa ttagctgggc gtggtggcac     1260
acacctatag tcccagctac tcgggagggt gaggcaggag aatcgcttga acctaggagg     1320

cggaggttgc actgagccga gatcacgcca ctgcactcca gcctgggcca cagagcgaga     1380
ctccatctca aaaaaaaaaa aaaaagaaag atcccagttt atcccagttt atcccttatt     1440
cttcctcaat tctcaagatt tgtttttaag ttaacataac ttaggttaac acactctttg     1500
taaaatacac tgttcaatct acagactcag tggttagctt cctgttaact aatttctggt     1560
gacagggtact tggatatttt atttagaaag tggttgcaa taaattagtt ataagtcgcc     1620
agtttctact ccttgtgaac acataattat tgtggtctca gtattcccta tgggtggcttc     1680
tcctgctcct ggtattgccc tgaaatgggc caaaagccgt ggctcccca tgcctcaggtt     1740
atagaacatt gtccagggtac cacctaggag agcccagcct cactgaaagt attcaaattt     1800
aggaatgggt ttgagaagta ggtagctggt atgtgcttag cacaagaatc tctcttcctt     1860
gggttagtct gtttcaaac tgaaaacact gtcattcctt aagaaaatag gaaaaagtat     1920
tccaaacctc tgtcactaga aaatttgcca tattaccaa tctcaaaaac ctctcaggaa     1980
atgagaaagt cccagtttct ggtaaactat ttgggccctt ttctcaagtt ctcttccag     2040
tgctatttcc ttgaggtgag gcaaagttac tcaagatcat cgctgccact caaggccttg     2100
atagggcaag tgaaaggcat ggaccattat tatattgatc acagcataag ctgtgaaaac     2160
ccacatcttc tccaaacatc tgcttgagc attatcatcg catagtttgc tctggtgttc     2220
agggaaatcg ctgtttcata ggaaatcaca tggcagtggt atgggagtgt ttcttgacct     2280
gccgatggta ctggcacctg agcaagcatt cctagtcctt tttggtctgg gcctcttggt     2340
ctatcacaac cacaagctgt ttaaaataaa aacgtcaagt cacaggcagg tcattttatc     2400
ctgctgaat caattgaag                                     2419

```

```

<210> 47
<211> 297
<212> DNA
<213> Homo sapiens

```

<400> 47  
 tcctcagtgc acagtgtctgc ctctgtctgag gggacaggag gatcaccctc ttcgtctgctt 60  
 cggccagtgt gtcgggctgg gccctgacaa gccacctgag gagaggctcg gagccggggc 120  
 cggaccccg cgattgccgc ccgcttctct ctagtctcac gaggggtttc ccgcctcgca 180  
 cccccacctc tggacttgcc tttccttctc ttctccggt gtggaggag ccagcgctta 240  
 ggccggagcg agcctggggg ccgccgcgcg tgaagacatc gcggggaccg attcacc 297

<210> 48  
 <211> 1192  
 <212> DNA  
 <213> Homo sapiens

<400> 48  
 tgagcttttt ctttaatttca ttcctttttt tggacactgg tggctcacta cctaaagcag 60  
 tctatttata ttttctacat ctaatttttag aagcctggct acaatactgc acaaacttgg 120  
 ttagttcaat ttttgatccc ctttctactt aatttacatt aatgctcttt tttagtatgt 180  
 tctttaatgc tggatcacag acagctcatt ttctcagttt tttggtattt aaaccattgc 240  
 attgcagtag catcatttta aaaaatgcac ctttttattt atttattttt ggctagggag 300  
 tttatccctt tttcgaatta tttttaagaa gatgccata taatttttgt aagaaggcag 360  
 taacctttca tcatgatcat aggcagttga aaaattttta cacctttttt ttcacatttt 420  
 acataaataa taatgctttg ccagcagtac gtggtagcca caattgcaca atatatatttc 480  
 ttaaaaaata ccagcagtta ctcatggaat atattctgcg tttataaaac tagtttttaa 540  
 gaagaaattt tttttggcct atgaaattgt taaacctgga acatgacatt gttaatcata 600  
 taataatgat tcttaaatgc tgtatggttt attattttaa tgggttaaagc catttacata 660  
 atatagaaaag atatgcatat atctagaagg tatgtggcat ttatttggat aaaattctca 720  
 attcagagaa atcatctgat gtttctatag tcactttgcc agctcaaaag aaaacaatac 780  
 cctatgtagt tgtggaagtt tatgctaata ttgtgtaact gatattaaac ctaaattgttc 840  
 tgcctaccct gttggtataa agatatatttg agcagactgt aaacaagaaa aaaaaaatca 900  
 tgcattctta gcaaaattgc ctagtatgtt aatttgctca aaatacaatg tttgatttta 960  
 tgcactttgt cgctattaac atcctttttt tcatgtagat ttcaataatt gagtaatttt 1020  
 agaagcatta ttttaggaat atatagttgt cacagtaa atcttggttt ttctatgtac 1080  
 attgtacaaa tttttcattc cttttgctct ttgtggttgg atctaact aactgtattg 1140  
 ttttgttaca tcaaataaac atcttctgtg gaccaggaaa aaaaaaaaaa aa 1192

<210> 49  
 <211> 197

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 49

```

agacagcctt aacccacggg cgcgggagcg tcgtatgggc aggggcaggc gggagcgacg      60
tggggagcgc ctcacgaacg atcagagctg cgggagcgcg aacgaagccc ggaggccgca      120
ggctgagcgc tccctcgcag cagccgggagc ggcaaaagcc cccagtcctc ggccccgcgc      180
caagcgcgcg cgggaaa                                     197

```

&lt;210&gt; 50

&lt;211&gt; 3293

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 50

```

taattattta tattgtaaag aattttaaca gtcctgggga cttccttgaa ggatcatttt      60
cacttttgct cagaagaaag ctctggatct atcaaataaa gaagtccttc gtgtgggcta      120
catatataga tgttttcatg aagaggagtg aaaagccaga aggatataga caaatgaggc      180
ctaagacctt tcctgccagt aactatactg tcagtagccg gcaaagtgtta caagaaattc      240
gggaatccct taggaattta tctaaaccat ctgatgctgc taaggctgag cataacatga      300
gtaaaatgtc aaccgaagat cctcgacaag tcagaaatcc acccaaattt gggacgcac      360
ataaagcctt gcaggaaatt cgaaactctc tgcttccatt tgcaaataaa acaaattcct      420
ctcggagtac ttcagaagtt aatccacaaa tgcttcaaga cttgcaagct gctggatttg      480
atgaggatat gggtatacaa gctcttcaga aaactaaca cagaagtata gaagcagcaa      540
ttgaattcat tagtaaaatg agttaccaag atcctcgacg agagcagatg gctgcagcag      600
ctgccagacc tattaatgcc agcatgaaac cagggaatgt gcagcaatca gttaaccgca      660
aacagagctg gaaagggtct aaagaatcct tagttcctca gaggcattgg ccgccactag      720
gagaaagtgt ggctatcat tctgagagtc ccaactcaca gacagatgta ggaagacctt      780
tgtctggatc tggtatatca gcatttgctc aagctcacc tagcaacgga cagagagtga      840
acccccacc accacctcaa gtaaggagtg ttactcctcc accacctcca agaggccaga      900
ctccccctcc aagagggtaca actccacctc ccccttcctg ggaaccaaac tctcaaaca      960
agcgctattc tggaaacatg gaatacgtaa tctcccgaat ctctcctgtc ccacctgggg      1020
catggcaaga gggctatcct ccaccacctc tcaacacttc ccccatgaat cctcctaata      1080
aaggacagag aggcattagt tctgttcctg ttggcagaca accaatcatc atgcagagtt      1140
ctagcaaatt taactttcca tcaggagac ctggaatgca gaatggtact ggacaaactg      1200
atttcatgat acacaaaaat gttgtccctg ctggcactgt gaatcggcag ccaccacctc      1260

```

catatcctct	gacagcagct	aatggacaaa	gcccttctgc	ttacaaaaca	gggggatctg	1320
ctgctccttc	gtcatataca	aatggaagta	ttcctcagtc	tatgatgggtg	ccaaacagaa	1380
atagtcataa	catggaacta	tataacatta	gtgtacctgg	actgcaaaca	aattggcctc	1440
agtcctcttc	tgctccagcc	cagtcatccc	cgagcagtg	gcatgaaatc	cctacatggc	1500
aacctaacat	accagtgagg	tcaaattctt	ttaataaccc	attaggaaat	agagcaagtc	1560
actctgctaa	ttctcagcct	tctgctacaa	cagtcactgc	aattacacca	gctcctattc	1620
aacagcctgt	gaaaagtatg	cgtgtattaa	aaccagagct	acagactgct	ttagcaccta	1680
cacacccttc	ttggatacca	cagccaattc	aaactgttca	accagtcct	tttcctgagg	1740
gaaccgcttc	aaatgtgact	gtgatgccac	ctgttgctga	agctccaaac	tatcaaggac	1800
caccaccacc	ctacccaaaa	catctgctgc	acccaaaacc	atctgttcct	ccatacagagt	1860
caatcagtaa	gcctagcaaa	gaggatcagc	caagcttgcc	caaggaagat	gagagtgaaa	1920
agagttatga	aaatgttgat	agtggggata	aagaaaagaa	acagattaca	acttcacctc	1980
ttactgttag	gaaaaacaag	aaagatgaag	agcgaaggga	atctcgtatt	caaagttatt	2040
ctcctcaagc	atttaaattc	tttatggagc	aacatgtaga	aaatgtactc	aaatctcatc	2100
agcagcgtct	acatcgtaaa	aaacaattag	agaatgaaat	gatgcggggt	ggattatctc	2160
aagatgcca	ggatcaaag	agaaagatgc	tttgccaaaa	agaatcta	tacatccgtc	2220
ttaaaagggc	taaaatggac	aagtctatgt	ttgtgaagat	aaagacacta	ggaataggag	2280
catttggtga	agtctgtcta	gcaagaaaag	tagatactaa	ggctttgtat	gcaacaaaaa	2340
ctcttcgaaa	gaaagatgtt	cttcttcgaa	atcaagtcgc	tcattgttaag	gctgagagag	2400
atatcctggc	tgaagctgac	aatgaatggg	tagttcgtct	atattattca	ttccaagata	2460
aggacaattt	atactttgta	atggactaca	ttcctggggg	tgatatgatg	agcctattaa	2520
ttagaatggg	catctttcca	gaaagtctgg	cacgattcta	catagcagaa	cttacctgtg	2580
cagttgaaag	tgttcataaa	atgggtttta	ttcatagaga	tattaaacct	gataatattt	2640
tgattgatcg	tgatgggtcat	attaaattga	ctgactttgg	cctctgcact	ggcttcagat	2700
ggacacacga	ttctaagtac	tatcagagtg	gtgaccatcc	acggcaagat	agcatggatt	2760
tcagtaatga	atggggggat	ccctcaagct	gtcgatgtgg	agacagactg	aagccattag	2820
agcggagagc	tgacgcgcag	caccagcgat	gtctagcaca	ttctttgggt	gggactccca	2880
attatattgc	acctgaagtg	ttgctacgaa	caggatacac	acagttgtgt	gattgggtgga	2940
gtgttggtgt	tattcttttt	gaaatgttgg	tgggacaacc	tcctttcttg	gcacaaacac	3000
cattagaaac	acaaatgaag	gtcacctgct	gctatataca	tcattggctc	gagaagaaac	3060



tactgaacac cctgcgagag agaagcctag aaaagaaaga aagggccaaa aggttttgaa 3120  
 ctcttcatcc ctaatttgct acaactgatca aaaccaagta agggctcctg aagtccatga 3180  
 gtctatcatc aatcagcaca aatgctatac tagtttgtaa ctgcgggggtc agttgtgaag 3240  
 gggaaggaca gcagtcttat ccatattcca ggaagccaca gtaaactgct cga 3293

<210> 51  
 <211> 424  
 <212> DNA  
 <213> Homo sapiens

<400> 51  
 cctactctat tcagatatcc tccagattcc taaagattag agatcatttc tcattctcct 60  
 aggagtactc acttcaggaa gcaaccagat aaaagagagg tgcaacggaa gccagaacat 120  
 tcctcctgga aattcaacct gtttcgcagt ttctcgagga atcagcattc agtcaatccg 180  
 ggccggggagc agtcatctgt ggtgaggctg attggctggg caggaacagc gccggggcgt 240  
 gggctgagca cagcgcttcg ctctctttgc cacaggaagc ctgagctcat tcgagtagcg 300  
 gctcttccaa gctcaaagaa gcagaggccg ctgttcgttt cctttagggtc tttccactaa 360  
 agtcggagta tcttcttcca agatttcacg tcttggtggc cgttccaagg agcgcgaggc 420  
 cggg 424

<210> 52  
 <211> 706  
 <212> DNA  
 <213> Homo sapiens

<400> 52  
 tgaactctga ctgtatgaga tgttaaatac tttttaatat ttgttttagat atgacattta 60  
 ttcaaagtta aaagcaaaca cttacagaat tatgaagagg tatctgttta acatttcctc 120  
 agtcaagttc agagtcttca gagacttcgt aattaaagga acagagtgag agacatcatc 180  
 aagtggagag aaatcatagt ttaaactgca ttataaattt tataacagaa ttaaagtaga 240  
 ttttaaaaga taaaatgtgt aattttgttt atattttccc atttggactg taactgactg 300  
 ccttgctaaa agattataga agtagcaaaa agtattgaaa tgtttgcata aagtgtctat 360  
 aataaaaacta aactttcatg tgactggagt catcttgtcc aaactgcctg tgaatatatc 420  
 ttctctcaat tggaatattg tagataactt ctgctttaa aaagttttct ttaaataatc 480  
 ctactcattt ttgtgggaat ggttaagcag tttaaataat tcctgtgtat atgtctatca 540  
 cataggggtc taacagaaca atctggattc attatttcta ggacttgatc ctgctgatgc 600  
 tgaatttgca cattaagggtg tgttaacaac caaaacacag atcgatataa gaagtaagga 660

ggtggggaga ggcaaattat gatgtgctat gagttagatg tatagt

706

<210> 53  
 <211> 239  
 <212> DNA  
 <213> Homo sapiens

<400> 53  
 agtccgcggc gttccccggc tgcagccggg agggggccga ggagtgactg agccccgggc 60  
 tgtgcagtcc gacgccgact gaggcacgag cgggtgacgc tgggcctgca gcgcggagca 120  
 gaaagcagaa cccgcagagt cctccctgct gctgtgtgga cgacacgtgg gcacaggcag 180  
 aagtgggccc tgtgaccagc tgcactgggt tcgtggaagg aagctccagg actggcggg 239

<210> 54  
 <211> 641  
 <212> DNA  
 <213> Homo sapiens

<400> 54  
 tgaggcagct gctatcccca tctccctgcc tggcccccaa cctcagggct cccaggggtc 60  
 tccctggctc cctcctccag gcctgcctcc cacttcactg cgaagaccct cttgcccacc 120  
 ctgactgaaa gtagggggct ttctggggcc tagcgatctc tcctggccta tccgctgcca 180  
 gccttgagcc ctggctgttc tgtgggttcct ctgctcaccg cccatcaggg ttctcttacc 240  
 aactcagaga aaaatgctcc ccacagcgtc cctggcgcag gtgggctgga cttctacctg 300  
 ccctcaaggg tgtgtatatt gtataggggc aactgtatga aaaattgggg aggagggggc 360  
 cgggcgcggt gctcacgcct gtaatcccag cactttggga ggccgaggcg ggtggatcac 420  
 gaggtcagga gatcgagacc atcctggcta acatggtgaa acccgtctc tactaaaaat 480  
 aaaaaaaaaa ttagccggg cgcggtggcg ggcacctgta gtcccagcta cttgggaggc 540  
 tgaggcagga gaatggtgtg aaccggggag cggagggttc agtgagctga gatcgtgcta 600  
 ctgcactcca gcctggggga cagaaagaga ctccgtctca a 641

<210> 55  
 <211> 493  
 <212> DNA  
 <213> Homo sapiens

<400> 55  
 tttctgtgaa gcagaagtct gggaatcgat ctggaaatcc tcctaatttt tactccctct 60  
 cccccgact cctgattcat tgggaagttt caaatcagct ataactggag agagctgaag 120  
 attgatggga tcgttgccct atgcctttgt tttggtttta caaaaaggaa acttgacaga 180  
 ggatcatgct atacttaaaa aataacaacat cgcagaggaa gtagactcat attaaaaata 240

cttactaata ataacgtgcc tcatgaagta aagatccgaa aggaattgga ataaaacttt 300  
 cctgcatctc aagccaaggg ggaaacacca gaatcaagtg ttccgcgtga ttgaagacac 360  
 cccctcgtcc aagaatgcaa agcacatcca ataaaagagc tggattataa ctccctcttct 420  
 ttctctgggg gccgtggggg gggagctggg gcgagaggtg ccgttggccc ccgttgcttt 480  
 tcctctggga ggg 493

<210> 56  
 <211> 5282  
 <212> DNA  
 <213> Homo sapiens

<400> 56  
 tgaagtcaac atgcctgccc caaacaata tgcaaaaggt tactaaagc agtagaaata 60  
 atatgcattg tcagtgatgt tccatgaaac aaagctgcag gctgtttaag aaaaaataac 120  
 acacatataa acatcacaca cacagacaga cacacacaca cacaacaatt aacagtcttc 180  
 aggcaaaacg tcgaatcagc tatttactgc caaagggaaa tatcatttat tttttacatt 240  
 attaagaaaa aaagatttat ttatttaaga cagtcccatc aaaactcctg tctttggaaa 300  
 tccgaccact aattgccaag caccgcttcg tgtggctcca cctggatggt ctgtgcctgt 360  
 aaacatagat tcgctttcca tgttgttggc cggatcacca tctgaagagc agacggatgg 420  
 aaaaaggacc tgatcattgg ggaagctggc tttctggctg ctggaggctg gggagaaggt 480  
 gttcattcac ttgcatttct ttgccctggg ggctgtgata ttaacagagg gagggttcct 540  
 gtggggggaa gtccatgcct ccctggcctg aagaagagac tctttgcata tgactcacat 600  
 gatgcatacc tgggtgggagg aaaagagttg ggaacttcag atggacctag taccactga 660  
 gatctccacg ccgaaggaca gcgatgggaa aaatgccctt aaatcatagg aaagtatttt 720  
 ttaagctac caattgtgcc gagaaaagca ttttagcaat ttatacaata tcatccagta 780  
 ccttaagccc tgattgtgta tattcatata ttttggatac gcacccccca actcccaata 840  
 ctggctctgt ctgagtaaga aacagaatcc tctggaactt gaggaagtga acatttcggt 900  
 gacttccgca tcaggaaggc tagagttacc cagagcatca ggccgccaca agtgccctgct 960  
 tttaggagac cgaagtccgc agaacctgcc tgtgtcccag cttggaggcc tggtcctgga 1020  
 actgagccgg ggccctcact ggcctcctcc agggatgata aacagggcag tgtggtctcc 1080  
 gaatgtctgg aagctgatgg agctcagaat tccactgtca agaaagagca gtagaggggt 1140  
 gtggctgggc ctgtcacccct ggggccctcc aggtaggccc gttttcacgt ggagcatggg 1200  
 agccacgacc cttcttaaga catgtatcac tgtagaggga aggaacagag gccctgggcc 1260  
 cttcctatca gaaggacatg gtgaaggctg ggaacgtgag gagaggcaat ggccacggcc 1320

catttttggct gtagcacatg gcacgttggc tgtgtggcct tggcccacct gtgagtttaa 1380  
agcaaggctt taaatgactt tggagaggggt cacaaatcct aaaagaagca ttgaagtgag 1440  
gtgtcatgga ttaattgacc cctgtctatg gaattacatg taaaacatta tcttgtcact 1500  
gtagtttgggt tttatttgaa aacctgacaa aaaaaaagtt ccaggtgtgg aatatggggg 1560  
ttatctgtac atcctggggc attaaaaaaa aaatcaatgg tggggaacta taaagaagta 1620  
acaaaagaag tgacatcttc agcaaataaa ctaggaaatt ttttttctt ccagtttaga 1680  
atcagccttg aaacattgat ggaataactc tgtggcatta ttgcattata taccatttat 1740  
ctgtattaac tttggaatgt actctgttca atgtttaatg ctgtggttga tatttcgaaa 1800  
gctgctttta aaaaatacat gcatctcagc gtttttttgt ttttaattgt atttagttat 1860  
ggcctataca ctatttgtga gcaaagggtga tcgttttctg tttgagattt ttatctcttg 1920  
attcttcaaa agcattctga gaagggtgaga taagccctga gtctcagcta cctaagaaaa 1980  
acctggatgt cactggccac tgaggagctt tgtttcaacc aagtcatgtg catttccacg 2040  
tcaacagaat tgtttattgt gacagttata tctgttgtcc ctttgacctt gtttcttgaa 2100  
ggtttcctcg tccctgggca attccgcatt taattcatgg tattcaggat tacatgcatg 2160  
tttggttaaa cccatgagat tcattcagtt aaaaatccag atggcaaagt accagcagat 2220  
tcaaatctat ggtgggttga ccttttagaga gttgctttac gtggcctgtt tcaacacaga 2280  
cccaccaga gccctcctgc cctccttccg cgggggcttt ctcatggctg tccttcaggg 2340  
tcttcctgaa atgcagtgggt gcttacgctc caccaagaaa gcaggaaacc tgtggtatga 2400  
agccagacct ccccgcgagg cctcagggaa cagaatgatc agacctttga atgattctaa 2460  
tttttaagca aaatattatt ttatgaaagg tttacattgt caaagtgatg aatatggaat 2520  
atccaatcct gtgctgctat cctgccaaaa tcattttaat ggagtcagtt tgcagtatgc 2580  
tccacgtgggt aagatcctcc aagctgcttt agaagtaaca atgaagaacg tggacgcttt 2640  
taatataaag cctgttttgt cttctgttgt tgttcaaacg ggattcacag agtatttgaa 2700  
aaatgtatat atattaagag gtcacggggg ctaattgctg gctggctgcc ttttgctgtg 2760  
gggttttgtt acctggtttt aataacagta aatgtgccca gcctcttggc ccagaaactg 2820  
tacagtattg tggctgcact tgctctaaga gtagttgatg ttgcattttc cttattgtta 2880  
aaaacatggt agaagcaatg aatgtatata aaagcctcaa ctagtcattt ttttctctc 2940  
ttcttttttt tcattatata taattatttt gcagttgggc aacagagAAC catccctatt 3000  
ttgtattgaa gagggattca catctgcac ctaactgctc tttatgaatg aaaaaacagt 3060  
cctctgtatg tactcctctt tacactggcc agggtcagag ttaaataagag tatatgcact 3120

ttccaaattg gggacaaggg ctctaaaaaa agccccaaaa ggagaagaac atctgagaac	3180
ctcctcggcc ctcccagtc ctcgctgcac aaatactccg caagagagggc cagaatgaca	3240
gctgacaggg tctatggcca tcgggtcgtc tccgaagatt tggcaggggc agaaaactct	3300
ggcaggctta agatttgaa taaagtcaca gaatcaagga agcacctcaa tttagttcaa	3360
acaagacgcc aacattctct ccacagctca cttacctctc tgtgttcaga tgtggccttc	3420
catttatatg tgatctttgt tttattagta aatgcttata atctaaagat gtagctctgg	3480
cccagtggga aaaattagga agtgattata aatcgagagg agttataata atcaagatta	3540
aatgtaaata atcagggcaa tcccaacaca tgtctagctt tcacctccag gatctattga	3600
gtgaacagaa ttgcaaatag tctctatctg taattgaact tatcctaaaa caaatagttt	3660
ataaatgtga acttaaaact taattaattc caactgtact ttttaaggcag tggctgtttt	3720
tagactttct tatcacttat agttagtaat gtacacctac tctatcagag aaaaacagga	3780
aaggctcgaa atacaagcca ttctaaggaa attagggagt cagttgaaat tctattctga	3840
tcttattctg tgggtgtctt tgcagcccag acaaatgtgg ttacacactt ttttaagaaat	3900
acaattctac attgtcaagc ttatgaagg tccaatcaga tctttattgt tattcaattt	3960
ggatctttca gggatttttt ttttaaatta ttatgggaca aaggacattt gttggagggg	4020
tgggagggag gaacaatttt taaatataaa acattcccaa gtttggatca gggagtggga	4080
agttttcaga ataaccagaa ctaagggtat gaaggacctg tattgggggc gatgtgatgc	4140
ctctgcgaag aaccttgtgt gacaaatgag aaacattttg aagtttgtgg tacgacctt	4200
agattccaga gacatcagca tggctcaaag tgcagctccg tttggcagtg caatggtata	4260
aatttcaagc tggatatgtc taatgggtat ttaacaata aatgtgcagt ttttaactaac	4320
aggatattta atgacaacct tctgggttgg aggacatct gtttctaaat gtttattatg	4380
tacaatacag aaaaaaattt tataaaatta agcaatgtga aactgaattg gagagtgata	4440
atacaagtcc tttagtctta cccagtgaat cattctgttc catgtctttg gacaaccatg	4500
accttggaca atcatgaaat atgcatctca ctggatgcaa agaaaatcag atggagcatg	4560
aatggtactg taccggttca tctggactgc ccagaaaaa taacttcaag caaacatcct	4620
atcaacaaca aggttgttct gcataccaag ctgagcacag aagatgggaa cactggtgga	4680
ggatggaaag gctcgtcaa tcaagaaaat tctgagacta ttaataaata agactgtagt	4740
gtagatactg agtaaatacca tgcacctaaa ccttttggaa aatctgccgt gggccctcca	4800
gatagctcat ttcattaagt ttttccctcc aaggtagaat ttgcaagagt gacagtggat	4860
tgcatttctt ttggggaagc tttcttttgg tggttttgtt tattatacct tcttaagttt	4920
tcaaccaagg tttgcttttg ttttgagtta ctgggggttat ttttgtttta aataaaaaata	4980

agtgtacaat aagtgttttt gtattgaaag cttttgttat caagattttc atactttttac 5040  
 cttccatggc tctttttaag attgatactt ttaagagggtg gctgatattc tgcaacactg 5100  
 tacacataaa aaatacggta aggatacttt acatgggttaa ggtaaagtaa gtctccagtt 5160  
 ggccaccatt agctataatg gcactttgtt tgtgttgttg gaaaaagtca cattgccatt 5220  
 aaactttcct tgtctgtcta gttaatattg tgaagaaaaa taaagtacag tgtgagatac 5280  
 tg 5282

<210> 57  
 <211> 117  
 <212> DNA  
 <213> Homo sapiens

<400> 57  
 attcggggcg agggaggagg aagaagcgga ggaggcggct cccgctcgca gggccgtgca 60  
 cctgcccgcg cgcccgcctg ctgctcgcc cgccgcgcg cgctgccgac cgccagc 117

<210> 58  
 <211> 430  
 <212> DNA  
 <213> Homo sapiens

<400> 58  
 tgatccaggg agccccacc atccgggggg accccgagtg tcatctcttc tacaatgagc 60  
 agcaggaggc ttgcgggggtg cacaccagc ggatgcagta gaccgcagcc agccggtgcc 120  
 tggcgcccct gcccccgcc cctctccaaa caccggcaga aaacggagag tgcttgggtg 180  
 gtgggtgctg gaggattttc cagtcttgac acacgtatth atatttgga agagaccagc 240  
 accgagctcg gcacctcccc ggctctctc tcccagctg cagatgccac acctgctcct 300  
 tcttgctttc cccgggggag gaagggggtt gtggtcgggg agctggggta caggtttggg 360  
 gagggggaag agaaatthtt atthttgaac ccctgtgtcc cthttgcata agattaaagg 420  
 aaggaaaagt 430

<210> 59  
 <211> 192  
 <212> DNA  
 <213> Homo sapiens

<400> 59  
 tcctaggcgg cggccgcggc ggcggaggca gcagcggcgg cggcagtggc ggcggcgaag 60  
 gtggcggcgg ctgggccagt actccggcc cccgccatth cggactggga gcgagcgcg 120  
 cgcaggcact gaaggcggcg gcggggccag aggctcagcg gctcccaggt gcgggagaga 180  
 ggctgctga aa 192

<210> 60  
 <211> 4172  
 <212> DNA  
 <213> Homo sapiens

<400> 60  
 taaataacaat ttgtactttt ttcttaaggc atactagtac aagtggtaat ttttgtacat 60  
 tacactaaat tattagcatt tgtttttagca ttacctaatt tttttcctgc tccatgcaga 120  
 ctggttagctt ttaccttaaa tgcttatattt aaaatgacag tggaagtttt tttttcctcg 180  
 aagtgccagt attcccagag ttttggtttt tgaactagca atgcctgtga aaaagaaaact 240  
 gaatacctaa gatttctgtc ttgggggtttt tgggtgcatgc agttgattac ttcttatattt 300  
 tcttaccaag tgtgaatggt ggtgtgaaac aaattaatga agcttttgaa tcatccctat 360  
 tctgtgtttt atctagtcac ataaatggat taattactaa tttcagttga gaccttctaa 420  
 ttggtttttta ctgaaacatt gagggacaca aatttatggg cttcctgatg atgattcttc 480  
 taggcacatcat gtcctatagt ttgtcatccc tgatgaatgt aaagttacac tgttcacaaa 540  
 gggttttgtct cttttccact gctattagtc atgggtcactc tccccaaaat attatatattt 600  
 ttctataaaa agaaaaaaat ggaaaaaaat tacaaggcaa tggaaaactat tataaggcca 660  
 tttccttttc acattagata aattactata aagactccta atagcttttt cctgttaagg 720  
 cagacccagt atgaatggga ttattatagc aaccattttg gggctatatt tacatgctac 780  
 taaattttta taataattga aaagatttta acaagtataa aaaaattctc ataggaatta 840  
 aatgtagtct ccctgtgtca gactgctctt tcatagtata actttaaatc ttttcttcaa 900  
 cttgagtctt tgaagatagt ttttaattctg cttgtgacat taaaagatta tttgggccag 960  
 ttatagctta ttaggtgttg aagagaccaa ggttgcaagc caggccctgt gtgaaccttg 1020  
 agctttcata gagagtttca cagcatggac tgtgtgcccc acggtcaccc gagtgggtgt 1080  
 acgatgcatt ggtagtcaa aaatggggag ggactagggc agtttgata gctcaacaag 1140  
 atacaatctc actctgtggt ggtcctgctg acaaatacaag agcattgctt ttgtttctta 1200  
 agaaaacaaa ctctttttta aaaattactt ttaaataatta actcaaaagt tgagattttg 1260  
 ggggtggtggt gtgccaagac attaatattt tttttaaaca atgaagtga aaagttttac 1320  
 aatctctagg tttggctagt tctcttaaca ctgggttaaat taacattgca taaacacttt 1380  
 tcaagtctga tccatattta ataatgcttt aaaataaaaa taaaaacaat ctttttgata 1440  
 aatttaaaat gttacttatt ttaaaataaa tgaagtgaga tggcatggtg aggtgaaagt 1500  
 atcactggac taggttggtg gtgacttagg ttctagatag gtgtctttta ggactctgat 1560

tttgaggaca	tcacttacta	tccattttctt	catgttaaaa	gaagtcattct	caaactctta	1620
gtttttttttt	tttactactat	gtgattttata	ttccattttac	ataaggatac	actttatttgt	1680
caagctcagc	acaatctgta	aattttttaac	ctatgtttaca	ccatcttcag	tgccagttctt	1740
gggcaaaatt	gtgcaagagg	tgaagttttat	atgttgaaat	ccattctctgt	tttaggactc	1800
ttctttccata	ttagtgtcat	cttgccctccc	tacctttccac	atgccccatg	acttgatgca	1860
gttttaatac	ttgtaattcc	cctaaccata	agattttactg	ctgctgtgga	tatctccatg	1920
aagtttttccc	actgagtcac	atcagaaatg	ccctacattct	tatttttctc	aggggtcaag	1980
agaatctgac	agataccata	aagggtatttg	acctaattcac	taatttttcag	gtgggtggctg	2040
atgcttttgaa	catctctttg	ctgccccatc	cattagcgcac	agtaggattt	ttcaaccctg	2100
gtatgaatag	acagaaccct	atccagtggg	aggagaattt	aataaagata	gtgcagaaaag	2160
aattccttag	gtaattctata	actaggacta	ctcctggtaa	cagtaataca	ttccattgtt	2220
ttagtaacca	gaaatcttca	tgcaatgaaa	aatacttttaa	ttcatgaagc	ttactttttt	2280
tttttttggtg	tcagagtctc	gctcttgtca	cccaggctgg	aatgcagtgg	cgccattctca	2340
gctcactgca	acctttccatc	ttcccagggt	caagcgattc	tcgtgcctcg	gcctcctgag	2400
tagctgggat	tacaggcgtg	tgcaactacac	tcaactaatt	tttgtatttt	taggagagac	2460
ggggttttcac	ctgttggcca	ggctgggtctc	gaactcctga	cctcaagtga	ttcaccacc	2520
ttggcctcat	aaacctgttt	tgcaagaactc	atttattcag	caaataattta	ttgagtgcct	2580
accagatgcc	agtcaccgca	caaggcactg	ggtatatggt	atccccaac	aagagacata	2640
atcccgggtcc	ttagggtactg	ctagtgtggt	ctgtaatatc	ttactaaggc	ctttggtata	2700
cgaccagag	ataacacgat	gcgtattttta	gttttgcaaa	gaaggggttt	ggtctctgtg	2760
ccagctctat	aattgttttg	ctacgattcc	actgaaactc	ttcgatcaag	ctactttatg	2820
taaatcactt	cattgtttta	aaggaataaa	cttgattata	ttgttttttt	atttggcata	2880
actgtgattc	ttttaggaca	attactgtac	acattaaggt	gtatgtcaga	tattcatatt	2940
gacccaaatg	tgtaatatcc	cagttttctc	tgcataagta	attaaaatat	acttaaaaat	3000
taatagtttt	atctgggtac	aaataaacag	tgctgaact	agttcacaga	caagggaaac	3060
ttctatgtaa	aaatcactat	gattttctgaa	ttgctatgtg	aaactacaga	tctttggaac	3120
actgtttagg	taggggtgta	agacttgaca	cagtacctcg	ttctacaca	gagaaagaaa	3180
tgccatact	tcaggaactg	cagtgtttat	gaggggatat	ttaggcctct	tgaatttttg	3240
atgtagatgg	gcattttttt	aaggtagtgg	tttaattacct	ttatgtgaac	tttgaatggt	3300
ttaacaaaag	atttgttttt	gtagagattt	taaaggggga	gaattctaga	aataaatggt	3360
acctaattat	tacagcctta	aagacaaaaa	tccttgttga	agttttttta	aaaaaagact	3420



aaattacata gacttaggca ttaacatggt tgtggaagaa tatagcagac gtatatgtga 3480  
 tcatttgagt gaatgttccc aagtaggcat tctaggtctt atttaactga gtcacactgc 3540  
 ataggaattt agaacctaac ttttataggt tatcaaaact gttgtcacca ttgcacaatt 3600  
 ttgtcctaata atatacatag aaactttgtg gggcatgtta agttacagtt tgcacaagtt 3660  
 catctcattt gtattccatt gatttttttt tttcttctaa acattttttt ttcaaaacag 3720  
 tataataaac ttttttttagg ggattttttt tagacagcaa aaaactatct gaagatttcc 3780  
 atttgtcaaa aagtaatgat ttcttgataa ttgtgtagtg aatgtttttt agaaccagc 3840  
 agttaccttg aaagctgaat ttatatattag taacttctgt gttaatactg gatagcatga 3900  
 attctgcatt gagaaactga atagctgtca taaaatgctt tctttcctaa agaaagatac 3960  
 tcacatgagt tcttgaagaa tagtcataac tagattaaga tctgtgtttt agtttaatag 4020  
  
 tttgaagtgc ctgtttggga taatgatagg taatttagat gaatttaggg gaaaaaaaaag 4080  
 ttatctgcag ttatgttgag ggcccatctc tccccccaca cccccacaga gctaactggg 4140  
 ttacagtgtt ttatccgaaa gtttccaatt cc 4172

<210> 61  
 <211> 238  
 <212> DNA  
 <213> Homo sapiens

<400> 61  
 ccattgtgct ggaaaggcgc gcaacggcgg cgacggcggc gacccaccg cgcattcctgc 60  
 caggcctccg cgcccagccg cccacgcgcc cccgcgcccc gcgccccgac cctttcttcg 120  
 cgcccccgcc cctcgccccg ccaggccccc ttgccggcca cccgccaggc cccgcgccgg 180  
 cccgcccgcc gcccaggacc ggcccgcgcc ccgcaggccg cccgccgcc gcgcccgc 238

<210> 62  
 <211> 547  
 <212> DNA  
 <213> Homo sapiens

<400> 62  
 ggccccgcag ctctggccac agggacctct gcagtgcccc ctaagtgacc cggacacttc 60  
 cgaggggggc atcacgcct gtgtatataa cgtttccggg attactctgc tacacgtagc 120  
 ctttttactt ttgggggttt gtttttgttc tgaactttcc tgttaccttt tcagggtga 180  
 tgtcacatgt aggtggcgtg tatgagtga gacgggcctg ggtcttgggg actggagggc 240  
 aggggtcctt ctgcccctgg ggtcccaggg tgctctgcct gctcagccag gcctctcctg 300  
 ggagccactc gccagagac tcagcttggc caacttgggg ggctgtgtcc acccagccc 360

cccgtcctgt gggctgcaca gtcaccttg ttccctcctg ccccggttcg agagccgagt 420  
ctgtggggcac tctctgcctt catgcacctg tcttttctaa cacgtcgcct tcaactgtaa 480  
tcacaacatc ctgactccgt catttaataa agaaggaaca tcaggcatgc taaaaaaaaa 540  
aaaaaaa 547

<210> 63  
<211> 102  
<212> DNA  
<213> Homo sapiens

<400> 63  
gaattccggc aaacatgagg cagctgccag ccggcctggg cagtcttgtc tgcctcggct 60  
gtgaagtggg gaggtggca acagttttct tcagcgccca gg 102

<210> 64  
<211> 2017  
<212> DNA  
<213> Homo sapiens

<400> 64  
gacacgtcca aaggagtgca tggccacagc cacctccacc cccaagaaac ctccatcctg 60  
ccaggagcag cctccaagaa acttttaaaa aatagatttg caaaaagtga acagattgct 120  
acacacacac acacacacac acacacacac acacacagcc attcatctgg gctggcagag 180  
gggacagagt tcagggaggg gctgagctctg gctagggggc gagtccagag gccccagcca 240  
gcccttccca ggccagcgag gcgaggctgc ctctgggtga gtggctgaca gagcaggctc 300  
gcaggccacc agctgctgga tgtcaccaag aaggggctcg agtgccctgc aggagggtcc 360  
aatcctccgg tccacctcg tcccggtcat ccattctgct ttcttgccac acagtggccg 420  
gccaggctc ccctgggtct ctccccgtag ccactctctg cccactacct atgcttctag 480  
aaagcccctc acctcaggac ccagaggac cagctggggg gcagggggga gagggggtaa 540  
tggaggccaa gcctgcagct ttctggaaat tcttccctgg gggctccagt atcccctgct 600  
actccactga cctggaagag ctgggtacca ggccaccac tgtggggcaa gcctgagtgg 660  
tgagggggcca ctggcatcat tctccctcca tggcaggaag gcgggggatt tcaagtttag 720  
ggattgggtc gtgggtggaga atctgagggc actctgccag ctccacaggt ggatgagcct 780  
ctccttgccc cagtccctgg tcaagtggaa tgcagtgggt ggggctgtac acaccctcca 840  
gcacagactg ttccctccaa ggtcctctta ggtcccgggg aggaacgtgg ttcagagact 900  
ggcagccagg gagccccggg cagagctcag aggagtctgg gaagggggcgt gtccctcctc 960  
ttcctgtagt gccctccca tggcccagca gcttggctga gccctctcc tgaagcagct 1020

```

gtgcgcgcgtc cctctgcctt gcacaaaaag cacaagacat tccttagcag ctacagcgcag 1080
ccctagtggg agcccagcac actgctttct ggaggccagg ccctcctgct ggctgagctt 1140
gggcccgggtg gcccgaatat ggtggccctg gggaagaggc cttgggggtc tgctctgtgc 1200
ctgggatcag tggggcccca aagcccagcc cggtgacca acattcaaaa gcacaaaccc 1260
tggggactct gcttggtgtt cccctccatc tggggatgga gaatgcagcc caaagctgga 1320
gccaatggtg agggctgaga gggctgtggc tgggtggtca gcagaaaccc caggaggaga 1380
gagatgtgtc tccgcctga ttggggcctc acccagaagg aaccgggtcc cagccgcatg 1440
gcccctccag gaacattccc acataatata ttccatcaca gccagcccag ctccactcag 1500
ggctggcccc gggagtcccc gtgtgcccc aagaggctagc ccagggtga gcagggccct 1560
cagaggaaa gtagtatggc ggaggccatg ggggcccctc ggcattcaca cacagcctgg 1620
cctccccctg ggagctgcat ggagcctgg ctccaggctc caggctgact ggggcctctg 1680
cctccaggag ggcacagct ttccctggct cagggatctt ctccctcccc tcaccgctg 1740
cccagccctc ccagctgatg tcaactctgc tctaagccaa ggctcagga gagcatcacc 1800
accacaccct gcggccttgc cttggggcca gactggctgc acagcccaac caggaggggt 1860
ctgcctccca cgctgggaca cagaccggcc gcatgtctgc atggcagaag cgtctccctt 1920
gccacggcct gggaggggtg ttctgtttct cagcatccac taatattcag tcctgtatat 1980
tttaataaaa taaacttgac aaaggaaaaa aaaaccg 2017

```

<210> 65  
 <211> 97  
 <212> DNA  
 <213> Homo sapiens

```

<400> 65
gtccaggaa tcctcagcag cgcctccttc agctccacag ccagacgccc tcagacagca 60
aagcctaccc ccgcgcgcgc ccctgcccgc cgctgcg 97

```

<210> 66  
 <211> 1474  
 <212> DNA  
 <213> Homo sapiens

```

<400> 66
aagtctaattg atcatattta tttatttata tgaaccatgt ctattaattt aattatttaa 60
taatatttat attaaactcc ttatgttact taacatcttc tgtaacagaa gtcagtactc 120
ctgttgcgga gaaaggagtc atacttgtga agacttttat gtcactactc taaagatttt 180
gctgttgctg ttaagtttgg aaaacagttt ttattctgtt ttataaacca gagagaaatg 240
agttttgacg tctttttact tgaatttcaa cttatattat aaggacgaaa gtaaagatgt 300

```

```

ttgaatactt aaacactatc acaagatgcc aaaatgctga aagtttttac actgtcgatg 360
tttccaatgc atcttccatg atgcattaga agtaactaat gtttgaaatt ttaaagtact 420
tttgggtatt tttctgtcat caaacaaaac aggtatcagt gcattattaa atgaatattt 480
aaattagaca ttaccagtaa tttcatgtct acttttttaa atcagcaatg aaacaataat 540
ttgaaatttc taaattcata gggtagaatc acctgtaaaa gcttgtttga tttcttaaag 600
ttattaaact tgtacatata ccaaaaagaa gctgtcttgg atttaaactc gtaaaatcag 660
atgaaatttt actacaattg cttgttaaaa tattttataa gtgatgttcc tttttcacca 720
agagtataaa ccttttttagt gtgactgtta aaacttcctt ttaaatacaa atgccaaatt 780
tattaagggtg gtggagccac tgcagtgtta tctcaaaata agaatacctt gttgagatat 840
tccagaatct gtttatatgg ctggtaacat gtaaaaacc cataaccccg ccaaaagggg 900
tcctaccctt gaacataaag caataaccaa aggagaaaag cccaaattat tggttccaaa 960
tttaggggtt aaactttttg aagcaaactt ttttttagcc ttgtgcactg cagacctggt 1020
actcagatth tgctatgagg ttaatgaagt accaagctgt gcttgaataa cgatatgttt 1080
tctcagatth tctgttgtac agtttaattt agcagtccat atcacattgc aaaagtagca 1140
atgacctcat aaaatacctc ttcaaaatgc ttaaaattcat ttcacacatt aattttatct 1200
cagtcttgaa gccaatcag taggtgcatt ggaatcaagc ctggctacct gcattgtgtt 1260
ccttttcttt tcttctttta gccattttgc taagagacac agtcttctca aacacttcgt 1320
ttctcctatt ttgttttact agttttaaga tcagagttca ctttctttgg actctgccta 1380
tattttctta cctgaacttt tgcaagtttt caggtaaacc tcagctcagg actgctatth 1440
agctcctctt aagaagatta aaaaaaaaaa aaaa 1474

```

<210> 67  
<211> 99  
<212> DNA  
<213> Homo sapiens

```

<400> 67
gcgcccggcc cccaccctc gcagcacc cgcccccgc cctcccagc cgggtccagc 60
cggagccatg gggccggagc cgcagtgag accatggag 99

```

<210> 68  
<211> 614  
<212> DNA  
<213> Homo sapiens

```

<400> 68
tgaaccagaa ggccaagtcc gcagaagccc tgatgtgtcc tcaggagca gggaaggcct 60

```

gactttctgct ggcatacaaga ggtgggaggg ccctccgacc acttccaggg gaacctgcca 120  
 tgccaggaac ctgtcctaag gaaccttctt tctgtcttga gttcccagat ggctggaagg 180  
 ggtccagcct cgttggaaga ggaacagcac tggggagtct ttgtggattc tgaggccctg 240  
 cccaatgaga ctctagggtc cagtggatgc cacagcccag cttggccctt tccttccaga 300  
 tcctgggtac tgaaagcctt agggaagctg gcctgagagg ggaagcggcc ctaaggagg 360  
 gtctaagaac aaaagcgacc cattcagaga ctgtccctga aacctagtag tgccccccat 420  
 gaggaaggaa cagcaatggt gtcagtatcc aggctttgta cagagtgtct ttctgttttag 480  
 tttttacttt ttttgttttg ttttttttaa gacgaaataa agaccaggg gagaatgggt 540  
 gttgtatggg gaggaagtg tgggggggtcc ttctccacac ccactttgtc catttgcaaa 600  
 tatatttttg aaaa 614

<210> 69  
 <211> 36  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 69  
 aaagtcgacg taatcgcgga ggcttggggc agccgg 36

<210> 70  
 <211> 30  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 70  
 tttgcgactg gtcagctgcg ggatcccaag 30

<210> 71  
 <211> 33  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 71  
 aagtcgacgt aagagctcca gagagaagtc gag 33

<210> 72  
 <211> 33  
 <212> DNA  
 <213> Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer

&lt;400&gt; 72

aaacccgggc agcaaggcaa ggctccaatg cac

33

&lt;210&gt; 73

&lt;211&gt; 39

&lt;212&gt; DNA

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer

&lt;400&gt; 73

gccgggcagg aggaaggagc ctccctcagg gtttcggga

39

&lt;210&gt; 74

&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer

&lt;400&gt; 74

ctgcactaga gacaaagacg tgatgttaat

30

&lt;210&gt; 75

&lt;211&gt; 66

&lt;212&gt; DNA

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Polylinker

&lt;400&gt; 75

gaacaaatgt cgacgggggc ccctagcaga tctagcgctg gatcccccg ggagctcaug

60

gaagac

66

&lt;210&gt; 76

&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer

&lt;400&gt; 76

cgggtgttggg cgcgttatatt atcggagttg

30

&lt;210&gt; 77

&lt;211&gt; 30

&lt;212&gt; DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: Primer

<400> 77  
ttggcgaaga atgaaaatag ggttggtact 30

<210> 78  
<211> 22  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: Primer

<400> 78  
ggtgaaggtc ggagtcaacg ga 22

<210> 79  
<211> 21  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: Primer

<400> 79  
gagggatctc gctcctggaa g 21

<210> 80  
<211> 55  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: Primer

<400> 80  
aaagtgcagc taaccgccag atttgaatcg cgggaccgct tggcagaggt ggcgg 55

<210> 81  
<211> 54  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: Primer

<400> 81  
aaaggatccg ggcaacgtcg gggcacccat gccgccgccg ccacctctgc caac 54

<210> 82  
<211> 40  
<212> DNA  
<213> Artificial

<211> 31  
 <212> DNA  
 <213> Artificial  
  
 <220>  
 <223> Description of Artificial Sequence: Primer  
  
 <400> 87  
 agcccatggt gctcactgcg gctccggccc c 31  
  
 <210> 88  
 <211> 22  
 <212> DNA  
 <213> Artificial  
  
 <220>  
 <223> Description of Artificial Sequence: Primer  
  
 <400> 88  
 agactctgaa ccagaaggcc aa 22  
  
 <210> 89  
 <211> 36  
 <212> DNA  
 <213> Artificial  
  
 <220>  
 <223> Description of Artificial Sequence: Primer  
  
 <400> 89  
 ctcggtacca gttttccaaa atatatttgc aaatgg 36  
  
 <210> 90  
 <211> 58  
 <212> DNA  
 <213> Artificial  
  
 <220>  
 <223> Description of Artificial Sequence: Primer  
  
 <400> 90  
 cccaagcttc gcgcccggcc ccccaccct cgcagcacc cgcgcccgc gccctccc 58  
  
 <210> 91  
 <211> 61  
 <212> DNA  
 <213> Artificial  
  
 <220>  
 <223> Description of Artificial Sequence: Primer  
  
 <400> 91  
 ggcccatgg ctccggctgg acccggtgg gaccggctg ggagggcgcg ggagggcgcg 60  
 g 61



<210> 92  
 <211> 7008  
 <212> DNA  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: Expression Vector

<400> 92

```

gacggatcgg gagatctccc gatcccctat ggtgcactct cagtacaatc tgctctgatg      60
ccgcatagtt aagccagtat ctgctccctg cttgtgtggt ggaggtcgct gagtagtgcg      120
cgagcaaaat ttaagctaca acaaggcaag gcttgaccga caattgcatg aagaatctgc      180
ttagggttag gcgtttttgcg ctgcttcgcg atgtacgggc cagatatacg cgttgacatt      240
gattattgac tagttattaa tagtaatcaa ttacgggggc attagttcat agcccatata      300
tgaggattccg cgttacataa cttacggtaa atggcccgcc tggctgaccg cccaacgacc      360
ccgcccatt  gacgtcaata atgacgtatg ttcccatagt aacgccaata gggactttcc      420
attgacgtca atgggtggag tatttacggt aaactgccca cttggcagta catcaagtgt      480
atcatatgcc aagtacgccc cctattgacg tcaatgacgg taaatggccc gcctggcatt      540
atgcccagta catgacctta tgggactttc ctacttggca gtacatctac gtattagtca      600
tcgctattac catggtgatg cggttttggc agtacatcaa tgggcgtgga tagcggtttg      660
actcacgggg atttccaagt ctccacccca ttgacgtcaa tgggagtttg ttttggcacc      720
aaaatcaacg ggactttcca aaatgtcgta acaactccgc ccattgacg caaatgggcg      780
gtaggcgtgt acggtgggag gtctatataa gcagagctct ctggctaact aagctttcgg      840
cgcgccgagg taccatggga tccgaagacg ccaaaaacat aaagaaaggc ccggcgccat      900
tctatcctct agaggatgga accgctggag agcaactgca taaggctatg aagagatacg      960
ccctggttcc tggaacaatt gcttttacag atgcacatat cgaggtgaac atcacgtacg     1020
cggaatactt cgaaatgtcc gttcggttgg cagaagctat gaaacgatat gggctgaata     1080
caaatcacag aatcgtcgta tgcagtgaaa actctcttca attctttatg ccggtgttgg     1140
gcgcgttatt tatcgagatt gcagttgcgc ccgcgaacga catttataat gaacgtgaat     1200
tgctcaacag tatgaacatt tcgcagccta ccgtagtggt tgtttccaaa aaggggttgc     1260
aaaaaatttt gaacgtgcaa aaaaaattac caataatcca gaaaattatt atcatggatt     1320
ctaaaacgga ttaccaggga tttcagtcga tgtacacgtt cgtcacatct catctacctc     1380
ccggttttaa tgaatacgat tttgtaccag agtcctttga tcgtgacaaa acaattgcac     1440
tgataatgaa ttcctctgga tctactgggt tacctaaggg tgtggccctt ccgcatagaa     1500
ctgcctgcgt cagattctcg catgccagag atcctatttt tggcaatcaa atcattccgg     1560

```

atactgcgat	tttaagtgtt	gttccattcc	atcacggttt	tggaatgttt	actacactcg	1620
gatattttgat	atgtggattt	cgagtcgtct	taatgtatag	atttgaagaa	gagctgtttt	1680
tacgatccct	tcaggattac	aaaattcaaa	gtgcgttgct	agtaccaacc	ctattttcat	1740
tcttcgcaa	aagcactctg	attgacaaat	acgatttatc	taatttacac	gaaattgctt	1800
ctgggggagc	acctctttcg	aaagaagtcg	gggaagcggg	tgcaaaacgc	ttccatcttc	1860
cagggatagc	acaaggatat	gggctcactg	agactacatc	agctattctg	attacacccg	1920
aggggggatga	taaaccgggc	gcggtcggta	aagttgttcc	atTTTTtgaa	gcgaagggtg	1980
tggatctgga	taccgggaaa	acgctgggag	ttaatcagag	aggcgaatta	tgtgtcagag	2040
gacctatgat	tatgtccggg	tatgtaaaca	atccggaagc	gaccaacgcc	ttgattgaca	2100
aggatggatg	gctacattct	ggagacatag	cttactggga	cgaagacgaa	cacttcttca	2160
tagttgaccg	cttgaagtct	ttaattaaat	acaaaggata	tcagggtggcc	cccgtggaat	2220
tggaatcgat	attgtttaca	caccccaaca	tcttcgacgc	gggctgggca	ggtcttcccg	2280
acgatgacgc	cgggtgaactt	cccgccgccc	ttgttgTTTT	ggagcacgga	aagacgatga	2340
cggaaaaaga	gatcgtggat	tacgtcgcca	gtcaagtaac	aaccgcgaaa	aagttgcgcg	2400
gaggagtgtg	gtttgtggac	gaagtaccga	aaggctttac	cggaaaactc	gacgcaagaa	2460
aatcagaga	gatcctcata	aaggccaaga	agggcggaag	gtccaaattg	cgcgccgctt	2520
aactcgagaa	taaaatgagg	aaattgcata	gcattgtctg	agtaggtgtc	attctattct	2580
gggggggtggg	gtggggcagg	acagcaaggg	ggaggattgg	gaagacaata	gcaggcatgc	2640
tggggatgag	gtgggctcta	tggcttctga	ggcggaaaga	accagctggg	gctctagggg	2700
gtatccccac	gcgccctgta	gcggcgcatc	aagcgcgggc	ggtgtgggtg	ttacgcgcag	2760
cgtgaccgct	acacttgcca	gcgccctagc	gcccgtctct	ttcgttttct	tcccttcctt	2820
tctcgccacg	ttcgccgggt	ttccccgtca	agctctaaat	cgggggctcc	ctttaggggt	2880
ccgatttagt	gctttacggc	acctcgaccc	caaaaaactt	gattaggggtg	atggttcacg	2940
tagtgggcca	tcgccctgat	agacggTTTT	tcgccctttg	acgttgaggt	ccacgttctt	3000
taatagtggg	ctcttgttcc	aaactggaac	aacactcaac	cctatctcgg	tctattcttt	3060
tgatttataa	gggattttgc	cgatttcggc	ctattgggtta	aaaaatgagc	tgatttaaca	3120
aaaatttaac	gcgaattaat	tctgtggaat	gtgtgtcagt	taggggtgtg	aaagtcccca	3180
ggctccccag	caggcagaag	tatgcaaagc	atgcatctca	attagtcagc	aaccagggtg	3240
ggaaagtccc	caggctcccc	agcaggcaga	agtatgcaaa	gcatgcatct	caattagtca	3300
gcaaccatag	ttccgcccc	aactccgccc	atcccgcccc	taactccgcc	cagttccgcc	3360
cattctccgc	cccatgggtg	actaattttt	tttatttatg	cagaggccga	ggccgcctct	3420

gcctctgagc	tattccagaa	gtagtgagga	ggcttttttg	gaggcctagg	cttttgcaaa	3480
aagctcccgg	gagcttgtat	atccattttc	ggatctgatc	agcacgtgat	gaaaaagcct	3540
gaactcaccg	cgacgtctgt	cgagaagttt	ctgatcgaaa	agttcgacag	cgtctccgac	3600
ctgatgcagc	tctcggaggg	cgaagaatct	cgtgctttca	gcttcgatgt	aggagggcgt	3660
ggatatgtcc	tgcgggtaaa	tagctgcgcc	gatggtttct	acaaagatcg	ttatgtttat	3720
cggcactttg	catcggccgc	gctcccgatt	ccggaagtgc	ttgacattgg	ggaattcagc	3780
gagagcctga	cctattgcat	ctcccgccgt	gcacagggtg	tcacgttgca	agacctgcct	3840
gaaaccgaac	tgcccgtctg	tctgcagccg	gtcgcggagg	ccatgggatgc	gatcgctgcg	3900
gccgatctta	gccagacgag	cgggttcggc	ccattcggac	cgcaaggaat	cggatcaatac	3960
actacatggc	gtgatttcat	atgcgcgatt	gctgatcccc	atgtgtatca	ctggcaaact	4020
gtgatggacg	acaccgtcag	tgcgtccgtc	gcgagggtc	tcgatgagct	gatgctttgg	4080
gccgaggact	gccccgaagt	ccggcacctc	gtgcacgcgg	atttcggctc	caacaatgtc	4140
ctgacggaca	atggccgcat	aacagcggtc	attgactgga	gcgaggcgat	gttcggggat	4200
tccaatacgc	aggtcgcaa	catcttcttc	tggaggccgt	ggttggtctg	tatggagcag	4260
cagacgcgct	acttcgagcg	gaggcatccg	gagcttgacg	gatcgccgcg	gctccgggcg	4320
tatatgctcc	gcattggtct	tgaccaactc	tatcagagct	tgggttgacgg	caatttcgat	4380
gatgcagctt	gggcgcaggg	tcgatgcgac	gcaatcgctc	gatccggagc	cgggactgtc	4440
gggcgtacac	aaatcgcccc	cagaagcgcg	gccgtctgga	ccgatggctg	tgtagaagta	4500
ctcgccgata	gtggaaaccg	acgccccagc	actcgtccga	gggcaaagga	atagcacgtg	4560
ctacgagatt	tcgattccac	cgcgccttc	tatgaaaggt	tgggcttcgg	aatcgttttc	4620
cgggacgcgc	gctggatgat	cctccagcgc	ggggatctca	tgctggagtt	cttcgcccac	4680
cccaacttgt	ttattgcagc	ttataatggt	tacaaataaa	gcaatagcat	cacaaatttc	4740
acaaataaag	catttttttc	actgcattct	agttgtgggt	tgtccaaact	catcaatgta	4800
tcttatcatg	tctgtatacc	gtcgacctct	agctagagct	tggcgtaatc	atggatcatg	4860
ctgtttcctg	tgtgaaattg	ttatccgctc	acaattccac	acaacatacg	agccggaagc	4920
ataaagtgtg	aagcctgggg	tgcctaatag	gtgagctaac	tcacattaat	tgcgttgccg	4980
tcactgcccc	ctttccagtc	gggaaacctg	tcgtgccagc	tgcattaatg	aatcggccaa	5040
cgcgcgggga	gaggcggttt	gcgtattggg	cgctcttcgc	cttcctcgct	cactgactcg	5100
ctgcgctcgg	tcgttcgggt	gcggcgagcg	gtatcagctc	actcaaaggc	ggtaatacgg	5160
ttatccacag	aatcagggga	taacgcagga	aagaacatgt	gagcaaaagg	ccagcaaaag	5220

gccaggaacc	gtaaaaaggc	cgcgttgctg	gcgtttttcc	ataggctccg	ccccctgac	5280
gagcatcaca	aaaatcgacg	ctcaagtcag	aggtggcgaa	acccgacagg	actataaaga	5340
taccaggcgt	ttccccctgg	aagctccctc	gtgcgctctc	ctgttccgac	cctgccgctt	5400
accggatacc	tgtccgcctt	tctcccttcg	ggaagcgtgg	cgctttctca	tagctcacgc	5460
tgtaggtatc	tcagttcggg	gtaggtcgtt	cgctccaagc	tgggctgtgt	gcacgaaccc	5520
cccgttcagc	ccgaccgctg	cgccttatcc	ggtaactatc	gtcttgagtc	caacccggtg	5580
agacacgact	tatcgccact	ggcagcagcc	actggtaaca	ggattagcag	agcgagggtat	5640
gtaggcggtg	ctacagagtt	cttgaagtgg	tggcctaact	acggctacac	tagaagaaca	5700
gtatttggtg	tctgcgctct	gctgaagcca	gttaccttcg	gaaaaagagt	tggtagctct	5760
tgatccggca	aacaaaccac	cgctggtagc	ggtttttttg	tttgcaagca	gcagattacg	5820
cgcagaaaaa	aaggatctca	agaagatcct	ttgatctttt	ctacgggggtc	tgacgctcag	5880
tggaacgaaa	actcacgtta	agggattttg	gtcatgagat	tatcaaaaag	gatcttcacc	5940
tagatccttt	taaattaaaa	atgaagtttt	aatcaatct	aaagtatata	tgagtaaact	6000
tggtctgaca	gttaccaatg	cttaatcagt	gaggcaccta	tctcagcgat	ctgtctattt	6060
cgttcatcca	tagttgcctg	actccccgtc	gtgtagataa	ctacgatacg	ggagggctta	6120
ccatctggcc	ccagtgcctg	aatgataccg	cgagaccac	gctcaccggc	tccagattta	6180
tcagcaataa	accagccagc	cggaagggcc	gagcgcagaa	gtggctcctgc	aactttatcc	6240
gcctccatcc	agtctattaa	ttgttgccgg	gaagctagag	taagtagttc	gccagttaat	6300
agtttgcgca	acgttggttg	cattgctaca	ggcatcgtgg	tgtcacgctc	gtcgtttggt	6360
atggcttcac	tcagctccgg	ttcccaacga	tcaaggcgag	ttacatgata	ccccatgttg	6420
tgcaaaaaag	cggttagctc	cttcggctct	ccgatcgttg	tcagaagtaa	gttggccgca	6480
gtgttatcac	tcattggttat	ggcagcactg	cataattctc	ttactgtcat	gccatccgta	6540
agatgctttt	ctgtgactgg	tgagtactca	accaagtcac	tctgagaata	gtgtatgcgg	6600
cgaccgagtt	gctcttgccc	ggcgtaata	cgggataata	ccgcgccaca	tagcagaact	6660
ttaaaagtgc	tcattcattgg	aaaacgttct	tcggggcgaa	aactctcaag	gatcttaccg	6720
ctgttgagat	ccagttcgat	gtaaccact	cgtgcacca	actgatcttc	agcatctttt	6780
actttcacca	gcgtttctgg	gtgagcaaaa	acaggaaggc	aaaatgccgc	aaaaaaggga	6840
ataagggcga	cacggaaatg	ttgaatactc	atactcttcc	tttttcaata	ttattgaagc	6900
atztatcagg	gttattgtct	catgagcgga	tacatatttg	aatgtattta	gaaaaataaa	6960
caaatagggg	ttccgcgcac	atttccccga	aaagtgccac	ctgacgtc		7008

<210> 93  
 <211> 11693  
 <212> DNA  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: Expression Vector

<400> 93

```

gttgacattg attattgact agttattaat agtaatcaat tacgggggtca ttagttcata      60
gcccatatat ggagttccgc gttacataac ttacggtaaa tggcccgccct ggctgaccgc      120
ccaacgaccc ccgccattg acgtcaataa tgacgtatgt tcccatagta acgccaatag      180
ggactttcca ttgacgtcaa tgggtggagt atttacggta aactgcccac ttggcagtac      240
atcaagtgtg tcatatgcca agtccgcccc ctattgacgt caatgacggt aaatggccccg      300
cctggcatta tgcccagtag atgaccttac gggactttcc tacttggcag tacatctacg      360
tattagtcac cgctattacc atggtgatgc ggttttggca gtacaccaat gggcgtggat      420
agcggtttga ctacggggga tttccaagtc tccaccccat tgacgtcaat gggagtttgt      480
tttggcacca aaatcaacgg gactttccaa aatgtcgtaa taaccccgcc ccgttgacgc      540
aaatgggccc taggcgtgta cgggtggagg tctatataag cagagctcgt ttagtgaacc      600
gtaagctttc ggcgcgccac ggtaccatgg gatccgaaga cgccaaaaaac ataaagaaag      660
gcccgggccc attctatcct ctagaggatg gaaccgctgg agagcaactg cataaggcta      720
tgaagagata cgccctgggt cctggaacaa ttgcttttac agatgcacat atcgagggtga      780
acatcacgta cgcggaatac ttcgaaatgt ccgttcgggt ggcagaagct atgaaacgat      840
atgggctgaa taaaaatcac agaatcgtcg tatgcagtga aaactctctt caattcttta      900
tgccggtggt gggcgcggtt tttatcggag ttgcagttgc gcccggaac gacatttata      960
atgaacgtga attgctcaac agtatgaaca tttcgcagcc taccgtagtg tttgtttcca     1020
aaaagggggt gcaaaaaatt ttgaacgtgc aaaaaaatt accaataatc cagaaaatta     1080
ttatcatgga ttctaaaacg gattaccagg gatttcagtc gatgtacacg ttcgtcacat     1140
ctcatctacc tcccggtttt aatgaatacg attttgtacc agagtccttt gatcgtgaca     1200
aaacaattgc actgataatg aattcctctg gatctactgg gttacctaag ggtgtggccc     1260
ttccgcatag aactgcctgc gtcagattct cgcagccag agatcctatt tttggcaatc     1320
aaatcattcc ggatactgcg attttaagtg ttgttcatt ccatcacggt tttggaatgt     1380
ttactacact cggatatttg atatgtggat ttcgagtcgt cttaatgtat agatttgaag     1440
aagagctggt ttacgatcc cttcaggatt acaaaattca aagtgcgttg ctagtaccaa     1500
ccctattttc attcttcgcc aaaagcactc tgattgacaa atacgattta tctaatttac     1560

```

acgaaattgc ttctgggggc gcacctcttt cgaaagaagt cggggaagcg gttgcaaaac 1620  
 gcttccatct tccagggata cgacaaggat atggggtcac tgagactaca tcagctattc 1680  
 tgattacacc cgagggggat gataaaccgg gcgcggtcgg taaagttgtt ccattttttg 1740  
 aagcgaaggt tgtggatctg gataccggga aaacgctggg cgttaatcag agaggcgaat 1800  
 tatgtgtcag aggacctatg attatgtccg gttatgtaaa caatccggaa gcgaccaacg 1860  
 ccttgattga caaggatgga tggctacatt ctggagacat agcttactgg gacgaagacg 1920  
 aacacttctt catagttgac cgcttgaagt ctttaattaa atacaaagga tatcaggtgg 1980  
 cccccgctga attggaatcg atattgttac aacaccccaa catcttcgac gcgggcgtgg 2040  
 caggtcttcc cgacgatgac gccggtgaac ttcccgcgcg cgttgttgtt ttggagcacg 2100  
 gaaagacgat gacggaaaaa gagatcgtgg attacgtcgc cagtcaagta acaaccgcga 2160  
 aaaagttgcg cggaggaggt gtgtttgtgg acgaagtacc gaaaggtctt accggaaaaac 2220  
 tcgacgcaag aaaaatcaga gagatcctca taaaggccaa gaagggcgga aagtccaaat 2280  
 tgcgcggcgc ctaactcgag aataaacaag ttaacaacaa caattgcatt cattttatgt 2340  
 ttcaggttca gggggaggtg tgggaggttt tttaaagcaa gtaaaacctc taaaaatgtg 2400  
 gtatggctga ttatgatccg gctgcctcgc gcgtttcggg gatgacgggtg aaaacctctg 2460  
 acacatgcag ctcccggaga cggtcacagc ttgtctgtaa gcggatgccg ggagcagaca 2520  
 agcccgtcag gcgtcagcgg gtgttgccgg gtgtcggggc gcagccatga ggtcgactct 2580  
 agaggatcga tgcccgcgcc cggacgaact aaacctgact acgacatctc tgccccttct 2640  
 tcgcggggca gtgcatgtaa tcccttcagt tgggtggtac aacttgccaa ctggggccctg 2700  
 ttccacatgt gacacggggg gggaccaaac acaaaggggt tctctgactg tagttgacat 2760  
 ccttataaat ggatgtgcac atttgccaac actgagtggc tttcatcctg gagcagactt 2820  
 tgcagtctgt ggactgcaac acaacattgc ctttatgtgt aactcttggc tgaagctctt 2880  
 acaccaatgc tgggggacat gtacctcca ggggccagg aagactacgg gaggctacac 2940  
 caacgtcaat cagaggggcc tgtgtagcta ccgataagcg gaccctcaag agggcattag 3000  
 caatagtgtt tataaggccc ccttggttaac cctaaacggg tagcatatgc ttcccgggta 3060  
 gtagtatata ctatccagac taaccctaata tcaatagcat atgttaccca acgggaagca 3120  
 tatgctatcg aattaggggt agtaaaaggg tcctaaggaa cagcgatatc tcccaccca 3180  
 tgagctgtca cggttttatt tacatggggg caggattcca cgagggtagt gaaccatttt 3240  
 agtcacaagg gcagtggctg aagatcaagg agcgggcagt gaactctcct gaatcttcgc 3300  
 ctgcttcttc attctccttc gtttagctaa tagaataact gctgagttgt gaacagtaag 3360  
 gtgtatgtga ggtgctcgaa aacaagggtt caggtgacgc cccagaata aaatttgac 3420

gggggggttca gtggtggcat tgtgctatga caccaatata accctcacia accccttggg 3480  
 caataaatac tagtgtagga atgaaacatt ctgaatatct ttaacaatag aaatccatgg 3540  
 ggtggggaca agccgtaaag actggatgtc catctcacac gaatttatgg ctatgggcaa 3600  
 cacataatcc tagtgcaata tgatactggg gttattaaga tgtgtcccag gcagggacca 3660  
 agacaggtga accatgttgt tacactctat ttgtaacaag gggaaagaga gtggacgccg 3720  
 acagcagcgg actccactgg ttgtctctaa ccccccgaa aattaaacgg ggctccacgc 3780  
 caatggggcc cataaaciaa gacaagtggc cactcttttt tttgaaattg tggagtgggg 3840  
 gcacgcgtca gccccacac gccgcctgc ggttttggac tgtaaaataa ggggtgtaata 3900  
 acttggctga ttgtaacccc gctaaccact gcggtcaaac cacttgcca caaaaccact 3960  
 aatggcacc cggggaatac ctgcataagt aggtgggcgg gccaaagatag gggcgcgatt 4020  
 gctgcgatct ggaggacaaa ttacacacac ttgcgcctga gcgccaagca cagggttgtt 4080  
 ggtcctcata ttcacgaggt cgctgagagc acggtgggct aatgttgcca tgggtagcat 4140  
 atactacca aatatctgga tagcatatgc taccctaata tatatctggg tagcataggc 4200  
 taccctaata tatatctggg tagcatatgc taccctaata tatatctggg tagtatatgc 4260  
 taccctaatt tatatctggg tagcataggc taccctaata tatatctggg tagcatatgc 4320  
 taccctaata tatatctggg tagtatatgc taccctaata tgtatccggg tagcatatgc 4380  
 taccctaata gagattaggg tagtatatgc taccctaatt tatatctggg tagcatatac 4440  
 tacccaaata tctggatagc atatgctatc ctaatctata tctgggtagc atatgctatc 4500  
 ctaatctata tctgggtagc ataggctatc ctaatctata tctgggtagc atatgctatc 4560  
 ctaatctata tctgggtagt atatgctatc ctaatttata tctgggtagc ataggctatc 4620  
 ctaatctata tctgggtagc atatgctatc ctaatctata tctgggtagt atatgctatc 4680  
 ctaatctgta tccgggtagc atatgctatc ctcatgcata tacagtcagc atatgatacc 4740  
 cagtagtaga gtgggagtgct taccctttgc atatgccgcc acccccag ggggcgtgaa 4800  
 ttttcgctgc ttgtcctttt cctgctgggt gctcccatc ttaggtgaat ttaaggaggc 4860  
 caggctaaag ccgtcgcatt tctgattgct caccaggtaa atgtcgctaa tgttttccaa 4920  
 cgcgagaagg tggtgagcgc ggagctgagt gacgtgacaa catgggtatg cccaattgcc 4980  
 ccatgttggg aggacgaaaa tggtgacaag acagatggcc agaaatacac caacagcacg 5040  
 catgatgtct actggggatt tattcttttag tgcgggggaa tacacggctt ttaatacgat 5100  
 tgagggcgct tcctaacaag ttacatcact cctgcccttc ctacccctca tctccatcac 5160  
 ctcttcatc tccgtcatct ccgtcatcac cctccgcggc agcccttcc accatagggtg 5220

gaaaccaggg	aggcaaattct	actccatcgt	caaagctgca	cacagtcacc	ctgatattgc	5280
aggtaggagc	gggctttgtc	ataacaaggt	ccttaatcgc	atccttcaaa	acctcagcaa	5340
atatatgagt	ttgtaaaaag	accatgaaat	aacagacaat	ggactccctt	agcgggccag	5400
gttgtgggcc	gggtccaggg	gccattccaa	aggggagacg	actcaatggg	gtaagacgac	5460
atttgtggaat	agcaagggca	gttcctcgcc	ttaggttgta	aaggagggtc	ttactacctc	5520
catatacgaa	cacaccggcg	acccaagttc	cttcgtcggg	agtcctttct	acgtgactcc	5580
tagccaggag	agctcttaaa	ccttctgcaa	tgttctcaaa	tttcgggttg	gaacctcctt	5640
gaccacgatg	cttttccaaa	ccacctcctt	tttttgcgcc	ctgcctccat	cacctgacc	5700
ccgggggtcca	gtgcttgggc	cttctcctgg	gtcatctgcg	gggccctgct	ctatcgctcc	5760
cggggggcacg	tcaggctcac	catctgggcc	accttcttgg	tggtattcaa	aataatcggc	5820
ttcccctaca	gggtggaaaa	atggccttct	acctggaggg	ggcctgcgcg	gtggagaccc	5880
ggatgatgat	gactgactac	tgggactcct	gggcctcttt	tctccacgtc	cacgacctct	5940
ccccctggct	ctttcacgac	ttccccccct	ggctctttca	cgtcctctac	cccggcggcc	6000
tccactacct	cctcgacccc	ggcctccact	acctcctcga	ccccggcctc	cactgcctcc	6060
tcgaccccg	cctccacctc	ctgetcctgc	ccctcctgct	cctgcccctc	ctcctgctcc	6120
tgcccctcct	gcccctcctg	ctcctgcccc	tcttgcccct	cctgctcctg	ccccctcctg	6180
ccctcctgct	cctgcccctc	ctgcccctcc	tcttgctcct	gcccctcctg	ccccctcctcc	6240
tgctcctgcc	cctcctgccc	ctcctgctcc	tgcccctcct	gcccctcctg	ctcctgcccc	6300
tcttgcccct	cctgctcctg	ccccctcctg	tcttgcccct	cctgctcctg	ccccctcctg	6360
tcttgcccct	cctgcccctc	ctgcccctcc	tcttgctcct	gcccctcctg	ctcctgcccc	6420
tcttgcccct	cctgcccctc	ctgctcctgc	ccctcctcct	gctcctgccc	ctcctgcccc	6480
tcttgcccct	cctcctgctc	ctgcccctcc	tgcccctcct	cctgctcctg	ccccctcctcc	6540
tgctcctgcc	cctcctgccc	ctcctgcccc	tctcctgct	cctgcccctc	ctgcccctcc	6600
tcttgctcct	gcccctcctc	ctgetcctgc	ccctcctgcc	cctcctgccc	ctcctcctgc	6660
tcttgcccct	cctcctgctc	ctgcccctcc	tgcccctcct	gcccctcctg	ccccctcctcc	6720
tgctcctgcc	cctcctcctg	ctcctgcccc	tcttgctcct	gcccctcccg	ctcctgctcc	6780
tgctcctggt	ccaccgtggg	tccctttgca	gccaatgcaa	cttgagcgtt	tttgggggtct	6840
ccggacacca	tctctatgtc	ttggccctga	tctgagccg	cccggggctc	ctggtcttcc	6900
gctcctcgt	cctcgtcctc	ttccccgtcc	tgtccatgg	ttatcacccc	ctcttctttg	6960
aggtccactg	ccgccggagc	cttctgggtcc	agatgtgtct	cccttctctc	ctaggccatt	7020
tccaggtcct	gtacctggcc	cctcgtcaga	catgattcac	actaaaagag	atcaatagac	7080



atcttttatta gacgacgctc agtgaataca gggagtgcag actcctgccc cctccaacag	7140
cccccccacc ctcatcccct tcatggctgc tgtcagacag atccaggtct gaaaattccc	7200
catcctccga accatcctcg tctcatcac caattactcg cagcccgga aactcccgt	7260
gaacatcctc aagatttgcg tcttgagcct caagccaggc ctcaaattcc tcgtccccct	7320
ttttgctgga cggtagggat ggggattctc gggaccctc ctcttctct tcaaggctac	7380
cagacagaga tgctactggg gcaacggaag aaaagctggg tgcggcctgt gaggatcagc	7440
ttatcgatga taagctgtca aacatgagaa ttcttgaaga cgaaagggcc tcgtgatacg	7500
cctattttta taggttaatg tcatgataat aatggtttct tagacgtcag gtggcacttt	7560
tcggggaaat gtgcgcggaa cccctatttg tttatttttc taaatacatt caaatatgta	7620
tccgctcatg agacaataac cctgataaat gcttcaataa tattgaaaaa ggaagagtat	7680
gagtattcaa catttccgtg tcgcccttat tccctttttt gcggcatttt gccttctgt	7740
ttttgctcac ccagaaacgc tggtgaaagt aaaagatgct gaagatcagt tgggtgcacg	7800
agtgggttac atcgaactgg atctcaacag cggtaagatc cttgagagtt ttcgccccga	7860
agaacgtttt ccaatgatga gcacttttaa agttctgcta tgtggcgcggt tattatcccg	7920
tgttgacgcc gggcaagagc aactcggctc ccgcatacac tattctcaga atgacttggt	7980
tgagtactca ccagtcacag aaaagcatct tacggatggc atgacagtaa gagaattatg	8040
cagtgtgcc ataaccatga gtgataacac tgcggccaac ttacttctga caacgatcgg	8100
aggaccgaag gagctaaccg cttttttgca caacatgggg gatcatgtaa ctcgccttga	8160
tcgttgggaa ccggagctga atgaagccat accaaacgac gagcgtgaca ccacgatgcc	8220
tgcagcaatg gcaacaacgt tgcgcaaact attaaactggc gaactactta ctctagcttc	8280
ccggcaacaa ttaatagact ggatggaggc ggataaagtt gcaggaccac ttctgcgctc	8340
ggcccttccg gctggctggg ttattgctga taaatctgga gccggtgagc gtgggtctcg	8400
cggtatcatt gcagcactgg ggccagatgg taagccctcc cgtatcgtag ttatctacac	8460
gacggggagt caggcaacta tggatgaacg aaatagacag atcgtgaga taggtgcctc	8520
actgattaag cattggtaac tgtcagacca agtttactca tatatacttt agattgattt	8580
aaaacttcat ttttaattta aaaggatcta ggtgaagatc ctttttgata atctcatgac	8640
caaaatccct taacgtgagt tttcgttcca ctgagcgtca gacccgtag aaaagatcaa	8700
aggatcttct tgagatcctt tttttctgcg cgtaatctgc tgcttgcaaa caaaaaaacc	8760
accgctacca gcggtggttt gtttgccgga tcaagagcta ccaactcttt ttccgaaggt	8820
aactggcttc agcagagcgc agataccaaa tactgtcctt ctagtgtagc cgtagttagg	8880

ccaccacttc	aagaactctg	tagcaccgcc	tacatacctc	gctctgctaa	tcctgttacc	8940
agtggctgct	gccagtggcg	ataagtcgtg	tcttaccggg	ttggactcaa	gacgatagtt	9000
accggataag	gcgcagcggg	cgggctgaac	gggggggttcg	tgcacacagc	ccagcttgga	9060
gcgaacgacc	tacaccgaac	tgagatacct	acagcgtgag	ctatgagaaa	gcgccacgct	9120
tcccgaaggg	agaaaggcgg	acaggatatcc	ggtaagcggc	agggtcggaa	caggagagcg	9180
cacgagggag	cttccagggg	gaaacgcctg	gtatctttat	agtctctgtcg	ggtttcgccca	9240
cctctgactt	gagcgtcgat	ttttgtgatg	ctcgtcaggg	gggcggagacc	tatggaaaaa	9300
cgccagcaac	gcggcctttt	tacggttcct	ggccttttgc	tggccttgaa	gctgtccctg	9360
atggtcgtca	tctacctgcc	tggacagcat	ggcctgcaac	gcgggcatcc	cgatgccgcc	9420
ggaagcgaga	agaatcataa	tggggaaggc	catccagcct	cgcgtcgoga	acgccagcaa	9480
gacgtagccc	agcgcgtcgg	ccccgagatg	cgccgcgtgc	ggctgctgga	gatggcggac	9540
gcgatggata	tgttctgcca	agggttgggt	tgcgcattca	cagttctccg	caagaattga	9600
ttggctccaa	ttcttgaggt	ggtgaatccg	ttagcgaggt	gccgccctgc	ttcatccccg	9660
tggcccgttg	ctcgcgtttg	ctggcgggtg	ccccggaaga	aatatatattg	catgtcttta	9720
gttctatgat	gacacaaacc	cgcgccagcg	tcttgtcatt	ggcgaattcg	aacacgcaga	9780
tgcagtcggg	gcggcgcggt	ccgaggtcca	cttcgcatat	taagggtgacg	cgtgtggcct	9840
cgaacaccga	gcgaccctgc	agcgacccgc	ttaacagcgt	caacagcgtg	ccgcagatcc	9900
cggggggcaa	tgagatatga	aaaagcctga	actcaccgcg	acgtctgtcg	agaagtttct	9960
gatcgaaaag	ttcgacagcg	tctccgacct	gatgcagctc	tgggagggcg	aagaatctcg	10020
tgctttcagc	ttcgatgtag	gagggcgtgg	atatgtcctg	cgggtaaata	gctgcgccga	10080
tggtttctac	aaagatcggt	atgtttatcg	gcactttgca	tgggcgcgc	tcccgattcc	10140
ggaagtgctt	gacattgggg	aattcagcga	gagcctgacc	tattgcatct	cccgccgtgc	10200
acaggggtgc	acgttgcaag	acctgcctga	aaccgaactg	cccgctgttc	tgcagccggg	10260
cgcggaggcc	atggatgcga	tcgctgcggc	cgatcttagc	cagacgagcg	ggttcggccc	10320
attcggaccg	caaggaatcg	gtcaatacac	tacatggcgt	gatttcatat	gcgcgattgc	10380
tgatccccat	gtgtatcact	ggcaaactgt	gatggacgac	accgtcagtg	cgtccgtcgc	10440
gcaggctctc	gatgagctga	tgctttgggc	cgaggactgc	cccgaagtcc	ggcacctcgt	10500
gcacgcggat	ttcggctcca	acaatgtcct	gacggacaat	ggccgcataa	cagcgggtcat	10560
tgactggagc	gaggcgatgt	tcggggattc	ccaatacgag	gtcgccaaca	tcttcttctg	10620
gaggccgtgg	ttggcttgta	tggagcagca	gacgcgctac	ttcgagcgga	ggcatccgga	10680
gcttgcagga	tcgccgcggc	tccgggcgta	tatgctccgc	attggtcttg	accaactcta	10740

```

tcagagcttg gttgacggca atttcgatga tgcagcttgg gcgcagggtc gatgcgacgc 10800
aatcgtccga tccggagccg ggactgtcgg gcgtacacaa atcgcccgcga gaagcgcggc 10860
cgtctggacc gatggctgtg tagaagtact cgccgatagt ggaaaccgac gccccagcac 10920
tcgtccggat cgggagatgg gggaggctaa ctgaaacacg gaaggagaca ataccggaag 10980
gaaccgcgc tatgacggca ataaaaagac agaataaaac gcacgggtgt tgggtcgttt 11040
gttcataaac gcgggggttc gtcccagggc tggcactctg tcgatacccc accgagaccc 11100
cattggggcc aatacgcccg cgtttcttcc ttttccccac cccaccccc aagttcgggt 11160
gaaggcccag ggctcgacg caacgtcggg gcggcaggcc ctgccatagc cactggcccc 11220
gtgggttagg gacgggggtcc cccatgggga atgggttatg gttcgtgggg gttattattt 11280
gggcgttgcg tggggtcagg tccacgactg gactgagcag acagacccat ggtttttgga 11340
tggcctgggc atggaccgca tgtactggcg cgacacgaac accgggcgct tgtggctgcc 11400
aaacaccccc gacccccaaa aaccaccgcg cggatttctg gcgtgccaag ctagtcgacc 11460
aattctcatg tttgacagct tatcatcgca gatccgggca acgttggtgc cattgctgca 11520
ggcgcagaac tggtaggtat ggaagatcta tacattgaat caatattggc aattagccat 11580
attagtcatt gggtatatag cataaatcaa tattggctat tggccattgc atacgttgta 11640
tctatatcat aatatgtaca tttatattgg ctcatgtcca atatgaccgc cat 11693

```

```

<210> 94
<211> 4825
<212> DNA
<213> Artificial

```

```

<220>
<223> Description of Artificial Sequence: Expression vector

```

```

<400> 94
gacggatcgg gagatctccc gatcccctat ggtgcactct cagtacaatc tgctctgatg 60
ccgcatagtt aagccagtat ctgctccctg cttgtgtgtt ggaggtcgct gagtagtgcg 120
cgagcaaaat ttaagctaca acaaggcaag gcttgaccga caattgcatg aagaatctgc 180
ttagggttag gcgtttttgcg ctgcttcgcg atgtacgggc cagatatacg cgttgacatt 240
gattattgac tagttattaa tagtaatcaa ttacggggtc attagttcat agcccatata 300
tggagtcccg cgttacataa cttacggtaa atggcccgc tggctgaccg cccaacgacc 360
cccgccatt gacgtcaata atgacgtatg ttcccatagt aacgccaata gggactttcc 420
attgacgtca atgggtggag tattttacgg aaactgcccc cttggcagta catcaagtgt 480
atcatatgcc aagtacgcc cctattgacg tcaatgacgg taaatggccc gcctggcatt 540

```

atgcccagta	catgacctta	tgggactttc	ctacttggca	gtacatctac	gtattagtca	600
tcgctattac	catggtgatg	cggttttggc	agtacatcaa	tgggcgtgga	tagcggtttg	660
actcacgggg	atttccaagt	ctccacccca	ttgacgtcaa	tgggagtttg	ttttggcacc	720
aaaatcaacg	ggactttcca	aaatgtcgta	acaactccgc	cccattgacg	caaattgggcg	780
gtaggcgtgt	acggtggggag	gtctatataa	gcagagctct	ctggctaact	aagctttcgg	840
cgcgccgagg	taccatggga	tccgaagacg	caaaaaacat	aaagaaaggc	cgggcgccat	900
tctatcctct	agaggatgga	accgctggag	agcaactgca	taaggctatg	aagagatacg	960
ccctggttcc	tggaacaatt	gcttttacag	atgcacatat	cgaggatgaac	atcacgtacg	1020
cggaataactt	cgaaatgtcc	gttcggttgg	cagaagctat	gaaacgatat	gggctgaata	1080
caaatcacag	aatcgctcgta	tgcagtgaag	actctcttca	attcttttatg	cgggtgttgg	1140
gcgcgttatt	tatcggagtt	gcagttgcgc	ccgcgaacga	catttataat	gaacgtgaat	1200
tgctcaacag	tatgaacatt	tgcgagccta	ccgtagtgtt	tgtttccaaa	aaggggttgc	1260
aaaaaatttt	gaacgtgcaa	aaaaaattac	caataatcca	gaaaattatt	atcatggatt	1320
ctaaaacgga	ttaccaggga	tttcagtcga	tgtacacgtt	cgtcacatct	catctaccto	1380
cgggttttaa	tgaatacgat	tttgtaccag	agtcctttga	tcgtgacaaa	acaattgcac	1440
tgataatgaa	ttcctctgga	tctactgggt	tacctaaggg	tgtggccctt	ccgcatagaa	1500
ctgcctgcgt	cagattctcg	catgccagag	atcctatttt	tggcaatcaa	atcattccgg	1560
atactgcgat	tttaagtgtt	gttcatttcc	atcacggttt	tggaatgttt	actacactcg	1620
gatatttgat	atgtggattt	cgagtcgtct	taatgtatag	atttgaagaa	gagctgtttt	1680
tacgatccct	tcaggattac	aaaattcaaa	gtgcgttgct	agtaccaacc	ctattttcat	1740
tcttcgcaa	aagcactctg	attgacaaat	acgatttato	taattttacac	gaaattgctt	1800
ctggggggcg	acctctttcg	aaagaagtcg	gggaagcggg	tgcaaaacgc	ttccatcttc	1860
cagggatagc	acaaggatat	gggctcactg	agactacatc	agctattctg	attacacccg	1920
agggggatga	taaaccgggc	gcggtcggtg	aagttgttcc	attttttgaa	gcgaagggtg	1980
tggatctgga	taccgggaaa	acgctggggc	ttaatcagag	aggcgaatta	tgtgtcagag	2040
gacctatgat	tatgtccggt	tatgtaaaca	atccggaagc	gaccaacgcc	ttgattgaca	2100
aggatggatg	gctacattct	ggagacatag	cttactggga	cgaagacgaa	cacttcttca	2160
tagttgaccg	cttgaagtct	ttaattaaat	acaaaggata	tcagggtggcc	cccgtgaat	2220
tggaatcgat	attgtttaca	caccccaaca	tcttcgacgc	gggcgtggca	ggtcttcccg	2280
acgatgacgc	cgggtgaactt	cccgcgcgcg	ttgttgtttt	ggagcacgga	aagacgatga	2340
cggaaaaaga	gatcgtggat	tacgtcgcca	gtcaagtaac	aaccgcgaaa	aagttgcgcg	2400

gaggagttgt	gtttgtggac	gaagtaccga	aaggtcttac	cgaaaaactc	gacgcaagaa	2460
aaatcagaga	gatcctcata	aaggccaaga	agggcgga	gtccaaattg	cgcgcccgct	2520
aactcgagaa	taaaatgagg	aaattgcac	gcattgtctg	agtaggtgtc	attctattct	2580
ggggggtggg	gtggggcagg	acagcaaggg	ggaggattgg	gaagacaata	gcaggcatgc	2640
tggggatgcg	gtgggctcta	tggcttctga	ggcggaaga	accagctggg	gctctagggg	2700
gtatccccac	gcgccctgta	gcggcgcat	aagcgcgcg	ggtgtgggtg	ttacgcgcag	2760
cgtgaccgct	acacttgcca	gcgccctagc	gcccgcctct	ttcgctttct	tcccttccct	2820
tctcgccacg	ttcgccggct	ttccccgtca	agctctaaat	cggggggtccc	tttaggggtc	2880
cgatttagtg	ctttacggca	cctcgacccc	aaaaaacttg	attaggggtga	tggttcacgt	2940
acctagaagt	tcctattccg	aagttcctat	tctctagaaa	gtataggaac	ttccttggcc	3000
aaaaagcctg	aactcaccgc	gacgtctgtc	gagaagtttc	tgatcgaaaa	gttcgacagc	3060
gtctccgacc	tgatgcagct	ctcggagggc	gaagaatctc	gtgctttcag	cttcgatgta	3120
ggagggcgctg	gatatgtcct	gcgggtaaat	agctgcgccg	atggtttcta	caaagatcgt	3180
tatgtttatc	ggcactttgc	atcgcccgcg	ctcccgattc	cggaagtgtc	tgacattggg	3240
gaattcagcg	agagcctgac	ctattgcac	tcccgccgtg	cacagggtgt	cacgttgcaa	3300
gacctgcctg	aaaccgaact	gcccgcgtgt	ctgcagccgg	tcgcggaggc	catggatgcg	3360
atcgctgcgg	ccgatcttag	ccagacgagc	gggttcggcc	cattcggacc	gcaaggaatc	3420
ggtcaataca	ctacatggcg	tgatttcata	tgcgcgattg	ctgatcccca	tgtgtatcac	3480
tggcaaaactg	tgatggacga	caccgtcagt	gcgtccgtcg	cgagggtctc	cgatgagctg	3540
atgctttggg	ccgaggactg	ccccgaagtc	cggcacctcg	tgacgcaaac	aaaccaccgc	3600
tggtagcggg	ttttttgttt	gcaagcagca	gattacgcgc	agaaaaaag	gatctcaaga	3660
agatcctttg	atcttttcta	cggggtctga	cgctcagtgg	aacgaaaact	cacgttaagg	3720
gattttgggtc	atgagattat	caaaaaggat	cttcacctag	atccttttaa	attaaaaatg	3780
aagttttaaa	tcaatctaaa	gtatatatga	gtaaacttgg	tctgacagtt	accaatgctt	3840
aatcagtgag	gcacctatct	cagcgatctg	tctatttcgt	tcacccatag	ttgcctgact	3900
ccccgtcgtg	tagataacta	cgatacggga	gggcttacca	tctggcccca	gtgctgcaat	3960
gataccgcga	gacccacgct	caccggctcc	agatttatca	gcaataaacc	agccagccgg	4020
aagggccgag	cgcagaagtg	gtcctgcaac	tttatccgcc	tccatccagt	ctattaattg	4080
ttgccgggaa	gctagagtaa	gtagtccgcc	agttaatagt	ttgcgcaacg	ttgttgccat	4140
tgctacaggc	atcgtgggtg	cacgctcgtc	gtttgggatg	gcttcattca	gctccgggtc	4200

ccaacgatca aggcgagtta catgatcccc catgttgtgc aaaaaagcgg ttagctcctt	4260
cggtcctccg atcgttgtca gaagtaagtt ggccgcagtg ttatcactca tggttatggc	4320
agcactgcat aattctctta ctgtcatgcc atccgtaaga tgcttttctg tgactgggtga	4380
gtactcaacc aagtcattct gagaatagtg tatgcccga ccgagttgct cttgcccggc	4440
gtcaatacgg gataatacgg cgccacatag cagaacttta aaagtgtca tcattggaaa	4500
acgttcttcg gggcgaaaac tctcaaggat cttaccgctg ttgagatcca gttcgatgta	4560
acccactcgt gcacccaact gatcttcagc atcttttact ttcaccagcg tttctgggtg	4620
agcaaaaaca ggaaggcaaa atgccgcaaa aaagggaata agggcgacac ggaaatggtg	4680
aatactcata ctcttccttt ttcaatatta ttgaagcatt tatcagggtt attgtctcat	4740
gagcggatac atatttgaat gtatttagaa aaataaacia ataggggttc cgcgcacatt	4800
tccccgaaaa gtgccacctg acgtc	4825